

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 1

December 30, 1976

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.),
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family LUCANIDAE クワガタムシ科

By

Yoshihiko KUROSAWA

Subfamily CERUCHINAE ツヤハダクワガタ亜科

Genus *Ceruchus* MACLEAY, 1819

Ceruchus MACLEAY, 1819, Horae ent., 1, p. 115 (type-species: *Lucanus tenebroides* FABRICIUS, 1792).

Platycerus: LATREILLE, 1807, Gen. Crust. et Ins., 2, p. 113 (nec GEOFFROY, 1762).

Tarandus DEJEAN, 1837, Cat. ed. 3, p. 194.

C. lignarius LEWIS, 1883 ツヤハダクワガタ

LEWIS, 1883, Trans. ent. Soc. London, 1883, p. 339, pl. 14, fig. 4 (Sapporo).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Subfamily AESALINAE マダラクワガタ亜科

Tribe AESALINI マダラクワガタ族

Genus *Aesalus* FABRICIUS, 1801 マダラクワガタ属

Aesalus FABRICIUS, 1801, Syst. Eleuth., 2, p. 154 (type-species: *Lucanus scarabaeoides* PANZER, 1794).

A. asiaticus LEWIS, 1883 マダラクワガタ

LEWIS, 1883, Trans. ent. Soc. London, 1883, p. 340, pl. 14, fig. 5 (Miyanoshiba).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Mikura-jima, Tsushima).

Subfamily FIGULINAE チビクワガタ亜科

Tribe FIGULINI チビクワガタ族

Genus *Figulus* MACLEAY, 1819 チビクワガタ属

Figulus MACLEAY, 1819, Horae ent., 1, p. 109 (type-species: *Lucanus striatus* OLIVIER, 1789).

Platycerus: BOISDUVAL, 1835, Voy. de l'Astrolabe, Coléopt., 2, p. 239 (nec GEOFFROY, 1762).

Eudora CASTELNAU, 1840, Hist. Nat. Ins. Coléopt., 2, p. 174 (pars).

F. binodulus WATERHOUSE, 1873 チビクワガタ

WATERHOUSE, 1873, Ent. mon. Mag., 9, p. 277 (S. Japan).

Distr.: Japan (Honshu, Shikoku, Kyushu, Izu Is.), China, Formosa, Tonkin.

F. boninensis NAKANE et Y. KUROSAWA, 1953 オガサワラチビクワガタ

NAKANE & Y. KUROSAWA, 1953, Bull. Natn. Sci. Mus., Tokyo, (33), p. 110, fig. (Chichi-jima).

Distr.: Bonin Is. (Chichi-jima, Haha-jima).

F. punctatus WATERHOUSE, 1873 マメクワガタ

WATERHOUSE, 1873, Ent. mon. Mag., 9, p. 278 (S. Japan).

Distr.: Japan (Honshu, Shikoku, Kyushu, Izu Is., Tsushima, Yakushima), Ryukyu Is. (Amami-Ōshima), Formosa.

Tribe NIGIDIINI ツノヒョウタンクワガタ族

Genus *Nigidius* MACLEAY, 1819 ツノヒョウタンクワガタ属

Nigidius MACLEAY, 1819, Horae ent., 1, p. 108 (type-species: *Nigidius cornatus* MACLEAY, 1819).

Platycerus: GUÉRIN-MÉNEVILLE, 1829-'44, Icon. Règne Anim., 3, p. 109 (nec GEOFFROY, 1762).

Eudora CASTELNAU, 1840, Hist. Nat. Ins. Coléopt., 2, p. 174 (pars).

Hadronigidius KRAATZ, 1869, Dtsch. ent. Z., 40, p. 65 (pars).

N. lewisi BOILEAU, 1905 ルイスツノヒョウタンクワガタ

BOILEAU, 1905, Le Naturaliste, 27, p. 61 (Iles Liou-Kiou).

Distr.: Japan (Shikoku, Kyushu, Yakushima), Ryukyu Is. (Tokara, Amami and Okinawa groups), Hainan.

Subfamily LUCANINAE クワガタムシ亜科

Tribe PLATYCERINI ルリクワガタ族

Genus *Platycerus* GEOFFROY, 1762 ルリクワガタ属

Platycerus GEOFFROY, 1762, Hist. Ins. environs Paris, 1, p. 62 (type-species: *Scarabaeus caraboides* LINNÉ, 1758).

Systemocerus WEISE, 1883, Cat. Coleopt. Europae et Caucase, p. 93.

Systemus SHARP et MUIR, 1912, Trans. ent. Soc. London, 1912, p. 573.

P. acuticollis Y. KUROSAWA, 1969 コルリクワガタ

Y. KUROSAWA, 1969, Bull. Natn. Sci. Mus., Tokyo, 12, p. 478, figs. (Hōshi Spa, Gumma Pref.).

Distr.: Japan (Honshu, Shikoku, Kyushu), Korea.

P. delicatulus LEWIS, 1883 ルリクワガタ

LEWIS, 1883, Trans. ent. Soc. London, 1883, p. 338, pl. 14, fig. 3 (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Tribe LUCANINI ミヤマクワガタ族

Genus *Lucanus* SCOPOLI, 1763 ミヤマクワガタ属

Lucanus SCOPOLI, 1763, Ent. Carn., p. 1 (type-species: *Scarabaeus cervus* LINNÉ, 1758).

Hexaphyllus MULSANT, 1839, Ann. Soc. Agr., Lyon, **2**, p. 119.
Pseudolucanus HOPE et WESTWOOD, 1845, Cat. Lucan. Coleopt., p. 30.

Subgenus *Lucanus* s. str. ミヤマクワガタ亜属

- L. (L.) ferriei* PLANET, 1898 オオシマミヤマクワガタ
 PLANET, 1898, Le Naturaliste, **20**, p. 251, fig. (Iles Liou-Kiou).
 Distr.: Ryukyu Is. (Amami-Ōshima).
- L. (L.) gamunus* SAWADA et WATANABE, 1960 ミクラミヤマクワガタ
 SAWADA & WATANABE, 1960, J. Agr. Sci. Tokyo Nogyo Daigaku, **6**, p. 99, fig.
 (Mikura-jima).
 Distr.: Japan (Mikura-jima, Kōzu-jima).
- L. (L.) maculifemoratus* MOTSCHULSKY, 1861 ミヤマクワガタ
 MOTSCHULSKY, 1861, Étud. ent., **10**, p. 9 (Japan).
 Distr.: Japan, Kuril Is. (Kunashiri).
 forma *maculifemoratus* MOTSCHULSKY, s. str. (ミヤマクワガタ)
 MOTSCHULSKY, 1861, Étud. ent., **10**, p. 9 (Japan).
cantori: MOTSCHULSKY, 1860, Étud. ent., **9**, p. 16 (nec HOPE, 1842) (Japan).
sericans SNELLEN VAN VOLLENHOVEN, 1861, Tijdschr. Ent., **4**, p. 103 (Java).
montivagus MÖLLENKAMP, 1912, Int. ent. Z. Guben, (6), p. 128 (S.-O. Borneo).
balachowskyi LACROIX, 1968, Bull. Soc. ent. France, **73**, p. 180, fig. (Mukoda,
 Kyushu).
 Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu).
 forma *hopei* PARRY, 1862 (エゾミヤマクワガタ)
 PARRY, 1862, Proc. ent. Soc. London, **1862**, p. 108 (Japan).
elegans PLANET, 1898, Le Naturaliste, **20**, p. 19, fig. (*maculifemoratus* var.) (Yeso).
 Distr.: Japan (Hokkaido, Honshu, Sado, Kyushu), Kuril Is. (Kunashiri).
 forma *nakanei* Y. KUROSAWA, 1976 (フジミヤマクワガタ)
 Y. KUROSAWA, 1976, Bull. Natn. Sci. Mus., Tokyo, (A), **2**, p. 189.
maculifemoratus MOTSCHULSKY, auct.
 Distr.: Japan (Honshu, Izu-Ōshima, Kōzu-jima, Shikoku).

Tribe ODONTOLABINI マルバネクワガタ族

Genus *Neolucanus* THOMSON, 1862 マルバネクワガタ属

- Neolucanus* THOMSON, 1862, Ann. Soc. ent. France, (4), **2**, p. 415 (type-species:
Odontolabis baladeva HOPE, 1842).
Odontolabis HOPE et WESTWOOD, 1845, Cat. Lucan. Coleopt., p. 5 (pars).
Anoplocnemus BURMEISTER, 1847, Handb. d. Ent., **5**, p. 357 (pars).
Chalcodes: ARROW, 1935, Trans. ent. Soc. London, **83**, p. 107 (pars).
- N. insularis* MIWA, 1929 チャイロマルバネクワガタ
 MIWA, 1929, Ins. Matsum., **3**, p. 73, fig. (Ishigaki-jima).
 Distr.: Ryukyu Is. (Yaéyama group).
- N. saundersi* PARRY, 1864 タテヅノマルバネクワガタ
 PARRY, 1864, Trans. ent. Soc. London, (3), **2**, p. 20, pl. 9, fig. 3 (India).
baladeva: ARROW, 1953, Fauna Brit. India, Coleopt., Lammell., (4), p. 105, pl. 18,
 fig. 3 (nec HOPE, 1842).
 Distr.: Assam, Burma, Tonkin, S. China, Formosa (?), Ryukyu Is.

- subsp. *insulicola* Y. KUROSAWA, 1976 (ヤエヤママルバネクワガタ)
 Y. KUROSAWA, 1976, Bull. Natn. Sci. Mus., Tokyo, (A), 2, p. 190 (Iriomote-jima).
 Distr.: Ryukyu Is. (Yaéyama group).
- subsp. *protogenetivus* Y. KUROSAWA, 1976 (アマミマルバネクワガタ)
 Y. KUROSAWA, 1976, Bull. Natn. Sci. Mus., Tokyo, (A), 2, p. 190 (Amami-
 Ōshima).
 Distr.: Ryukyu Is. (Amami group).

Tribe CLADOGNATHINI ノコギリクワガタ族

Genus *Prismognathus* MOTSCHULSKY, 1860 オニクワガタ属

- Prismognathus* MOTSCHULSKY, 1860, Schrenk's Reisen in Amur-Lande, 2, p. 138 (type-species: *Prismognathus subaeneus* MOTSCHULSKY, 1860).
Cyclorasis THOMSON, 1860, Ann. Soc. ent. France, (4), 2, p. 397.
Dorcus: ARROW, 1935, Trans. ent. Soc. London, 88, p. 109 (pars).

- P. angularis* WATERHOUSE, 1874 オニクワガタ
 WATERHOUSE, 1874, Ent. mon. Mag., 11, p. 6 (Kawachi).
 Distr.: Japan, Sakhalin.
- subsp. *angularis* WATERHOUSE, s. str. (オニクワガタ)
 WATERHOUSE, 1874, Ent. mon. Mag., 11, p. 6 (Kawachi).
 Distr.: Japan (Hokkaido, Rishiri-tô, Honshu, Shikoku, Kyushu), Sakhalin.
- subsp. *morimotoi* Y. KUROSAWA, 1975 (キュウシュウオニクワガタ)
 Y. KUROSAWA, 1975, Mem. Natn. Sci. Mus., Tokyo, (8), p. 157 (Mt. Aoi-dake,
 Miyazaki Pref.).
 Distr.: Japan (S. Kyushu).
- P. dauricus* (MOTSCHULSKY, 1860) キンオニクワガタ
 MOTSCHULSKY, 1860, Schrenk's Reisen in Amur-Lande, 2, p. 137, pl. 9, fig. 11 (♀)
 (*Metopodontus*) (env. du fort Mariinsk, Daourie).
subaeneus MOTSCHULSKY, 1860, Schrenk's Reisen in Amur-Lande, 2, p. 138, pl. 9,
 fig. 12 (♂) (bords du fl. Amour).
jekeli PARRY, 1864, Trans. ent. Soc. London, (3), 2, p. 41, pl. 9, fig. 4 (*Cyclorasis*)
 (Chowsan, Korea).
 Distr.: Japan (Tsushima), Korea, Quel-part, Manchuria, E. Siberia., N. China.
- P. tokui* Y. KUROSAWA, 1975 ヤクシマオニクワガタ
 Y. KUROSAWA, 1975, Mem. Natn. Sci. Mus., Tokyo, (8), p. 156 (Yakushima).
 Distr.: Japan (Yakushima).

Genus *Prosopocoilus* HOPE et WESTWOOD, 1845 ノコギリクワガタ属

- Prosopocoilus* HOPE et WESTWOOD, 1845, Cat. Lucan. Coleopt., p. 4 (type-species:
Lucanus caviformis HOPE et WESTWOOD, 1845).
Metopodontus HOPE et WESTWOOD, 1845, ibid., p. 4.
Cladognathus BURMEISTER, 1847, Handb. d. Ent., 5, p. 364.
Psalidognathus MOTSCHULSKY, 1857, Étud. ent., 6, p. 29 (nec GREY, 1832).
Psalidoremus MOTSCHULSKY, 1861, ibid., 10, p. 13 (nom. nov. for *Psalidognathus*
 MOTSCHULSKY, 1857).
Hoplitocranum JAKOWLEW, 1896, Horae Soc. ent. Ross., 30, p. 172.
Prosopocoelus VAN ROON, 1910, in JUNK, Coleopt. Cat., (8), p. 25.

- Metopotropus* OBERTHÜR et HOULBERT, 1913, *Insecta*, **3**, p. 416.
Cyclotropus OBERTHÜR et HOULBERT, 1913, *ibid.*, **3**, p. 449.
Pelecognathus OBERTHÜR et HOULBERT, 1915, *ibid.*, **5**, p. 28.
Homoderinus KRIESCHE, 1926, *Sttet. ent. Zgt.*, **87**, p. 384.
Dorcus: ARROW, 1935, *Trans. ent. Soc. London*, **83**, p. 109 (pars).
Cladognathinus DIDIER et SÉGUY, 1952, *Rev. fr. Ent.*, **19**, p. 223.

- P. dissimilis* (BOILEAU, 1898) リュウキュウノコギリクワガタ
 BOILEAU, 1898, *Bull. Soc. ent. France*, **1898**, p. 95, fig. (*Psalidoremus*) (Iles Liou-Kiou).
 Distr.: Ryukyu Is.
 subsp. *elegans* (INAHARA, 1958) (トカラノコギリクワガタ)
 INAHARA, 1958, *Ent. mon. Mag.*, **94**, p. 12, pl. 1, figs. 1b, 3b (*Psalidoremus*) (Takara-jima).
 Distr.: Ryukyu Is. (Tokara group).
 subsp. *dissimilis* (BOILEAU), s. str. (リュウキュウノコギリクワガタ)
 BOILEAU, 1898, *Bull. Soc. ent. France*, **1898**, p. 95, fig. (*Psalidoremus*) (Iles Liou-Kiou).
 Distr.: Ryukyu Is. (Amami group).
 subsp. *okinawanus* NOMURA, 1962 (オキナワノコギリクワガタ)
 NOMURA, 1962, *Tôhō Gakuhō*, (12), p. 37.
 Distr.: Ryukyu Is. (Okinoerabu-jima, Okinawa, Kerama-jima, Kume-jima).
- P. inclinatus* (MOTSCHULSKY, 1857) ノコギリクワガタ
 MOTSCHULSKY, 1857, *Étud. ent.*, **6**, p. 19, pl. 1, fig. 11 (*Lucanus*) (Shimoda).
mandibularis THOMSON, 1862, *Ann. Soc. ent. France*, (4), **2**, p. 417 (*Cladognathus*) (Japan).
inflexus HAROLD, 1875, *Abh. naturw. Ver. Bremen*, **4**, p. 288 (*Psalidoremus*) (Hiogo).
 Distr.: Japan, Korea.
 subsp. *inclinatus* (MOTSCHULSKY), s. str. (ノコギリクワガタ)
 MOTSCHULSKY, 1857, *Étud. ent.*, **6**, p. 19, pl. 1, fig. 11 (*Lucanus*) (Shimoda).
 Distr.: Japan (Hokkaido, Honshu, Sado, Izu Is. excl. Hachijō-jima, Shikoku, Kyushu, Yakushima, Tanegashima, Kuroshima, Tsushima), Korea.
 subsp. *hachijoensis* NOMURA, 1860 (ハチジョウノコギリクワガタ)
 NOMURA, 1960, *Tôhō Gakuhō*, (10), p. 40 (Hachijō-jima).
 Distr.: Japan (Hachijō-jima).
- P. motschulskyi* (WATERHOUSE, 1869) タカサゴノコギリクワガタ
 WATERHOUSE, 1869, *Trans. ent. Soc. London*, **1869**, p. 16 (*Cladognathus*) (Japan or Indian Archipelago).
 Distr.: Ryukyu Is., Formosa.
 subsp. *pseudodissimilis* Y. KUROSAWA, 1976 (ヤエヤマノコギリクワガタ)
 Y. KUROSAWA, 1976, *Bull. Natn. Sci. Mus., Tokyo*, (A), **2**, p. 190, fig. 4 (Ishigaki-jima and Iriomote-jima).
 Distr.: Ryukyu Is. (Yaéyama group).

Tribe PSEUDODORCINI ネプトクワガタ族

Genus *Aegus* MACLEAY, 1819 ネプトクワガタ属

- Aegus* MACLEAY, 1819, *Horae ent.*, **1**, p. 112 (type-species: *Aegus chelifera* MACLEAY, 1819).

- Alcimus* FAIRMAIRE, 1849, Rev. Mag. Zool., (2), 1, p. 416 (nec LOEW, 1848).
Paraegus GAHAN, 1888, Proc. ent. Soc. London, 1888, p. 539.
Xenostomus BOILEAU, 1898, Bull. Soc. ent. France, 1898, p. 164.
Eubussea ZACHER, 1913, Arb. Biol. Anst. Berlin, 9, p. 93 (nom. nov. for *Alcimus* FAIRMAIRE, 1849).
Elsion KRIESCHE, 1920, Arch. Naturg., A, (86), p. 95.
Malietoa KRIESCHE, 1920, *ibid.*, A, (86), p. 104.
Odontaegus KRIESCHE, 1935, Koleopt. Rundschau, 21, p. 174.
- A. formosae** BATES, 1866 タイワンネプトクワガタ
 BATES, 1866, Proc. zool. Soc. London, 1866, p. 374 (Formosa).
 Distr.: Ryukyu Is., Formosa.
 subsp. *ishigakiensis* NOMURA, 1960 (ヤエヤマネプトクワガタ)
 NOMURA, 1960, Tôhō Gakuhō, (10), p. 43 (subsp. of *laevicollis* E. SAUNDERS, 1854) (Ishigaki-jima).
 Distr.: Ryukyu Is. (Yaéyama group).
- A. laevicollis** E. SAUNDERS, 1854 ネプトクワガタ
 E. SAUNDERS, 1854, Trans. ent. Soc. London, (2) 3, p. 54, pl. 4, fig. 8 (Tea districts of China).
punctiger E. SAUNDERS, 1854, *ibid.*, (2), 3, p. 55, pl. 3, fig. 6 (Tea districts of China).
labilis E. SAUNDERS, 1864, *ibid.*, (3), 2, p. 54, pl. 11, fig. 8 (Himalaya).
 Distr.: Japan, Ryukyu Is., China, Burma, Assam, Himalaya.
 subsp. *subnitidus* WATERHOUSE, 1873 (ネプトクワガタ)
 WATERHOUSE, 1873, Ent. mon. Mag., 9, p. 177 (Japan).
abei ICHIKAWA et IMANISHI, 1976, Elytra, 3, p. 10, pl. 1, fig. 3, pl. 2, figs. 3a, 3b, 3c (subsp. of *laevicollis* E. SAUNDERS, 1854) (Akuseki-jima, Tokara Is.).
fujitai ICHIKAWA et IMANISHI, 1976, *ibid.*, 3, p. 10, pl. 1, fig. 4, pl. 2, figs. 4a, 4b (subsp. of *laevicollis* E. SAUNDERS, 1854) (Hachijō-jima).
 Distr.: Japan (Honshu, Izu Is., Shikoku, Kyushu, Yakushima, Tanegashima, Tsushima), Ryukyu Is. (Tokara Group).
 subsp. *taurus* DIDIER, 1928 (アマミネプトクワガタ)
 DIDIER, 1928, Étud. Coleopt. Luc., (2), p. 55 (Iles Lieou-k'ieous: Oshima).
 Distr.: Ryukyu Is. (Amami group).
- A. nakanei** ICHIKAWA et IMANISHI, 1976 オキナワネプトクワガタ
 ICHIKAWA & IMANISHI, 1976, Elytra, 3, p. 9, pl. 1 fig. 1, pl. 2, figs. 1a, 1b (subsp. of *laevicollis* E. SAUNDERS, 1854) (Okinawa).
 Distr.: Ryukyu Is.
 subsp. *nakanei* ICHIKAWA et IMANISHI, s. str. (オキナワネプトクワガタ)
 ICHIKAWA & IMANISHI, 1976, Elytra, 3, p. 9, pl. 1, fig. 1, pl. 2, figs. 1a, 1b (subsp. of *laevicollis* E. SAUNDERS, 1854) (Okinawa).
 Distr.: Ryukyu Is. (Okinawa).
 subsp. *tamanukii* ICHIKAWA et IMANISHI, 1976 (エラブネプトクワガタ)
 ICHIKAWA & IMANISHI, 1976, Elytra, 3, p. 10, pl. 1, fig. 2, pl. 2, figs. 2a, 2b (subsp. of *laevicollis* E. SAUNDERS, 1854) (Okinoerabu-jima).
 Distr.: Ryukyu Is. (Okinoerabu-jima).
- A. ogasawarensis** OKAJIMA et KOBAYASHI, 1975 オガサワラネプトクワガタ
 OKAJIMA & KOBAYASHI, 1975, Bull. Natn. Sci. Mus., Tokyo, (A), 1, p. 119 (Haha-jima).

Distr.: Bonin Is. (Haha-jima).

Tribe DORCINI クワガタムシ族

Genus *Rhaetulus* WESTWOOD, 1871 シカクワガタ属

Rhaetulus WESTWOOD, 1871, Trans. ent. Soc. London, 1871, p. 353, pl. 8, fig. 4 (type-species: *Rhaetulus crenatus* WESTWOOD, 1871).

Dorcus: ARROW, 1943, Proc. ent. Soc. London, (B), 12, p. 142 (pars).

Subgenus *Rhaetulus* s. str. シカクワガタ亜属

R. (R.) recticornis Y. KUROSAWA, 1964 アマミシカクワガタ

Y. KUROSAWA, 1964, Kontyû, Tokyo, 32, p. 444, fig. 2 (subsp. of *crenatus* WESTWOOD, 1871) (Tokuno-shima).

Distr.: Ryukyu Is. (Amami group).

Genus *Macrodercas* MOTSCHULSKY, 1861 コクワガタ属

Macrodercas MOTSCHULSKY, 1861, Étud. ent., 10, p. 15 (type-species: *Psolidostomus ? rectus* MOTSCHULSKY, 1857).

Hemisodercus THOMSON, 1862, Ann. Soc. ent. France, (4), 2, p. 421.

Macrogathus HEYNE et TASCHEBERG, 1908, Exot. Käfer, p. 54 (nec LACEPEDE, 1800).

Macrodercus VAN ROON, 1910, in JUNK, Coleopt. Cat., 19, (8), p. 33.

Dorcus: ARROW, 1950, Fauna Brit. India, Coleopt., Lammel., (4), p. 78 (pars).

Pogonodercus SÉGUY, 1954, Rev. fr. Ent., 21, p. 185.

M. okinawanus NOMURA, 1964 リュウキュウコクワガタ

NOMURA, 1964, Ent. Rev. Japan, 17, p. 47, pl. 3, fig. 1 (subsp. of *rectus* MOTSCHULSKY, 1857) (Yona, Okinawa).

Distr.: Ryukyu Is.

subsp. *amamianus* NOMURA, 1964 (アマミコクワガタ)

NOMURA, 1964, *ibid.*, 17, p. 47, pl. 3, fig. 2 (subsp. of *rectus* MOTSCHULSKY, 1857) (Ikari, Amami-Oshima).

Distr.: Ryukyu Is. (Amami group).

subsp. *okinawanus* NOMURA, s. str. (リュウキュウコクワガタ)

NOMURA, 1964, *ibid.*, 17, pl. 3, fig. 1 (subsp. of *rectus* MOTSCHULSKY, 1857) (Yona, Okinawa).

Distr.: Ryukyu Is. (Okinawa group).

M. rectus (MOTSCHULSKY, 1857) コクワガタ

MOTSCHULSKY, 1857, Étud. ent., 6, p. 29 (*Psolidostomus ?*) (Simoda).

binervis MOTSCHULSKY, 1860, *ibid.*, 9, p. 16 (Japon).

rugipennis MOTSCHULSKY, 1861, *ibid.*, 10, p. 16 (Japon).

niponensis SNELLEN VAN VOLLENHOVEN, 1861, Tijdschr. Ent., 4, p. 113, pl. 7, fig. 3 (*Dorcus*) (Japan).

diabolicus THOMSON, 1862, Ann. Soc. ent. France, (4), 2, p. 423 (*Eurytrachelus*) (Japon).

montivagus: MATSUMURA, 1906, Thous. Ins. Japan, 3, pl. 45, fig. 1 (*Eurytrachelus*) (nec LEWIS, 1883).

Distr.: Japan (Hokkaido, Honshu, Sado, Izu Is., Shikoku, Kyushu, Tsushima, Yakushima, Tanegashima, Kuroshima, Kuchinoerabu-jima), Ryukyu Is. (Tokara group), Korea, China.

M. striatipennis MOTSCHULSKY, 1861 スジクワガタ

MOTSCHULSKY, 1861, Étud. ent., 10, p. 17 (Japon).

cribellatus MOTSCHULSKY, 1861, *Étud. ent.*, **10**, p. 17 (Japon).

opacus WATERHOUSE, 1870, *Ent. mon. Mag.*, **6**, p. 208 (Japan).

vanvolxemi LEWIS, 1879, *Ann. Mag. nat. Hist.*, (5), **2**, p. 462 (Yezo).

Distr.: Japan (Hokkaido, Rishiri-tô, Rebun-tô, Honshu, Sado, Mikura-jima, Shikoku, Kyushu, Tsushima, Yakushima), Korea, China, Formosa.

Genus *Nipponodorcus* NOMURA et Y. KUROSAWA, 1960 アカアシクワガタ属

Nipponodorcus NOMURA et Y. KUROSAWA, 1960, *Tôhô Gakuhô*, (10), p. 41 (type-species: *Eurytrachelus rubrofemoratus* SNELLEN VAN VOLLENHOVEN, 1865).

N. montivagus (LEWIS, 1883) ヒメオオクワガタ

LEWIS, 1883, *Trans. ent. Soc. London*, **1883**, p. 337, pl. 14, fig. 2 (*Macrodercus*) (Chiuzenji, Junsai, Nanaye).

ezoensis MIWA, 1927, *Ins. Matsum.*, **2**, p. 27 (*Eurytrachelus*) (Jôzankei).

Distr.: Japan (Hokkaido, Honshu, Izu-Ôshima, Shikoku, Kyushu), Korea.

N. rubrofemoratus (SNELLEN VAN VOLLENHOVEN, 1865) アカアシクワガタ

SNELLEN VAN VOLLENHOVEN, 1865, *Tijdschr. Ent.*, **8**, p. 152, pl. 2, figs. 1, 2 (*Eurytrachelus*) (Japon).

Distr.: Japan, Korea, China, Formosa, Burma.

subsp. *rubrofemoratus* (SNELLEN VAN VOLLENHOVEN), s. str. (アカアシクワガタ)

SNELLEN VAN VOLLENHOVEN, 1865, *Tijdschr. Ent.*, **8**, p. 152, pl. 2, figs. 1, 2 (*Eurytrachelus*) (Japon).

arrowi BOILEAU, 1911, *Trans. ent. Soc. London*, **1911**, p. 441 (*Hemisodorcus*) (Burma).

haitschunus DIDIER et SÉGUY, 1952, *Rev. fr. Ent.*, **19**, (4), p. 227, fig. 5 (*Eurytrachelellus*) (Fokien, China).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima), Korea, China, Tibet, Burma.

Genus *Serrognathus* MOTSCHULSKY, 1861 ヒラタクワガタ属

Serrognathus MOTSCHULSKY, 1861, *Étud. ent.*, **10**, p. 12 (type-species: *Serrognathus castanicolor* MOTSCHULSKY, 1861).

Platyprosopus HOPE et WESTWOOD, 1845, *Cat. Lucan. Coleopt.*, p. 6, 31 (nec MANNERHEIM, 1830).

Dorcus: BURMEISTER, 1847, *Handb. d. Ent.*, **5**, p. 383 (pars).

Eurytrachelus THOMSON, 1862, *Ann. Soc. ent. France*, (4), **2**, p. 421 (nec MOTSCHULSKY, 1850).

Eurytrachelellus DIDIER, 1931, *Étud. Coleopt. Lucan.*, (8), p. 185 (nom. nov. for *Eurytrachelus* THOMSON, 1862).

Brontodorcus DIDIER, 1931, *ibid.*, (8), p. 196.

Eurydorcus DIDIER, 1931, *ibid.*, (8), p. 196.

Goniodorcus DIDIER, 1931, *ibid.*, (8), p. 196.

Lasiodorcus DIDIER, 1931, *ibid.*, (8), p. 196.

Telodorcus DIDIER, 1931, *ibid.*, (8), p. 196.

S. consentaneus (ALBERS, 1886) チョウセンヒラタクワガタ

ALBERS, 1886, *Dtsch. ent. Z.*, **30**, p. 28 (*Eurytrachelus*) (Pekin).

fasolt: NOMURA, 1962, *Tôhô Gakuhô*, (12), p. 35 (pars).

Distr.: Japan (Tsushima), Korea, China.

S. costatus (BOILEAU, 1898) スジブトヒラタクワガタ

BOILEAU, 1898, *Le Naturaliste*, **20**, p. 45, fig. (*Eurytrachelus*) (L'île d'Oshima, archipel Liou Kiou).

Distr.: Ryukyu Is. (Amami group).

S. platymelus (E. SAUNDERS, 1854) ヒラタクワガタ

E. SAUNDERS, 1854, *Trans. ent. Soc. London*, (2), **3**, p. 50, pl. 3, fig. 7 (*Platyprosopus*) (Tea districts of China).

lateralis E. SAUNDERS, 1854, *ibid.*, (2), **3**, p. 46 (*Dorcus*) (nec DEJEAN, 1836) (err.).

obscurus E. SAUNDERS, 1854, *ibid.*, (2), **3**, p. 52, pl. 4, fig. 7 (*Dorcus*) (Tea districts of China).

marginalis E. SAUNDERS, 1854, *ibid.*, (2), **3**, p. 53, pl. 4, fig. 6 (*Dorcus*) (Tea districts of China).

Distr.: Japan, Korea, S. Manchuria, China, Ryukyu Is., Formosa, Tonkin, Burma.

subsp. *pilifer* (SNELLEN VAN VOLLENHOVEN, 1861) (ヒラタクワガタ)

SNELLEN VAN VOLLENHOVEN, 1861, *Tijdschr. Ent.*, **6**, p. 112, pl. 6, fig. 4 (*Dorcus*) (Japan).

Distr.: Japan (Honshu, N. Izu Is., Shikoku, Kyushu, Yakushima, Tanegashima), Ryukyu Is. (Tokara group).

subsp. *castanicolor* MOTSCHULSKY, 1861 (ツシマヒラタクワガタ)

MOTSCHULSKY, 1861, *Étud. ent.*, **10**, p. 12 (Tsushima).

fasolt KRIESCHE, 1920, *Arch. Naturg.*, **86**, p. 118 (*Eurytrachelus*) (Korea) (subsp. of *E. titanus* BOISDUVAL, 1835).

Distr.: Japan (Tsushima), Korea, S. Manchuria.

subsp. *elegans* (BOILEAU, 1899) (アマミヒラタクワガタ)

BOILEAU, 1899, *Bull. Soc. ent. France*, **1899**, p. 177 (*Eurytrachelus*) (Iles Liou Kiou).

Distr.: Ryukyu Is. (Amami group).

subsp. *okinawanus* (KRIESCHE, 1922) (オキナワヒラタクワガタ)

KRIESCHE, 1922, *Stett. ent. Ztg.*, **83**, p. 124 (*Eurytrachelus*) (subsp. of *E. titanus* BOISDUVAL, 1835) (Okinawa).

Distr.: Ryukyu Is. (Okinawa group).

subsp. *sakishimanus* NOMURA, 1864 (サキシマヒラタクワガタ)

NOMURA, 1964, *Tôhō Gakuhō*, (14), p. 62, pl. 1, figs. 1-3.

Distr.: Ryukyu Is. (Yaéyama group, Yonaguni-jima).

Genus **Dorcus** MACLEAY, 1819 オオクワガタ属

Dorcus MACLEAY, 1819, *Horae ent.*, **1**, p. 111 (type-species: *Scarabaeus parallelipipedus* LINNÉ, 1758).

Dynodorcus DIDIER, 1931, *Étud. Coleopt. Lucan.*, (8), p. 197.

D. hopei (E. SAUNDERS, 1854) オオクワガタ

E. SAUNDERS, 1854, *Trans. ent. Soc. London*, (2), **3**, p. 50, pl. 3, fig. 8 (*Platyprosopus*) (Tea districts of China).

striato-punctatus E. SAUNDERS, 1854, *ibid.*, (2), **3**, p. 51, pl. 4, fig. 5 (Tea districts of China).

striatus E. SAUNDERS, 1854, *ibid.*, (2), **3**, p. 53, pl. 4, fig. 4 (Tea districts of China).

binodulus WATERHOUSE, 1874, *Ent. mon. Mag.*, **11**, p. 6 (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima), Korea, China, Formosa(?).