

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 9

December 30, 1977

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.),
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family ELMINTHIDAE ヒメドロムシ科

By

Masataka SATÔ

Subfamily LARINAE ハバビロドロムシ亜科

Tribe POTAMOPHILINI ハバビロドロムシ族

Genus *Dryopomorphus* HINTON, 1936 ハバビロドロムシ属

Dryopomorphus HINTON, 1936, Entomologist, **69**, p. 165 (type-species: *Dryopomorphus extraneus* HINTON, 1936).

D. extraneus HINTON, 1936 ハバビロドロムシ
HINTON, 1936, Entomologist, **69**, p. 166, figs. 1, 2 (Japan).
Distr.: Japan (Honshu, Kyushu).

D. nakanei NOMURA, 1958 ヒメハバビロドロムシ
NOMURA, 1958, Tôhō-Gakuhô, (8), p. 45, pl. 1, fig. 4 (Noziri, Nagano Pref.).
Distr.: Japan (Honshu).

Subfamily ELMINTHINAE ヒメドロムシ亜科

Tribe ELMINTHINI ヒメドロムシ族

Subtribe STENELMINTHINA アシナガミゾドロムシ亜族

Genus *Stenelmis* DUFOUR, 1835 アシナガミゾドロムシ属

Stenelmis DUFOUR, 1835, Ann. Sci. Nat. Zool., **3**, p. 158 (type-species: *Limnius canaliculatus* GYLLENHAL, 1808).

S. aritai M. SATÔ, 1965 ヨナグニアシナガミゾドロムシ
M. SATÔ, 1965, Bull. Jap. ent. Acad., **1**, p. 32 (Tabaru-gawa, Yonaguni-jima).
Distr.: Ryukyu Is. (Yonaguni-jima).

S. hisamatsui M. SATÔ, 1960 アカハラアシナガミゾドロムシ
M. SATÔ, 1960, Kontyû, Tokyo, **28**, p. 253, figs. 1, 2 (Shin-mura, Amami-Ôshima).

Distr.: Ryukyu Is. (Amami-Ōshima, Tokuno-shima, Okinawa).

- S. ishiharai** M. SATŌ, 1965 ヤエヤマアシナガミゾドロムシ
M. SATŌ, 1965, Bull. Jap. ent. Acad., 1, p. 31 (Takeda, Ishigaki-jima).
Distr.: Ryukyu Is. (Ishigaki-jima, Iriomote-jima).
- S. miyamotoi** NOMURA et NAKANE, 1958 ミヤモトアシナガミゾドロムシ
NOMURA & NAKANE, 1958, Akitu, 7, p. 81, figs. a, b (Nagaragawa, Gifu Pref.).
Distr.: Japan (Honshu, Shikoku, Kyushu).
- S. nipponica** NOMURA, 1958 イブシアシナガミゾドロムシ
NOMURA, 1958, Ent. Rev. Japan, 9, p. 41, pl. 8, fig. 1 (Matsuyama, Ehime Pref.).
Distr.: Japan (Honshu, Shikoku, Kyushu).
- S. vulgaris** NOMURA, 1958 アシナガミゾドロムシ
NOMURA, 1958, Ent. Rev. Japan, 9, p. 44, pl. 8, fig. 3 (Wakamatsu, Fukushima Pref.).
Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Ordobrevia* SANDERSON, 1953 ミゾドロムシ属

Ordobrevia SANDERSON, 1953, J. Kansas ent. Soc., 26, p. 159 (type-species: *Stenelmis nubifera* FALL, 1901).

- O. amamiensis** (NOMURA, 1957) アマミミゾドロムシ
NOMURA, 1957, Akitu, 6, p. 4. (*Stenelmis*) (Nishinakama, Amami-Ōshima).
Distr.: Ryukyu Is. (Amami-Ōshima).
- subsp. *okinawana* NOMURA, 1959 (オキナワミゾドロムシ)
NOMURA, 1959, Tōhō-Gakuhō, (9), p. 34 (Nago, Okinawa).
Distr.: Ryukyu Is. (Okinawa).
- O. foveicollis** (SCHÖNFELDT, 1888) キスジミゾドロムシ
SCHÖNFELDT, 1888, Ent. Nachr., 14, p. 139 (*Stenelmis*) (Hiogo).
flavovittatus KŌNO, 1934, Ins. Matsum., 8, p. 126 (*Stenelmis*) (Iwate, Tokio).
freyi BOLLOW, 1941, Mitt. Münchner ent. Ges., 31, p. 21 (*Stenelmis*) (Hondo, Mte. Rokko, Nähe Tokio).
japonica JANSSENS, 1956, Bull. Inst. r. Sci. nat. Belg., 32, p. 2, fig. 1 (*Stenelmis*) (Kobé).
Distr.: Japan (Honshu, Shikoku, Kyushu).
- O. gotoi** NOMURA, 1959 ゴトウミゾドロムシ
NOMURA, 1959, Tōhō-Gakuhō, (9), p. 33, pl. 1, figs. 1, 2 (Inunaki, Wakayama Pref.).
Distr.: Japan (Honshu).
- O. maculata** (NOMURA, 1957) アカモンミゾドロムシ
NOMURA, 1957, Ent. Rev. Japan, 8, p. 43 (*Stenelmis*) (Yaku-shima).
Distr.: Japan (Honshu, Shikoku, Yaku-shima).

Genus *Leptelmis* SHARP, 1888 ヨコミゾドロムシ属

Leptelmis SHARP, 1888, Ann. Mag. nat. Hist., (6), 2, p. 243 (type-species: *Leptelmis gracilis* SHARP, 1888).

Lepthelmis: ZAITZEV, 1910, Junk, Col. Cat. Pars 17, Dryopidae, p. 21.

- L. gracilis** SHARP, 1888 ヨコミゾドロムシ
SHARP, 1888, Ann. Mag. nat. Hist., (6), 2, p. 244 (Tokyo).
Distr.: Japan (Honshu, Shikoku, Kyushu).

- L. parallela** NOMURA, 1962 ホソヨコミゾドロムシ
 NOMURA, 1962, Tōhō-Gakuhō, (12), p. 47 (Ikenobe, Kagawa Pref.).
 Distr.: Japan (Honshu, Shikoku, Kyushu).
- Genus **Graphelmis** DELÈVE, 1968 アヤスジミゾドロムシ属
Graphelmis DELÈVE, 1968, Ann. hist.-nat. Mus. natn. Hung., 60 Zool, p. 169 (type-species: *Graphelmis scapularis* DELÈVE, 1968).
- G. shirahatai** (NOMURA, 1958) アヤスジミゾドロムシ
 NOMURA, 1958, Ent. Rev. Japan, 9, p. 42, pl. 8, fig. 2 (*Stenelmis*) (Jimmachi, Yamagata Pref.).
 Distr.: Japan (Honshu).
- Subtribe ELMINTHINA ヒメドロムシ亜族
 Genus **Nomuraelmis** M. SATŌ, 1964 ノムラヒメドロムシ属
Nomuraelmis M. SATŌ, 1964, Bull. Jap. ent. Acad., 1, p. 11 (type-species: *Nomuraelmis amamiensis* M. SATŌ, 1964).
- N. amamiensis** M. SATŌ, 1964 ノムラヒメドロムシ
 M. SATŌ, 1964, Bull. Jap. ent. Acad., 1, p. 12 (Akaogi, Amami-Ōshima).
 Distr.: Ryukyu Is. (Amami-Ōshima).
- Genus **Neoriohelmis** NOMURA, 1958 クロサワドロムシ属
Neoriohelmis NOMURA, 1958, Tōhō-Gakuhō, (8), p. 45 (type-species: *Neoriohelmis kurosawai* NOMURA, 1958).
- N. kurosawai** NOMURA, 1958 クロサワドロムシ
 NOMURA, 1958, Tōhō-Gakuhō, (8), p. 46, pl. 1, figs. 1a-d (Mamigasaki-gawa, Yamagata Pref.).
 Distr.: Japan (Honshu).
- N. kuwatai** M. SATŌ, 1963 シコククロサワドロムシ
 M. SATŌ, 1963, Trans. Shikoku ent. Soc., 7, p. 132 (Hijikawa, Ehime Pref.).
 Distr.: Japan (Shikoku).
- Genus **Cleptelmis** SANDERSON, 1954 セマルヒメドロムシ属
Cleptelmis SANDERSON, 1954, J. Kansas ent. Soc., 27, p. 4 (type-species: *Elmis ornata* SCHAEFFER, 1911).
- C. parvula** NOMURA et BABA, 1961 セマルヒメドロムシ
 NOMURA & BABA, 1961, Akitu, 10, p. 4 (Kurokawa, Niigata Pref.).
 Distr.: Japan (Honshu).
- Genus **Pseudamophilus** BOLLOW, 1940 ケスジドロムシ属
Pseudamophilus BOLLOW, 1940, Arch. f. Zool., 32 (A-13), p. 9 (type-species: *Pseudamophilus malaisei* BOLLOW, 1940).
- P. japonicus** NOMURA, 1957 ケスジドロムシ
 NOMURA, 1957, Akitu, 6, p. 3, fig. 1 (Wakamatsu, Fukushima Pref.).
 Distr.: Japan (Honshu).
- Genus **Optioservus** SANDERSON, 1954 ヒメドロムシ属
Optioservus SANDERSON, 1954, J. Kansas ent. Soc., 27, p. 8 (type-species: *Limnius*

trivittatus BROWN, 1930).

Subgenus *Optioservus* SANDERSON, 1954, s. str.

- O. (O.) hayashii** NOMURA, 1960 スジヒメドロムシ
NOMURA, 1960, *Akitu*, **9**, p. 34, fig. 1 (Yachi Spa, Aomori Pref.).
Distr.: Japan (Honshu).
- O. (O.) maculatus** NOMURA, 1958 セアカヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 50, pl. 2, fig. 1 (Yudonosan, Yamagata Pref.).
Distr.: Japan (Honshu).
- O. (O.) nitidus** NOMURA, 1958 ツヤヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 53, pl. 2, fig. 6 (Shiromizu-gawa, Yamagata Pref.).
Distr.: Japan (Honshu).
- O. (O.) rugulosus** NOMURA, 1958 ヨツモンヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 53, pl. 2, fig. 7 (Furuya, Saga Pref.).
Distr.: Japan (Honshu, Kyushu).
- O. (O.) variabilis** NOMURA, 1958 スネアカヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 51, pl. 2, figs. 2-4 (Tokura, Gunma Pref.).
Distr.: Japan (Honshu).

Subgenus *Cyclolimnius* NOMURA, 1958 マルヒメドロムシ亜属

Cyclolimnius NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 54 (type-species: *Optioservus kubotai* NOMURA, 1958).

- O. (C.) ater** NOMURA, 1958 クロマルヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 56 (Bôhira, Yamagata Pref.).
Distr.: Japan (Honshu).
- O. (C.) hagai** NOMURA, 1958 ハガマルヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 56 (Kokura, Fukuoka Pref.).
Distr.: Japan (Kyushu).
- O. (C.) kubotai** NOMURA, 1958 クボタマルヒメドロムシ
NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 54 (Nikko, Tochigi Pref.).
Distr.: Japan (Honshu).

Genus *Grouvellinus* CHAMPION, 1923 ナガアシドロムシ属

Grouvellinus CHAMPION, 1923, *Ent. mon. Mag.*, **59**, p. 168 (type-species: *Macronychus caucasicus* MOTSCHULSKY, 1839).

Microdes MOTSCHULSKY, 1857, *Étud. ent.*, **8**, p. 48.

Grouvelleus ZAITZEV, 1908, *Horae. Soc. ent. Ross.*, **38**, p. 311.

- G. marginatus** (KÔNO, 1934) キベリナガアシドロムシ
KÔNO, 1934, *Ins. Matsum.*, **8**, p. 127 (*Grouvelleus*) (Koyadaira, Shikoku).
Distr.: Japan (Honshu, Shikoku, Kyushu).
- G. nitidus** NOMURA, 1963 ツヤナガアシドロムシ
NOMURA, 1963, *Tôhō-Gakuhō*, (13), p. 52 (Ryûzu, Nikko).
Distr.: Japan (Honshu).

G. subopacus NOMURA, 1962 マルナガアシドロムシNOMURA, 1962, *Tôhō-Gakuhō*, (12), p. 48 (Taken, Amami-Ōshima).

Distr.: Ryukyu Is. (Amami-Ōshima, Tokuno-shima, Okinawa).

Tribe MACRONYCHINI ツヤドロムシ族

Genus *Paramacronychus* NOMURA, 1958 ツブスジドロムシ属*Paramacronychus* NOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 47 (type-species: *Paramacronychus granulatus* NOMURA, 1958).**P. granulatus** NOMURA, 1958 ツブスジドロムシNOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 48 (Higashiyama, Fukushima Pref.).

Distr.: Japan (Honshu).

Genus *Urumaelmis* M. SATŌ, 1963 ウルマツヤドロムシ属*Urumaelmis* M. SATŌ, 1963, *New Entomologist*, **12**, p. 39 (type-species: *Zaitzevia uenoi* NOMURA, 1961). (subgen. of *Zaitzevia* CHAMPION, 1923)**U. uenoi** (NOMURA, 1961) ウエノツヤドロムシNOMURA, 1961, *Akitu*, **10**, p. 2 (*Zaitzevia*) (Ta-yō Cave, Okinoerabu Is.)*uenoi amamiana* M. SATŌ, 1963, *New Entomologist*, **12**, p. 41 (Ohtana, Amami-Ōshima). (subsp. of *uenoi* NOMURA, 1961)

Distr.: Ryukyu Is. (Amami-Ōshima, Okino-erabu-shima).

subsp. *tokarana* M. SATŌ, 1963 (トカラツヤドロムシ)M. SATŌ, 1963, *New Entomologist*, **12**, p. 40 (*Zaitzevia*) (Kuchino-shima).

Distr.: Ryukyu Is. (Kuchino-shima).

Genus *Zaitzeviaria* NOMURA, 1959 ヒメツヤドロムシ属*Zaitzeviaria* NOMURA, 1959, *Tôhō-Gakuhō*, (9), p. 35 (type-species: *Zaitzevia brevis* NOMURA, 1958). (subgen. of *Zaitzevia* CHAMPION, 1923)**Z. brevis** (NOMURA, 1958) ヒメツヤドロムシNOMURA, 1958, *Tôhō-Gakuhō*, (8), p. 49, pl. 1, figs. 3a-c (*Zaitzevia*) (Okadamura, Ehime Pref.).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Z. gotoi (NOMURA, 1959) ホソヒメツヤドロムシNOMURA, 1959, *Tôhō-Gakuhō*, (9), p. 36, pl. 1, figs. 8-10 (*Zaitzevia*) (Kimitōge, Wakayama Pref.).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Z. ovata (NOMURA, 1959) マルヒメツヤドロムシNOMURA, 1959, *Tôhō-Gakuhō*, (9), p. 35, pl. 1, figs. 4-7 (*Zaitzevia*) (Kimitōge, Wakayama Pref.).

Distr.: Japan (Honshu, Shikoku).

Genus *Zaitzevia* CHAMPION, 1923 ツヤドロムシ属*Zaitzevia* CHAMPION, 1923, *Ent. mon. Mag.*, **59**, p. 170 (type-species: *Zaitzevia solidicornis* CHAMPION, 1923).*Awadoronus* KŌNO, 1934, *Ins. Matsum.*, **8**, p. 127 (type-species: *Awadoronus awana* KŌNO, 1934).