

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 13

December 30, 1977

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.),
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family COLYDIIDAE ホソカタムシ科

By

Hiroyuki SASAJI

Subfamily COLYDIINAE ホソカタムシ亜科 (新称)

Tribe SYNCHITINI ヒラタホソカタムシ族 (新称)

Genus *Bitoma* HERBST, 1793 ナガセスジホソカタムシ属 (新称)

Bitoma HERBST, 1793, Käfer, 5, p. 26 (type-species: *Bitoma crenata* HERBST, 1793).

Ditoma ILLIGER, 1807, Mag. Ins., 6, p. 320.

Xuthia PASCOE, 1863, J. Ent., 2, p. 122, 128.

Euditomum GISTEL, 1857, Vacuna, 2, p. 524.

Eulachus LECONTE, 1863, New Spec. N. Amer. Col., 1, p. 68.

B. niponia (LEWIS, 1879) ヒメナガセスジホソカタムシ

LEWIS, 1879, Ann. Mag. nat. Hist., (5), 4, p. 462 (*Xuthia*) (Nagasaki).

Distr.: Japan (Kyushu).

B. parallela (SHARP, 1885) ナガセスジホソカタムシ

SHARP, 1885, J. Linn. Soc. London, 19, p. 70, pl. 6 (*Xuthia*) (Hitoyoshi, Yuyama).

Distr.: Japan (Honshu, Kyushu), Ryukyu Is., Formosa, Philippines, Ceylon, Madagascar, Equatorial Guinea, Marquesas Is., Réunion.

Genus *Microprius* FAIRMAIRE, 1868 ヒメセスジホソカタムシ属 (新称)

Microprius FAIRMAIRE, 1868, Ann. Soc. ent. France, (4), 8, p. 779 (type-species:
Phormesa demissus PASCOE, 1863).

Trionus SHARP, 1885, J. Linn. Soc. London, 19, p. 70.

M. opacus (SHARP, 1885) ツヤケンヒメホソカタムシ

SHARP, 1885, J. Linn. Soc. London, 19, p. 70, pl. 6 (*Trionus*) (Nagasaki, Ichiuchi, Yumamoto).

Distr.: Japan (Honshu, Shikoku, Kyushu), Philippines, Ceylon.

Genus *Cicones* CURTIS, 1829 ヒメヒラタホソカタムシ属 (新称)

Cicones CURTIS, 1829, Brit. Ent., 4, p. 149 (type-species: *Cicones carpini* CURTIS, 1829).

- C. bitomoides* SHARP, 1885 ヨコモンヒメヒラタホソカタムシ
SHARP, 1885, J. Linn. Soc. London, 19, p. 69 (Nara, Nagasaki).
Distr.: Japan (Honshu, Kyushu), Ceylon.
- C. minimus* SHARP, 1885 チビヒメヒラタホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, 19, p. 69 (Yuyama).
Distr.: Japan (Kyushu), Ceylon.
- C. niveus* SHARP, 1885 クロモンヒメヒラタホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, 19, p. 68 (Kashiwagi, Chiuzenji).
Distr.: Japan (Honshu).
- C. oblongus* SHARP, 1885 ホソヒメヒラタホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, 19, p. 68 (Sapporo).
Distr.: Japan (Hokkaido).
- C. oculatus* SHARP, 1885 メダカヒメヒラタホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, 19, p. 67 (Nikko, Wadatoge).
Distr.: Japan (Honshu).
- C. angustissimus* NAKANE, 1968 ナガヒラタホソカタムシ
NAKANE, 1968, Fragm. Coleopt., (18), p. 74 (Uji, Kyoto).
Distr.: Japan (Honshu).
- C. tokarensis* NAKANE, 1968 クロヒメヒラタホソカタムシ
NAKANE, 1968, Fragm. Coleopt., (19), p. 75 (Nakanoshima).
Distr.: Ryukyu Is. (Nakanoshima).
- C. hayashii* SASAJI, 1971 ハヤシヒメヒラタホソカタムシ (新称)
SASAJI, 1971, Life Study, Fukui, 15, p. 43 (Yokohama).
Distr.: Japan (Honshu).

Genus *Colobicus* LATREILLE, 1807 ヒラタホソカタムシ属 (新称)

Colobicus LATREILLE, 1807, Gen. Crust. Ins., 2, p. 9 (type-species: *Colobicus marginatus* LATREILLE, 1807).

- C. marginatus* LATREILLE, 1807 ヒラタホソカタムシ
LATREILLE, 1807, Gen. Crust. Ins., 2, p. 10, t. 16 (Europe?).
emarginatus ERICHSON, 1845, Naturg. Ins. Dtschl. Col., 3, p. 268.
axillaris DUFTSCHMIDT, 1825, Fauna Austr., 3, p. 155 (*Monotoma*).
hirsutus BRULLÉ, 1835, Hist. nat. Ins. Col., 5, p. 410.
Distr.: Japan (Hokkaido, Honshu), Siberia, Europe.
- C. granulatus* SHARP, 1885 オニヒラタホソカタムシ
SHARP, 1885, J. Linn. Soc. London, 19, p. 65 (Nikko).
Distr.: Japan (Honshu).

Genus *Acolophus* SHARP, 1885 マメヒラタホソカタムシ属 (新称)

Acolophus SHARP, 1885, J. Linn. Soc. London, 19, p. 66 (type-species: *Acolophus*

debilis SHARP, 1885).

- A. *debilis*** SHARP, 1885 マメヒラタホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, **19**, p. 66 (Nikko, Wada-toge).
Distr.: Japan (Honshu).

Genus *Neotrichus* SHARP, 1885 サンゲホソカタムシ属 (新称)

Neotrichus SHARP, 1885, J. Linn. Soc. London, **19**, p. 60 (type-species: *Neotrichus hispidus* SHARP, 1885).

- N. *hispidus*** SHARP, 1885 サンゲホソカタムシ
SHARP, 1885, J. Linn. Soc. London, **19**, p. 61, pl. 3 (Nagasaki, Oyayama, Hitoyoshi, Kashiwagi, Nikko).
Distr.: Japan (Honshu, Kyushu).

Genus *Ithris* PASCOE, 1863 ユミセスジホソカタムシ属 (新称)

Ithris PASCOE, 1863, J. Ent., **2**, p. 134 (type-species: *Ithris decisa* PASCOE, 1863).

- I. *sculpturata*** SHARP, 1885 ユミセスジホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, **19**, p. 72, pl. 3 (Oyayama).
Distr.: Japan (Kyushu).

Genus *Endophloeus* ERICHSON, 1845 ノコギリホソカタムシ属 (新称)

Endophloeus ERICHSON, 1845, Naturg. Ins. Dtschl. Col., **3**, p. 256 (type-species: *Eledona spinulosa* LATREILLE, 1807).

- E. *serratus*** SHARP, 1885 ノコギリホソカタムシ
SHARP, 1885, J. Linn. Soc. London, **19**, p. 61 (Hitoyoshi, Yuyama, Idzu, Miyanoshita, Kurigahara).
Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Sympanotus* SHARP, 1885 ホソマダラホソカタムシ属 (新称)

Sympanotus SHARP, 1885, J. Linn. Soc. London, **19**, p. 62 (type-species: *Sympanotus pictus* SHARP, 1885).

- S. *pictus*** SHARP, 1885 ホソマダラホソカタムシ
SHARP, 1885, J. Linn. Soc. London, **19**, p. 62, pl. 3 (Oyayama, Oyama).
Distr.: Japan (Honshu, Kyushu).

Tribe COXELINI ダルマホソカタムシ族 (新称)

Genus *Glyphocryptus* SHARP, 1885 ヒサゴホソカタムシ属 (新称)

Glyphocryptus SHARP, 1885, J. Linn. Soc. London, **19**, p. 63 (type-species: *Glyphocryptus brevicollis* SHARP, 1885).

- G. *brevicollis*** SHARP, 1885 ヒサゴホソカタムシ
SHARP, 1885, J. Linn. Soc. London, **19**, p. 64 (Yuyama).
Distr.: Japan (Kyushu, Tsushima, Izu-Miyakejima).

Genus *Pseudotarphius* WOLLASTON, 1874 ダルマチビホソカタムシ属 (新称)

Pseudotarphius WOLLASTON, 1874, Trans. ent. Soc. London, **1873**, p. 1 (type-species: *Pseudotarphius lewisi* WOLLASTON, 1873).

P. lewisi WOLLASTON, 1873 ダルマチビホソカタムシ

WOLLASTON, 1873, Trans. ent. Soc. London, **1873**, p. 4 (Nagasaki, Yuyama, Hagi).
Distr.: Japan (Honshu, Kyushu).

Genus **Trachypholis** ERICHSON, 1845 マダラホソカタムシ属 (新称)

Trachypholis ERICHSON, 1845, Naturg. Ins. Dtschl. Col., **3**, p. 257 (type-species: *Opatrum hispidum* WEBER, 1801).

Tarphioides WOLLASTON, 1862, J. Ent., **1**, p. 372.

Chorites PASCOE, 1862, J. Ent., **1**, p. 114.

Achorites GROUVELLE, 1911, Notes Leyden Mus., **33**, p. 154.

Labromimus SHARP, 1885, J. Linn. Soc. London, **19**, p. 64.

Optis PASCOE, 1885, Trans. ent. Soc. London, **1885**, Proc., p. 13.

T. variegata (SHARP, 1885) マダラホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 65, pl. 3 (*Labromimus*) (Oyama).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Tribe PYCNOMERINI ツヤナガホソカタムシ族 (新称)

Genus **Penthelispa** PASCOE, 1860 ツヤナガホソカタムシ属 (新称)

Penthelispa PASCOE, 1860, J. Ent., **1**, p. 111 (type-species: *Penthelispa porosa* PASCOE, 1860).

Pycnomeroplesius GANGLBAUER, 1899, Käf. Mitteleur., **3**, p. 885.

Endectus LECONTE, 1861, Classif. Col. N. Amer., **1**, p. 91.

P. sculpturatus (SHARP, 1885) アバタツヤナガヒラタホソカタムシ (新称)

SHARP, 1885, J. Linn. Soc. London, **19**, p. 77 (*Pycnomerus*) (Tonosawa near Miyanoshita).

Distr.: Japan (Honshu, Tsushima), Formosa.

P. vilis (SHARP, 1885) ツヤナガヒラタホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 77 (*Pycnomerus*) (Kyushu, Main Island, Yezo).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima).

Tribe GEMPYLODINI ナガホソカタムシ族 (新称)

Genus **Gempylodes** PASCOE, 1863 ナガホソカタムシ属 (新称)

Gempylodes PASCOE, 1863, J. Ent., **2**, p. 132 (type-species: *Gempylodes macer* PASCOE, 1863).

G. lewisii SHARP, 1885 ルイスホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 72, pl. 3 (Yuyama).

Distr.: Japan (Honshu, Shikoku, Kyushu), Formosa.

Subfamily BOTHRIDERINAE ムキヒゲホソカタムシ亜科 (新称)

Tribe DERETARPHINI ツツホソカタムシ族 (新称)

Genus **Teredolaemus** SHARP, 1885 ツヤツツホソカタムシ属 (新称)

Teredolaemus SHARP, 1885, J. Linn. Soc. London, **19**, p. 74 (type-species: *Teredus politus* LEWIS, 1879, designated here).

- T. politus** (LEWIS, 1879) クロツヤツツホソカタムシ (新称)
LEWIS, 1879, Ann. Mag. nat. Hist., (5), 4, p. 462 (*Teredus*) (South Japan).
Distr.: Japan (Shikoku, Kyushu).
- T. guttatus** SHARP, 1885 アトキツツホソカタムシ
SHARP, 1885, J. Linn. Soc. London, 19, p. 74 (Kashiwagi, Yuyama).
Distr.: Japan (Honshu, Shikoku, Kyushu).
- Genus *Cylindromicus* SHARP, 1885 セスジツツホソカタムシ属 (新称)
Cylindromicus SHARP, 1885, J. Linn. Soc. London, 19, p. 73 (type-species: *Cylindromicus gracilis* SHARP, 1885).
- C. gracilis** SHARP, 1885. セスジツツホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, 19, p. 73, pl. 3 (Oyayama).
Distr.: Japan (Kyushu).
- Tribe BOTHRIDERINI スジバネホソカタムシ族 (新称)
Genus *Bothriderus* ERICHSON, 1845 スジバネホソカタムシ属 (新称)
Bothriderus ERICHSON, 1845, Naturg. Ins. Dtschl. Col., 3, p. 288 (type-species: *Lycetus contractus* FABRICIUS, 1792).
- B. takeii** NAKANE, 1968 ムネクボスジホソカタムシ
NAKANE, 1968, Fragm. Coleopt., (19), p. 75 (Gumma Pref.: Numata).
Distr.: Japan (Honshu).
- Genus *Erotylathris* MOTSCHULSKY, 1861 フカミゾホソカタムシ属 (新称)
Erotylathris MOTSCHULSKY, 1861, Bull. Soc. Nat. Mosc., 34, p. 130 (type-species: *Erotylathris septemcostatus* MOTSCHULSKY, 1861).
- E. costatus** SHARP, 1885 フカミゾホソカタムシ
SHARP, 1885, J. Linn. Soc. London, 19, p. 75, pl. 3 (Sapporo, Nara, Nishimura).
Distr.: Japan (Hokkaido, Honshu), Formosa.
- Genus *Leptoglyphus* SHARP, 1885 ミスジホソカタムシ属 (新称)
Leptoglyphus SHARP, 1885, J. Linn. Soc. London, 19, p. 75 (type-species: *Leptoglyphus vittatus* SHARP, 1885).
- L. vittatus** SHARP, 1885 ミスジホソカタムシ (新称)
SHARP, 1885, J. Linn. Soc. London, 19, p. 75 (Hitoyoshi).
Distr.: Japan (Kyushu).
- Tribe DASTARCINI オオホソカタムシ族 (新称)
Genus *Dastarcus* WALKER, 1858 オオホソカタムシ属 (新称)
Dastarcus WALKER, 1858, Ann. Mag. nat. Hist., (3), 2, p. 209 (type-species: *Dastarcus porosus* WALKER, 1858).
- D. longulus** SHARP, 1885 サビマダラオオホソカタムシ
SHARP, 1885, J. Linn. Soc. London, 19, p. 76, pl. 3 (Konose in Higo).
Distr.: Japan (Honshu, Kyushu, Tsushima).