

CHECK-LIST
OF
COLEOPTERA OF JAPAN

~~~~~  
**日本産甲虫目録**

No. 14

December 30, 1977

The Coleopterists' Association of Japan  
c/o Department of Zoology, National Science Museum (Nat. Hist.),  
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

**Family CERYLONIDAE** カクホソカタムシ科

By

**Hiroyuki SASAJI**

Subfamily MURMIDIINAE チビマルホソカタムシ亜科 (新称)

Tribe MURMIDIINI チビマルホソカタムシ族 (新称)

Genus *Murmidius* LEACH, 1822 チビマルホソカタムシ属

*Murmidius* LEACH, 1822, Trans. Linn. Soc. London, **13**, p. 41 (type-species: *Murmidius ferrugineus* LEACH, 1822).

*Cautocerus* SCHUEPPEL, 1829, in GERMAR, Ins. Spec. novae, **1**, p. 85 (type-species: *Cautocerus advena* SCHUEPPEL, 1829).

*M. ovalis* (BECK, 1817) チビマルホソカタムシ

BECK, 1917, Beitr. baier. Insektenfauna, **7**, p. 1, t. 1 (*Hister*) (Europe?).

*ferrugineus* LEACH, 1922, Trans. Linn. Soc. London, **13**, p. 41.

*advena* SCHUEPPEL, 1829, in GERMAR, Ins. Spec. novae, **1**, p. 85 (*Cautocerus*).

Distr.: Japan (Honshu, Kyushu, Tsushima), China, Europe, Africa, India, Australia, North America.

Subfamily CERYLONINAE カクホソカタムシ亜科 (新称)

Tribe CERYLONINI カクホソカタムシ族 (新称)

Genus *Cerylon* LATREILLE, 1802 カクホソカタムシ属

*Cerylon* LATREILLE, 1802, Hist. nat. Gen. Crust. Ins., **3**, p. 205 (type-species: *Lycus histerooides* FABRICIUS, 1792).

*C. crassipes* SHARP, 1885 アシプトカクホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 81 (Oyayama, Miyanoshita, Nikko).

Distr.: Japan (Honshu, Shikoku, Kyushu).

*C. curticolle* SHARP, 1885 ムネヨコカクホソカタムシ (新称)

SHARP, 1885, J. Linn. Soc. London, **19**, p. 81 (Ichiuchi).

Distr.: Japan (Kyushu).

**C. *minimum*** SHARP, 1885 ヒメカクホソカタムシ (新称)

SHARP, 1885, J. Linn. Soc. London, **19**, p. 81 (Sapporo).

Distr.: Japan (Hokkaido).

**C. *sharpi*** NAKANE, 1967 カクホソカタムシ

NAKANE, 1967, *Fragm. Coleopt.*, (18), p. 74 (Tokugo near Kamikochi).

Distr.: Japan (Honshu).

**C. *takara*** NAKANE, 1967 ケナガカクホソカタムシ

NAKANE, 1967, *Fragm. Coleopt.*, (18), p. 73 (Takarajima).

Distr.: Ryukyu Is. (Takara I.).

Genus *Philothermus* AUBÉ, 1843 ダエンカクホソカタムシ属

*Philothermus* AUBÉ, 1843, *Ann. Soc. ent. France*, (2), **1**, p. 93 (type-species: *Philothermus montandoni* AUBÉ, 1843).

**P. *depressus*** SHARP, 1885 ダエンカクホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 78 (Miyanoshta, Hakone, Junsai, Sapporo).

Distr.: Japan (Hokkaido, Honshu, Kyushu).

**P. *pubens*** (SHARP, 1885) アメイロカクホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 80 (*Ectomicrus*) (Hitoyoshi, Oyayama, Yuyama, Miyanoshta).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Ectomicrus* SHARP, 1885 アラメカクホソカタムシ属 (新称)

*Ectomicrus* SHARP, 1885, J. Linn. Soc. London, **19**, p. 78 (type-species: *Ectomicrus rugicollis* SHARP, 1885).

**E. *rugicollis*** SHARP, 1885 アラメカクホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 79 (Subashiri, Oyayama).

Distr.: Japan (Honshu, Kyushu).

Tribe ACULAGNATHINI クチボソカクホソカタムシ族 (新称)

Genus *Thyroderus* SHARP, 1885 アナムネカクホソカタムシ属 (新称)

*Thyroderus* SHARP, 1885, J. Linn. Soc. London, **19**, p. 82 (type-species: *Thyroderus porcatus* SHARP, 1885).

**T. *porcatus*** SHARP, 1885 アナムネカクホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 83, pl. 3 (Nagasaki).

Distr.: Japan (Honshu, Kyushu).

Genus *Cautomus* SHARP, 1885 ムネビロカクホソカタムシ属 (新称)

SHARP, 1885, J. Linn. Soc. London, **19**, p. 82 (type-species: *Cautomus hystriculus* SHARP, 1885).

**C. *hystriculus*** SHARP, 1885 ムネビロカクホソカタムシ

SHARP, 1885, J. Linn. Soc. London, **19**, p. 82, pl. 3 (Nagasaki).

Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima).