

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 16

June 30, 1979

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chō, Shinjuku, Tokyo 160, Japan

Family CERAMBYCIDAE カミキリムシ科
(DISTENIINAE, PARANDRINAE, PRIONINAE,
SPONDYLINAE, ASEMINAE)

By

Masao HAYASHI

Subfamily DISTENIINAE ホソカミキリ亜科

Tribe DISTENINI ホソカミキリ族

Genus *Distenia* SERVILLE, 1825 ホソカミキリ属

Distenia SERVILLE, 1825, Encycl. Meth., 5, p. 485 (type species: *Distenia columbina* SERVILLE, 1825).

D. gracilis (BLESSIG, 1872) ホソカミキリ

BLESSIG, 1872, Horae Soc. ent. Rossicae, 9, p. 168, pl. 8, fig. 1 (*Apheles*) (Amur).

japonica BATES, 1873, Ann. Mag. nat. Hist., (4)-12, p. 155 (Japan).

Distr.: Japan (Hokkaido to Kyushu, Rishirito, Sado, Oki, Awajishima, Tsushima, Izu-miyakejima), Kurile Is., Sakhalin, Korea, N. & E. China, Siberia.

subsp. *yakushimana* YOKOYAMA, 1966 屋久島亜種

YOKOYAMA, 1966, Ent. Rev. Japan, 18, p. 55 (Yakushima).

Distr.: Japan (Yakushima).

Genus *Tengius* MATSUSHITA, 1938 オオクボカミキリ属

Tengius MATSUSHITA, 1938, Ins. Matsumurana, 12, p. 104 (type species: *Tengius ohkuboi* MATSUSHITA, 1938).

T. ohkuboi MATSUSHITA, 1938 オオクボカミキリ

MATSUSHITA, 1938, ibid., p. 105 (Kochi, Shikoku).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Subfamily PARANDRINAE ニセクワガタカミキリ亜科

Genus *Parandra* LATREILLE, 1804 ニセクワガタカミキリ属

Parandra LATREILLE, 1804, Hist. Nat. Crust, Inst., 11, p. 252 (type species: *P. laevis* LATREILLE, 1804).

P. formosana MIWA et MITONO, 1939 タイワンニセクワガタカミキリ

MIWA & MITONO, 1939, Trans. nat. Hist. Soc. Formosa, 29, p. 92, fig. (Formosa).
Distr.: Ryukyu Is. (Ishigaki, Iriomote), Formosa.

P. shibatai HAYASHI, 1963 アマミニセクワガタカミキリ

HAYASHI, 1963, Ent. Rev. Japan, 15, p. 50 (Amami-Ôshima, Botel-Tobago=Lan Hsu Is.).

janus (nec BATES) KANO, 1938, Annot. zool. Japon., 17, p. 115, pl. 7, fig. 1 (Botel-Tobago).

Distr.: Ryukyu Is. (Amami-Ôshima), Formosa (Botel-Tobago=Lan Hsu Is.).

Subfamily PRIONINAE ノコギリカミキリ亜科

Tribe MACROTOMINI トゲフチオオウスバカミキリ族

Genus *Macrotoma* SERVILLE, 1832 トゲフチオオウスバカミキリ属

Macrotoma SERVILLE, 1832, Ann. Soc. ent. France, 1, p. 137 (type species: *Prionus serripes* FABRICIUS, 1781).

Subgenus *Bandar* LAMEERE, 1912

Bandar LAMEERE, 1912, Mem. Soc. ent. Belg., 21, p. 144 (type species: *Macrotoma fisheri* WATERHOUSE, 1884).

M. (B.) fisheri WATERHOUSE, 1884 トゲフチオオウスバカミキリ

WATERHOUSE, 1884, Ann. Mag. nat. Hist., (5), 14, p. 382. (Burma).

Distr.: Ryukyu Is., Formosa, S. China, Laos, Viet Nam, Malaya, Burma.

subsp. *obscuribrunnea* HAYASHI, 1962 (コゲチャトゲフチオオウスバカミキリ)

HAYASHI, 1962, Ent. Rev. Japan, 15, p. 2, pl. 1, fig 1 (Ishigakijima).

Distr.: Ryukyu Is. (Ishigaki, Iriomote).

Tribe EURYPODINI ヒラタカミキリ族

Genus *Eurypoda* SAUNDERS, 1853 ヒラタカミキリ属

Eurypoda SAUNDERS, 1853, Trans. ent. Soc. London, (2), 2, p. 109 (type species: *E. antennata* SAUNDERS, 1853).

Zarax PASCOE, 1867, Ann. Mag. nat. Hist., (3), 19, p. 410 (type species: *Z. eurypodoides* PASCOE, 1867).

Subgenus *Eurypoda* s. str.

E. (E.) unicolor HAYASHI, 1956 コゲチャヒラタカミキリ

HAYASHI, 1956, Ent. Rev. Japan, 7, p. 11, pl. 3, fig. 1 (Yakushima; *antennata* SAUNDERS subsp.).

forma *tsushima* OHBAYASHI, 1963

OHBAYASHI, 1963, Fragn. Col., (2), p. 7 (Tsushima; *E. (E.) tsushima*).

Distr.: Japan (Shikoku, Yakushima, Tsushima).

E. (E.) boninensis HAYASHI et KUSAMA, 1973 オガサワラヒラタカミキリ

HAYASHI & KUSAMA, 1973, in KUSAMA, NARA & KUSUI, Rep. Fac. Sci., Shizuoka

Univ., 8, p. 117, pl. 1, fig. 1 (Chichijima).
 Distr.: Ogasawara Is. (Chichijima, Hahajima).

Subgenus *Neoprion* LACORDAIRE, 1869

Neoprion LACORDAIRE, 1869, Gen. Col., 8, p. 131 (type species: *N. parandraeformis* LACORDAIRE, 1869).

E. (N.) batesi GAHAN, 1894 ペーツヒラタカミキリ

GAHAN, 1894, Ann. Mag. nat. Hist., (6), 19, p. 225 (Yamaguchiya, Honshu, Japan).
 Distr.: Japan (Honshu, Izu-Miyakejima, Mikurajima, Shikoku, Kyushu, Oki, Yakushima, Tokara-Nakanoshima), Ryukyu Is. (Amami-Ôshima, Okinawa, Tokunoshima), SW. China, N. Viet Nam, Laos.

Tribe MEGOPIDINI ウスバカミキリ族

Genus *Megopis* SERVILLE, 1832 ウスバカミキリ属

Megopis SERVILLE, 1832, Ann. Soc. ent. France, 1, p. 161 (type species: *M. mutica* SERVILLE, 1832).

Subgenus *Spinimegopis* OHBAYASHI, 1963 トゲウスバカミキリ亜属

Spinimegopis OHBAYASHI, 1963, Fragm. Col., (2), p. 7 (type species: *Megopis (Aegosoma) nipponica* MATSUSHITA, 1934).

M. (S.) nipponica MATSUSHITA, 1934 トゲウスバカミキリ

MATSUSHITA, 1934, Trans. nat. Hist. Soc. Formosa, 24, p. 538, figs. 1a, 2a (*Megopis (Aegosoma)*) (Japan: Shikoku).
 Distr.: Japan (Shikoku, Kyushu, Yakushima).

M. (S.) kawazoei HAYASHI, 1961 アマミトゲウスバカミキリ

HAYASHI, 1961, Ent. Rev. Japan, 13, p. 36, pl. 9, fig. 2 (*Megopis (Aegosoma)*) (Amami-Ôshima).
 Distr.: Ryukyu Is. (Amami-Ôshima).

M. (S.) ishigakiana YOSHINAGA et NAKAYAMA, 1972 イシガキトゲウスバカミキリ

YOSHINAGA & NAKAYAMA, 1972, Gensei, Kochi, 23, p. 10 (Ishigakijima).
 Distr.: Ryukyu Is. (Ishigaki).

Subgenus *Aegosoma* SERVILLE, 1832 ウスバカミキリ亜属

Aegosoma SERVILLE, 1832, Ann. Soc. ent. France, 1, p. 162 (type species: *Prionus scabri-cornis* SCOPOLI, 1763).

M. (A.) sinica (WHITE, 1853) ウスバカミキリ

WHITE, 1853, Cat. Col. Brit. Mus., 7, p. 30 (*Aegosoma sinicum*) (Shanghai).
 Distr.: Japan, Korea, Formosa, China, Hainan, S. Viet Nam, Laos, Cambodia, Malaya, Burma, N. India, Borneo, Sumatra.

subsp. *sinica* (WHITE, 1853), s. str. (ウスバカミキリ)

WHITE, 1853, ibid., p. 30.

Distr.: Japan (Hokkaido to Kyushu, Rebunto, Izu-Niijima, Miyakejima, Hachijo-jima, Yakushima, Oki, Tsushima, Tokara-Nakanoshima), Ryukyu Is. (Amami-Ôshima, Okinawa), Formosa, China, Korea, Burma, Laos, S. Viet Nam.

subsp. *savoryi* KUSUI, 1973 小笠原亜種

KUSUI, 1973, in KUSAMA, NARA & KUSUI, Rep. Fac. Sci., Shizuoka Univ., 8, p. 119, pl. 1, fig. 6 (Ogasawarajima).

Distr.: Ogasawara Is.

M. (A.) validicornis GRESSITT, 1951 フトヒゲウスバカミキリ
GRESSITT, 1951, Philipp. J. Sci., 79, p. 205 (Ishigakijima).
Distr.: Ryukyu Is. (Ishigaki).

Tribe PRIONINI ノコギリカミキリ族

Genus *Prionus* GEOFFROY, 1762 ノコギリカミキリ属

Prionus GEOFFROY, 1762, Hist. abr. ins. Paris, 1, p. 198 (type species: *Cerambyx coriarius* LINNÉ, 1758).

P. insularis MOTSCHULSKY, 1857 ノコギリカミキリ
MOTSCHULSKY, 1857, Étud. ent., 6, p. 36 (Shimoda, Honshu).
Distr.: Japan (Hokkaido to Kyushu, Sado, Oki, Izu-Niijima, Kōzushima, Mikurajima, Gotō), Kurile Is., Korea, NE. China, E. Siberia.

P. yakushimanus OHBAYASHI, 1964 ヤクシマノコギリカミキリ
OHBAYASHI, 1964, Bull. Jap. ent. Acad., 1, p. 19 (Yakushima).
Distr.: Japan (Yakushima, Tanegashima).

P. sejunctus HAYASHI, 1959 ニセノコギリカミキリ
HAYASHI, 1959, Ent. Rev. Japan, 10, p. 4 (Mt. Hiei, Kyoto, Honshu).
Distr.: Japan (Honshu, Shikoku, Kyushu, Oki, Tsushima, Yakushima, Tanegashima).

Tribe ANACOLINI コバネカミキリ族

Genus *Psephactus* HAROLD, 1879 コバネカミキリ属

Psephactus HAROLD, 1879, Dtsch. ent. Z., 23, p. 367 (type species: *P. remiger* HAROLD, 1879).

P. remiger HAROLD, 1879 コバネカミキリ
HAROLD, 1879, ibid, p. 368 (Moheji, Japan).
Distr.: Japan (Hokkaido to Kyushu, Sado, Oki, Tsushima).
subsp. *insularis* HAYASHI, 1961
HAYASHI, 1961, Ent. Rev. Japan, 13, p. 37, pl. 9, fig. 3 (Amami-Ōshima, Yakushima).
Distr.: Japan (Izu-Mikurajima, Hachijojima, Yakushima, Tokara-Nakanoshima), Ryukyu Is. (Amami-Ōshima).

P. scabripennis KUSAMA, 1973 オガサワラコバネカミキリ
KUSAMA, 1973, in KUSAMA, NARA & KUSUI, Rep. Fac. Sci., Shizuoka Univ., 8, p. 118, pl. 1, figs. 2–4 (Ogasawarajima: Hahajima).
forma *chichijimensis* KUSAMA, 1973
KUSAMA, 1973, in KUSAMA, NARA & KUSUI, ibid, p. 119, pl. 1, fig. 5 (Ogasawara-jima: Chichijima).
Distr.: Ogasawara Is. (Chichijima, Hahajima).

Subfamily SPONDYLINAE クロカミキリ亜科

Genus *Spondylis* FABRICIUS, 1775 クロカミキリ属

Spondylis FABRICIUS, 1775, Syst. Ent., p. 159 (type species: *Attelabus buprestoides* LINNÉ, 1758).

S. buprestoides (LINNÉ), 1758 クロカミキリ
LINNÉ, 1758, Syst. Nat., ed. 10, p. 388 (*Attelabus*) (Europe).

Distr.: Japan (Hokkaido to Kyushu, Sado, Oki, Awajishima, Izu-Ōshima, Nijima, Kōzushima, Miyakejima, Tsushima, Tanegashima, Yakushima, Tokara-Nakano-shima), Ryukyu Is. (Amami-Ōshima, Okinawa), Sakhalin, Korea, China, Mongolia, Siberia, Caucasia, Central Asia, Europe.

Subfamily ASEMINAE マルクビカミキリ亜科

Tribe ASEMINI マルクビカミキリ族

Genus *Asemum* ESCHSCHOLTZ, 1830 マルクビカミキリ属

Asemum ESCHSCHOLTZ, 1830, Bull. Moscou, 2, p. 66 (type species: *Cerambyx striatus* LINNÉ, 1758).

A. amurense KRAATZ, 1879 マルクビヒラタカミキリ

KRAATZ, 1879, Dtsch. ent. Z., 23, p. 97 (Amur).

striatum LINNÉ subsp. *japonicum* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34, p. 235, pl. 2, fig. 7 (Nikko, Iwate, Chichibu, Honshu).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Tsushima), Sakhalin, Korea, NE. China, Mongolia, SE. Siberia.

A. punctulatum BLESSIG, 1872 ヒメマルクビヒラタカミキリ

BLESSIG, 1872, Horae Soc. ent. Rossicae, 9, p. 182 (E. Siberia).

Distr.: Japan (Hokkaido, Honshu, Rishirito), Kurile Is., Sakhalin, Korea, N. & C. China, Mongolia, Siberia, Central Asia.

Genus *Arhopalus* SERVILLE, 1834 ムナクボカミキリ属

Arhopalus SERVILLE, 1834, Ann. Soc. ent. France, 3, p. 77 (type species: *Cerambyx rusticus* LINNÉ, 1758).

A. rusticus (LINNÉ, 1758) ムナクボカミキリ

LINNÉ, 1758, Syst. Nat., ed. 10, p. 395 (*Cerambyx*) (Europe).

Distr.: Japan (Hokkaido to Kyushu, Izu-Kōzushima, Tsushima), Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Caucasia, Europe.

A. tobirensis HAYASHI, 1968 シナノムナクボカミキリ

HAYASHI, 1968, Ent. Rev. Japan, 20, p. 20 (Nagano, Honshu).

Distr.: Japan (Honshu).

Genus *Megasemum* KRAATZ, 1879 オオクロカミキリ属

Megasemum KRAATZ, 1879, Dtsch. ent. Z., 23, p. 97 (type species: *M. quadricostulatum* KRAATZ, 1879).

M. quadricostulatum KRAATZ, 1879 オオクロカミキリ

KRAATZ, 1879, ibid., p. 98 (Amur) ♀.

4-costulatum Kr. var. *brevior* PIC, 1901, L'Échange, 18, p. 11 (Yokohama, Japan) ♂.

Distr.: Japan (Hokkaido to Kyushu), Kurile Is., Sakhalin, Korea, China, Siberia.

Genus *Cephalallus* SHARP, 1905 ツシマムナクボカミキリ属

Cephalallus SHARP, 1905, Trans. ent. Soc. London, 1905, p. 148 (type species: *Criocephalus* (*Cephalallus*) *oberthuri* SHARP, 1905).

C. unicolor GAHAN, 1906 ツシマムナクボカミキリ

GAHAN, 1906, Fauna Brit. India, Col., 1, p. 97 (Assam, Tsushima, Japan).

sharpi REITTER, 1912, Fauna Germanica, 4, p. 43, foot note (*Megasemum*) (Japan).

projectum OKAMOTO, 1927, Ins. Matsumurana, 2, p. 63 (*Megesemum*) ♀ (Korea).
 Distr.: Japan (Honshu, Izu-Ōshima, Niijima, Kōzushima, Miyakejima, Shikoku, Kyushu, Oki, Tsushima, Tanegashima, Yakushima), Ryukyu Is. (Amami-Ōshima, Okinawa), Korea, C. China, Burma, Laos, Assam.

Genus *Nothorhina* REDTENBACHER, 1845 ケブカヒラタカミキリ属

Nothorhina REDTENBACHER, 1845, Gatt. Dtsch. Käferf., p. 102 (type species: *Callidium punctatum* FABRICIUS, 1798).

N. punctata (FABRICIUS, 1798) ケブカヒラタカミキリ

FABRICIUS, 1798, Suppl. ent. Syst., p. 149 (*Callidium punctatum*) (Europe).

Distr.: Japan (Honshu, Sado, Kyushu), Siberia, Southwest Himalayas, Syria, Europe.

Genus *Tetropium* KIRBY, 1837 トドマツカミキリ属

Tetropium KIRBY, 1837, in RICHARDSON, Fauna Borealli-Americanica, 4, p. 174 (type species: *T. cinnamopterum* KIRBY, 1837).

T. castaneum (LINNÉ, 1758) トドマツカミキリ

LINNÉ, 1758, Syst. Nat., ed. 10, p. 396 (*Cerambyx*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Korea, NE. China, Siberia, Europe.

T. gracilicorne REITTER, 1889 ホソトドマツカミキリ

REITTER, 1889, Dtsch. ent. Z., 33, p. 287 (Amur).

Distr.: Japan (Hokkaido), Sakhalin, Korea, NE. China, Mongolia, E. Siberia.

Tribe ATIMINI ケブカマルクビカミキリ族

Genus *Atimia* HALDEMAN, 1847 ケブカマルクビカミキリ属

Atimia HALDEMAN, 1847, Trans. Amer. Philos. Soc., (2), 10, p. 56 (type species: *Atimia tristis* HALDEMAN, 1847).

A. okayamensis HAYASHI, 1972 ケブカマルクビカミキリ

HAYASHI, 1972, Ent. Rev. Japan, 24, p. 26 (Okayama, Honshu).

Distr.: Japan (Honshu).

A. fujimurai HAYASHI, 1972 ヒメケブカマルクビカミキリ

HAYASHI, 1972, ibid., p. 26 (Okutama, Tokyo, Honshu).

Distr.: Japan (Honshu).

