

CHECK-LIST  
OF  
COLEOPTERA OF JAPAN

---

---

日本産甲虫目録

---

No. 17

June 30, 1979

---

The Coleopterists' Association of Japan  
c/o Department of Zoology, National Science Museum (Nat. Hist.)  
3-23-1 Hyakunin-chō, Shinjuku, Tokyo 160, Japan

---

Family CHRYSOMELIDAE ハムシ科  
(HISPINAE トゲハムシ亜科)

By

Shinsaku KIMOTO

Subfamily HISPINAE トゲハムシ亜科

Genus *Leptispa* BALY, 1858 ホソヒラタハムシ属

*Leptomorpha* GERMAR, 1842, Fauna Ins. Eur., 22, p. 10 (nec FALDERMAN).

*Leptispa* BALY, 1858, Cat. Hisp., 1858, p. 1 (type species: *Leptomorpha filiformis* GERMAR, 1842).

*Parallelispa* FAIRMAIRE, 1884, Ann. Soc. ent. France, (6), 4, p. 238.

*Paradownesia* GESTRO, 1899, Ann. Mus. Stor. Nat. Genova, (2), 20, p. 220, p. 224.

*L. miyamotoi* KIMOTO, 1957 ミヤモトホソヒラタハムシ (ミヤモトトゲナシトゲトゲ)

KIMOTO, 1957, Ins. Matsumurana, 21, p. 77, fig. 1 (Yuwan in Amami-Ōshima).  
Distr.: Ryukyu Is. (Amami-Ōshima).

*L. taguchii* CHŪJŌ, 1956 タグチホソヒラタハムシ (タグチトゲナシトゲトゲ)

CHŪJŌ, 1956, Mem. Fac. Lib. Arts & Educ., 21 (31), p. 18, fig. 9 (Kawagoe-shi in Saitama Pref.).

*godwini*: Lewis, 1888, Ent. mon. Mag., 25, p. 94 (near Tokio).

Distr.: Japan (Honshu, Kyushu).

Genus *Agonita* STRAND, 1942 オキナワホソヒラタハムシ属

*Agonia* WEISE, 1905, Dtsch. ent. Z., 1905, p. 116 (nec FORSTER).

*Agonita* STRAND, 1942, Folia Zool. Hydrob., 11, p. 391 (n. n. for *Agonia* WEISE, nec FORSTER; type species: *Gonophora wallacei* BALY, 1858).

*A. omoro* TAKIZAWA, 1975 オキナワホソヒラタハムシ

TAKIZAWA, 1975, Ent. Rev. Japan, 28, p. 57, fig. 3 (Omotodake in Ishigaki I.).

Distr.: Ryukyu Is. (Ishigaki).

Genus *Hispellinus* WEISE, 1897 クロトゲハムシ属

*Monochirus* CHAPUIS, 1875, in LACORDAIRE, 1875, Gen. Col., **11**, p. 330 (nec RAFINESQUE).  
*Hispellinus* WEISE, 1897, Dtsch. ent. Z., **1897**, p. 144 (type species: *Hispa multispinosa* GERMAR, 1848).

*H. moerens* (BALY, 1874) クロトゲハムシ

BALY, 1874, Trans. ent. Soc. London, **1874**, p. 215 (*Hispa*) (N. China).  
Distr.: Japan (Honshu, Shikoku, Kyushu), E. Siberia, China, Taiwan.

Genus *Asamangulia* MAULIK, 1915 ヨナグニトゲハムシ属

*Asamangulia* MAULIK, 1915, Rec. Ind. Mus., **11**, p. 378 (type species: *Asamangulia cuspidata* MAULIK, 1915).

*A. yonakuni* KIMOTO et GRESSITT, 1966 ヨナグニトゲハムシ

KIMOTO & GRESSITT, 1966, Pacif. Ins., **8**, p. 570, fig. 48a (Yonakuni I.).  
Distr.: Ryukyu Is. (Yonaguni).

Genus *Rhadinosa* WEISE, 1905 クロルリトゲハムシ属

*Rhadinosa* WEISE, 1905, Dtsch. ent. Z., **1905**, p. 318 (type species: *Hispa nigrocyanea*

*R. nigrocyanea* MOTSCHULSKY, 1861 クロルリトゲハムシ

MOTSCHULSKY, 1861, Schrenck's Reisen Amurl., **2**, p. 237, pl. 11, fig. 26 (*Hispa*) (Amur).

Distr.: Japan (Honshu, Shikoku, Kyushu), S. Siberia, China, Hainan, Korea.

Genus *Dactylispa* WEISE, 1897 トゲハムシ属

*Dactylispa* WEISE, 1897, Dtsch. ent. Z., **1897**, p. 137 (type species: *Dactylispa andrewesi* WEISE, 1897, = *Hispa severinii* GESTRO, 1897).

*Monohispa* WEISE, 1897, ibid., p. 147.

*Triplispa* WEISE, 1897, ibid., p. 150.

*D. angulosa* (SOLSKY, 1871) ヒメキベリトゲハムシ

SOLSKY, 1871, Horae Soc. ent. Ross., **8**, p. 262 (*Hispa*) (Amur).

*japonica* BALY, 1874, Trans. ent. Soc. London, **1874**, p. 215 (*Hispa*) (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), E. Siberia, China, Korea.

*D. higoniae* (LEWIS, 1896) ヒゴトゲハムシ

LEWIS, 1896, Ann. Mag. nat. Hist., (6), **17**, p. 342 (*Hispa*) (Japan).

*corpulentina* UHMANN, 1927, Suppl. Ent., **16**, p. 113, fig. 2 (Formosa).

Distr.: Japan (Honshu, Shikoku, Kyushu), S. China, Hainan, Taiwan.

*D. issikii* CHÛJÔ, 1938 タケトゲハムシ

CHÛJÔ, 1938, Bull. Umeno ent. Lab., (6), p. 12, fig. 1 (Kyushu).

Distr. Japan (Kyushu), China.

*D. masonii* GESTRO, 1923 キベリトゲハムシ

GESTRO, 1923, Ann. Mus. Stor. Nat. Genova, **51**, p. 9 (China).

*flavomarginata* SHIRÔZU, 1957, Sieboldia, Fukuoka, **2**, p. 53, pl. 6, fig. (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima), E. Siberia, China.

*D. subquadrata* (BALY, 1874) カタビロトゲハムシ

BALY, 1874, Trans. ent. Soc. London, **1874**, p. 216 (*Hispa*) (Japan).

*adstricta* WEISE, 1922, Phil. J. Sci., 21, p. 81 (Japan).

Distr.: Japan (Honshu, Sado I., Shikoku, Kyushu, Tsushima), E. China.

Genus *Dicladispa* GESTRO, 1897 イネトゲハムシ属

*Dicladispa* GESTRO, 1897, Ann. Mus. Stor. Nat. Genova, (2), 18, p. 81 (type species: *Hispa testacea* LINNAEUS).

*D. boutani* (WEISE, 1905) イネトゲハムシ

WEISE, 1905, Archiv Naturg., 71, p. 101 (*Hispa*) (Tonkin).

*similis* UHMANN, 1927, Suppl. Ent., 16, p. 116 (*Hispa*) (Formosa).

Distr.: Ryukyu Is. (Iriomote), Vietnam, S. China, Taiwan.

Genus *Platypria* GUÉRIN-MÉNEVILLE, 1840 ヘリビロトゲハムシ属

*Lobacantha* KIRBY, 1837, in RICHARDSON, Fauna Boreal.-Amer. 4, p. 227 (nomen. suppr.; HEMMING, 1955, in Int. Comm. Zool. Nomencl. 11, Opin. 376, p. 393).

*Platypria* GUÉRIN-MÉNEVILLE, 1840, Rev. Zool. Paris, 1840, p. 139 (type species: *Platypria echidna* GUÉRIN-MÉNEVILLE, 1840).

*P. echidna* GUÉRIN-MÉNEVILLE, 1840 ヘリビロトゲハムシ

GUÉRIN-MÉNEVILLE, 1840, Rev. Zool. Paris, 1840, p. 139 (India).

Distr.: Ryukyu Is. (?Okinawa), India, Ceylon, Sikkim, Burma, Indo-China.

*P. melli* UHMANN, 1955 ツシマヘリビロトゲハムシ

UHMANN, 1955, Ent. Blätter, 50, p. 211, fig. 10 (S. China: Kwangtung, Fukien).

Distr.: Japan (Tsushima), S. China.