

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 18

August 30, 1980

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family CHRYSOMELIDAE ハムシ科
(EUMOLPINAЕ サルハムシ亜科)

By

Shinsaku KIMOTO

Subfamily EUMOLPINAЕ サルハムシ亜科

Genus *Abirus* CHAPUIS, 1874 ツツサルハムシ属

Abirus CHAPUIS, 1874, Gen. Col., 10, p. 310 (type-species: *Cryptocephalus aeneus* WIEDEMANN, 1821).

A. fortunei (BALY, 1864) ツツサルハムシ

BALY, 1864, J. Ent., 1, p. 283 (*Dermorrhytis*) (N. China).

recticollis PIC, 1927, Bull. Soc. Linn. Lyon, 6, p. 133 (China).

sinensis PIC, 1927, *ibid.* (Yunnan).

yashiroi YUASA, Proc. Imp. Acad. Tokyo, 6 (7), p. 294, fig. 2 (Okinawa).

kiotoensis PIC, 1944, Échange, 60, p. 8 ("Kioto").

sinensis var. *guerryi* PIC, 1946, *ibid.* 62, p. 14 (China).

Distr.: Vietnam, S. China, Taiwan, Korea, Ryukyu Is. (Okinoerabu, Okinawa).

Genus *Acrothinium* MARSHALL, 1864 アカガネサルハムシ属

Acrothinium MARSHALL, 1864, Proc. Linn. Soc. London, 8, p. 47 (type-species: *Chrysochus gaschkevitchii* MOTSCHULSKY, 1860).

A. gaschkevitchii (MOTSCHULSKY, 1860) アカガネサルハムシ

MOTSCHULSKY, 1860, Étud. ent., 9, p. 23 (*Chrysochus*) (Japan).

gaschkevitchii f. *miyakensis* OHNO, 1968, Ent. Rev. Japan, 21, p. 35 (Miyake-jima).

gaschkevitchii f. *obscurum* OHNO, 1968, *ibid.* (Miyake-jima).

Distr.: E. Siberia, China, Taiwan, Japan, Ryukyu Is.

subsp. *gaschkevitchii* (MOTSCHULSKY, 1860)MOTSCHULSKY, 1860, Étud. ent., **9**, p. 23 (*Chrysochus*) (Japan).

Distr.: E. Siberia, China, Taiwan, Japan (Hokkaido, Honshu, Awashima, Sado, Hachijō, Shikoku, Kyushu, Tsushima, Yakushima, Kuchinoerabu), Ryukyu Is. (Miyako).

subsp. *tokaraense* NAKANE, 1956NAKANE, 1956, Sci. Rept. Saikyo Univ., (A), **2** (3), p. 170 (Nakanoshima in Tokara)

Distr.: Ryukyu Is. (Tokara).

subsp. *shirakii* NAKANE, 1956NAKANE, 1956, Sci. Rept. Saikyo Univ., (A), **2** (3), p. 170 (Shimokawa and Shinmura in Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima, Okinawa).

subsp. *matsuii* NAKANE, 1956NAKANE, 1956, Sci. Rept. Saikyo Univ., (A), **2** (3), p. 170 (Okinoerabu).

Distr.: Ryukyu Is. (Okinoerabu).

Genus *Cleoporus* LEFÈVRE, 1884 サクラサルハムシ属*Cleoporus* LEFÈVRE, 1884, Bull. Soc. ent. France, (6), **4**, p. 76 (type-species: *Cleoporus cruciatus* LEFÈVRE, 1884).*C. variabilis* (BALY, 1874) サクラサルハムシ(ヒトミヒメサルハムシ)BALY, 1874, Trans. ent. Soc. London, **1874**, p. 166 (*Paria*) (Nagasaki, E. Siberia).*robustus* BALY, 1874, *ibid.* (*Paria*) (Nagasaki).*tibialis* LEFÈVRE, 1885, Ann. Soc. ent. France, (6), **5**, p. 65 (*Stethotes*) (Tonkin).*pallidipes* FAIRMAIRE, 1888, Ann. Soc. ent. Belg., **32**, p. 36 (*Stethotes*) (Kiangsi).*rufipes* LEFÈVRE, 1889, Ann. Soc. ent. France, (6), **9**, p. 293 (*Mouhotina*) (Cochin-China, Annam, Tonkin).*niger* WEISE, 1922, Tijdschr. Ent., **65**, p. 52 (Fukien).*pygmaeus* WEISE, 1922, *ibid.*, p. 53 (Fukien).*tibialis* var. *aeneipennis* CHEN, 1936, Sinensia, **6** (3), p. 286 (Luc-nam, Tonkin).

Distr.: E. Siberia, China, Korea, Taiwan, Vietnam, Cambodia, Japan (Honshu, Shikoku, Kyushu).

Genus *Colaspoides* LAPORTE, 1833 ミドリサルハムシ属*Colaspoides* LAPORTE, 1833, in Silbermann, Rev. Ent., **1**, p. 20 (type-species: *Cryptoccephalus limbatus* FABRICIUS, 1781).*Pleuraulaca* CHEVROLAT, 1837, in DEJEAN, Cat. Col. ed. **3**, p. 409.*Amasia* CHAPUIS, 1874, Gen. Col. **10**, p. 313.*C. fulva* (CHŪJŌ, 1935) シヤンボサルハムシCHŪJŌ, 1935, Trans. nat. Hist. Soc. Formosa, **25**, p. 76 (Iriomote, Ishigaki).*japana*: NAKANE & KIMOTO, 1961, Kontyû, Tokyo, **29**, p. 17 (Amami-Oshima).

Distr.: Japan (Yakushima), Ryukyu Is. (Amami-Oshima, Tokunoshima, Okinawa, Ishigaki, Iriomote, Yonaguni).

C. japana CHŪJŌ, 1956 ミドリサルハムシCHŪJŌ, 1956, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., **2** (31), p. 3, fig. 2 (Miyazaki, Kagoshima).

Distr.: Japan (Kyushu).

Genus *Platycorynus* CHEVROLAT, 1837 オオミドリサルハムシ属

- Platycorynus* CHEVROLAT, 1837, in DEJEAN, Cat. Col. ed. 3, p. 413 (type-species: *Eumolpus compressicornis* FABRICIUS, 1801).
Corynodes HOPE, 1840, Col. Manual., 3, p. 162.
Eudora LAPORTE, 1840, Hist. Nat. Ins. Col., 2, p. 513.
Corynoeides CLARK, 1865, Ann. Mag. nat. Hist., (3), 15, p. 139.
Theumorus MARSHALL, 1865, Proc. Linn. Soc. Zool. London, 8, p. 35.
Eurycorynus MARSHALL, 1865, *ibid.*, p. 36.
Omodon MARSHALL, 1865, *ibid.*, p. 44.
Erigenes MARSHALL, 1865, *ibid.*, p. 45.
Batycolpus MARSHALL, 1865, *ibid.*, p. 46.

P. japonicus (JACOBY, 1896) オオミドリサルハムシ

JACOBY, 1896, Entomolog., 29, p. 7 (*Corynodes*) (Amami-Oshima).
 Distr.: Ryukyu Is.

subsp. *japonicus* JACOBY, 1896, s. str.

JACOBY, 1896, Entomolog., 29, p. 7 (*Corynodes*) (Amami-Oshima).
 Distr.: Ryukyu Is. (Amami-Oshima, Okinawa, Ishigaki, Iriomote, Yonaguni).

subsp. *umebayashii* KIMOTO, 1974

KIMOTO, 1974, Kontyû, Tokyo, 42, p. 281 (Okinoerabu).
 Distr.: Ryukyu Is. (Okinoerabu).

Genus *Chrysochus* CHEVROLAT, 1837 オオサルハムシ属

Chrysochus CHEVROLAT, 1837, in DEJEAN, Cat. Col. ed. 3, p. 413 (type-species: *Cryptocephalus pretiosus* SCHNEIDER, 1791 = *Cryptocephalus asclepiadeus* PALLAS, 1776).

C. chinensis BALY, 1859 オオサルハムシ

BALY, 1859, Ann. Mag. nat. Hist., (3), 4, p. 125 (N. China).
singularis LEFÈVRE, 1884, C. R. Soc. ent. Belg., 28, p. ccv (China).
goniostoma WEISE, 1889, Horae Soc. ent. Ross., 23, p. 592 (C. Mongolia).
cyclostoma WEISE, 1889, *ibid.*, p. 593 (Ordos).
curtus PIC, 1927, Bull. Soc. ent. France, 1927, p. 154 (Korea).
singularis var. *coerulescens* PIC, 1927, Mém. exot. Ent., 48, p. 25 (China).
 Distr.: E. Siberia, China, Korea, Japan (Honshu, Kyushu).

Genus *Pagria* LEFÈVRE, 1884 ヒメキバネサルハムシ属

Colposcelis DEJEAN, 1835, Cat. Col. ed. 2, p. 408 (nec DEJEAN, 1834, *ibid.*, p. 185, Tenebrionidae) (type-species: *Colaspis viridiaenea* GYLLENHAL, 1808).

Pagria LEFÈVRE, 1884, Bull. Soc. ent. France, (6), 4, p. lxxvii (type-species: *Pagria suturalis* LEFÈVRE, 1884).

P. signata (MOTSCHULSKY, 1858) ヒメキバネサルハムシ

MOTSCHULSKY, 1858, Étud. ent., 7, p. 110 (*Metachroma*) (Burma).
consimile BALY, 1874, Trans. ent. Soc. London, 1874, p. 168 (*Nodostoma*) (Nagasaki).
flavopustulatum BALY, 1874, *ibid.*, p. 169 (*Nodostoma*) (Nagasaki, Tsushima).
flavopustulata ab. *bicolor* WEISE, 1922, Philip. J. Sci., 21, p. 451 (Philippines).
signata var. *anceyi* PIC, 1929, Mém. exot. Ent., 53, p. 35 (Japan).
signata var. *rufithorax* PIC, 1929, *ibid.* (Tonkin).

signata var. *innotata* PIC, 1929, *ibid.* (Tonkin).

signata ab. *ussuriensis* ROUBAL, 1929, *Boll. Soc. ent. Ital.*, **61** (5-6), p. 98 (Vladivostock).

signata grandis CHÛJÔ, 1956, *Philip. J. Sci.*, **85** (1), p. 61 (Formosa).

Distr.: India, Burma, Vietnam, China, Siberia, Korea, Taiwan, Philippines, Indonesia, Japan (Honshu, Sado, Hachijô, Shikoku, Kyushu), Ryukyu Is. (Tokara, Amami-Oshima, Yoron, Miyako, Ishigaki, Iriomote), Ogasawara Is. (Naka-Iôtô), Micronesia (Guam).

Genus *Nodina* MOTSCHULSKY, 1853 チビサルハムシ属

Nodina MOTSCHULSKY, 1853, *Étud. ent.*, **7**, p. 108 (type-species: *Nodina pusilla* MOTSCHULSKY, 1858).

N. chalcosoma BALY, 1874 アオガネヒメサルハムシ

BALY, 1874, *Trans. ent. Soc. London*, **1874**, p. 170 (Nagasaki, China).

chalcosoma var. *rufofulva* CHÛJÔ, 1951, *Trans. Shikoku ent. Soc.*, **2**, p. 37 (Tokushima)

chalcosoma ab. *ao* NAKANE, 1963, *Kyoto Pref. Univ. Sci. Rept. (Nat. Sci. & Liv. Sci.)*, (A), **3** (5), p. 226 (Nakanoshima in Tokara Is.).

Distr.: S. China, Taiwan, Ryukyu Is. (Tokara, Amami-Oshima, Ishigaki, Iriomote, Yonaguni), Japan (Honshu, Shikoku, Kyushu, Yakushima, Tanegashima, Kuchinoerabu).

N. kraussi KIMOTO et GRESSITT, 1966 クラウスチビサルハムシ

KIMOTO & GRESSITT, 1966, *Pacif. Ins.*, **8**, p. 502 (Okinawa).

Distr.: Ryukyu Is. (Okinawa).

N. morimotoi KIMOTO et GRESSITT, 1966 モリモトチビサルハムシ

KIMOTO & GRESSITT, 1966, *Pacif. Ins.*, **8**, p. 503 (Okinawa).

Distr.: Ryukyu Is. (Okinawa).

N. sauteri CHÛJÔ, 1956 ザウターチビサルハムシ

CHÛJÔ, 1956, *Philip. J. Sci.*, **85** (1), p. 16 (Formosa).

Distr.: Taiwan, Ryukyu Is. (Okinawa).

Genus *Parascela* BALY, 1887 ヒメムクゲサルハムシ属

Parascela BALY, 1887, *J. Linn. Soc. London*, **14**, p. 252 (type-species: *Pseudocolaspis cribrata* SCHAUFUSS, 1871).

P. cribrata (SCHAUFUSS, 1871) ヒメムクゲサルハムシ

SCHAUFUSS, 1871, *Nonquam Otiosus*, **1**, p. 200 (*Pseudocolaspis*) (Hong Kong).

hirta CHEN, 1935, *Sinensia*, **6** (6), p. 769, fig. 1 (*Basilepta*) (Yunnan).

Distr.: S. China, Ryukyu Is. (Iriomote).

Genus *Basilepta* BALY, 1860 アオパネサルハムシ属

Basilepta BALY, 1860 (April), *J. Ent.*, **1**, p. 23 (type-species: *Basilepta longipes* BALY, 1860).

Nodostoma MOTSCHULSKY, 1860 (Decem.), *Schrenck's Reisen Amurl.*, **2**, p. 176 (type-species: *Nodostoma fulvipes* MOTSCHULSKY, 1860).

B. amamiense CHÛJÔ, 1957 ヨツモンサルハムシ

CHÛJÔ, 1957, *Kontyû, Tokyo*, **25**, p. 13, fig. 1 (Shinmura in Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima).

- B. balyi** (HAROLD, 1877) チャイロサルハムシ(ハンノキサルハムシ)
 HAROLD, 1877, Dtsch. ent. Z., **21**, p. 361 (*Nodostoma*) (Hakone).
japonicum JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 205 (*Nodostoma*) (Kiso).
otsukae OTSUKA, 1911, Konchû Sekai, **15** (2), p. 61 (*Nodostoma*) (Mt. Togakushi).
balyi yeso NAKANE, 1963, Kyoto Pref. Univ. Sci. Rept. (Nat. Sci & Liv. Sci.), (A),
 3 (5), 225 (Hokkaido).
balyi kurilensis L. MEDVEDEV, 1966, Forest Ent.-Fauna Kuril., Kamch. & Magad.,
 p. 40 (Kurul).
 Distr.: Japan (Hokkaido, Honshu, Sado, Hachijô, Shikoku, Kyushu), Kuril Is.
- B. borodinense** KIMOTO, 1979 ダイトウサルハムシ
 KIMOTO, 1979, Ent. Rev. Japan, **33**, p. 41 (Minami-Daitojima).
 Distr.: Ryukyu Is. (Minami-Daito).
- B. davidi** (LEFÈVRE, 1877) ダビッドサルハムシ
 LEFÈVRE, 1877, Ann. Soc. ent. France, (5), **7**, p. 157 (*Nodostoma*) (China).
atriventris PIC, 1933, Mél. exot. Ent., **61**, p. 7 (Tonkin).
atriventris var. *approximatus* PIC, 1933, *ibid.*, p. 8 (China).
okinawense CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, **25**, p. 77 (*Nodostoma*)
 (Naha in Okinawa).
insulanum CHÛJÔ, 1935, *ibid.*, p. 204 (*Nodostoma*) (Ishigaki).
 Distr.: Vietnam, China, Taiwan, Ryukyu Is. (Tokara, Amami-Oshima, Okinawa,
 Miyako, Ishigaki, Iriomote).
- B. fulvipes** (MOTSCHULSKY, 1860) アオバネサルハムシ
 MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 176, pl. 11, fig. 1 (*Nodostoma*)
 (W. Siberia: Dauria, Amur).
aeneipenne MOTSCHULSKY, 1860, *ibid.*, p. 177 (*Nodostoma*) (E. Siberia: Dauria).
rufotestaceum MOTSCHULSKY, 1860, *ibid.* (*Nodostoma*) (E. Siberia: Dauria).
atripes MOTSCHULSKY, 1860, *ibid.* (*Nodostoma*) (Japan).
chinense LEFÈVRE, 1877, Ann. Soc. ent. France, (5), **7**, p. 158 (*Nodostoma*) (Kiangsi).
fulvipes var. *coerulescens* WEISE, 1889, Horae. Soc. ent. Ross. **23**, p. 597 (*Nodostoma*)
 (Japan).
fulvipes var. *picicollis* WEISE, 1889, *ibid.* (*Nodostoma*) (Japan, Ordos).
bicoloripes PIC, 1930, Mél. exot. Ent., **56**, p. 7 (*Nodostoma*) (Yunnan).
bicoloripes var. *guerryi* PIC, 1930, *ibid.* (*Nodostoma*) (Yunnan).
 Distr.: Siberia, Mongolia, Korea, China, Taiwan, Japan (Hokkaido, Honshu, Awa-
 shima, Sado, Shikoku, Kyushu, Yakushima, Kuchinoerabu).
- B. hirayamai** (CHÛJÔ, 1935) ニセウスイロサルハムシ
 CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, **25**, p. 77 (*Nodostoma*) (Iriomote,
 Ishigaki, Naha in Okinawa).
 Distr.: Japan (Kyushu, Yakushima), Ryukyu Is. (Tokara, Amami-Oshima, Tokuno-
 shima, Okinawa, Ishigaki, Iriomote, Yonaguni), Taiwan.
- B. hirticollis** (BALY, 1874) ムナゲクロサルハムシ
 BALY, 1874, Trans. ent. Soc. London, **1874**, p. 167 (*Nodostoma*) (Nagasaki).
 Distr.: Japan (Honshu, Shikoku, Kyushu, Yakushima).

- B. modestum** (JACOBY, 1885) ルリサルハムシ
 JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 206 (*Nodostoma*) (Hitoyoshi).
 Distr.: S. China, Japan (Kyushu), Taiwan.
- B. pallidulum** (BALY, 1874) ウスイロサルハムシ(スギハムシ)
 BALY, 1874, Trans. ent. Soc. London, **1874**, p. 169 (*Nodostoma*) (Nagasaki).
laeviusculum WEISE, 1910, Verh. Naturf. Ver. Brünn, **48**, p. 34 (*Nodostoma*) (Tokyo).
 Distr.: Japan (Honshu, Hachijō, Shikoku, Kyushu), S. China.
- B. ruficollis** (JACOBY, 1885) ムネアカサルハムシ
 JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 205, pl. 11, fig. 10 (*Nodostoma*)
 (Nikko, Fukushima).
nigripennis PIC, 1928, Bull. Soc. zool. France, **43**, p. 379 (*Nodostoma*) (China).
ruficollis var. *melanicollis* CHŪJŌ, 1956, Philip. J. Sci., **85** (1), p. 37 (Formosa).
ruficollis ab. *kuromune* NAKANE, 1963, Kyoto Pref. Univ. Sci. Rept. (Nat. Sci. & Liv.
 Sci.), (A), (5), p. 225 (Mie).
 Distr.: W. China, Taiwan, Japan (Honshu, Kyushu).
- B. uenoi** NAKANE, 1958 ドウイロムナゲサルハムシ
hirticollis uenoi NAKANE, 1958, Sci. Rept. Saikyo Univ., (A), **2** (5), p. 305, fig. 8
 (Nakanoshima in Tokara Is.).
uenoi: NAKANE & KIMOTO, 1961, Bull. Osaka Mus. nat. Hist., **13**, p. 73 (Nakanoshima
 in Tokara Is.).
 Distr.: Ryukyu Is. (Tokara, Amami-Oshima, Okinawa).
- B. varicolor** (JACOBY, 1885) シイサルハムシ
 JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 751 (*Nodostoma*) (Hitoyoshi).
maebarai CHŪJŌ, 1956, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., **2** (31), p. 1,
 fig. 1 (Cape Sata in Kagoshima Pref.).
 Distr.: S. China, Japan (Hachijō, Shikoku, Kyushu, Yakushima).

Genus *Colasposoma* LAPORTE, 1833 イモサルハムシ属

- Colasposoma* LAPORTE, 1833, in Silbermann, Rev. Ent., **1**, p. 22 (type-species: *Colasposoma senegalense* LAPORTE, 1833).
Acis CHEVROLAT, 1837, in DEJEAN, Cat. Col. ed. 3, p. 411.
Thysbe THOMSON, 1858, Arch. Ent. **2**, p. 370.
- C. auripennis** (MOTSCHULSKY, 1860) オキナワイモサルハムシ
 MOTSCHULSKY, 1860, Schrenck's Reisen Amurl. **2**, p. 179 (*Acis*) (India).
viridicoeruleum MOTSCHULSKY, 1860, *ibid.*, p. 173 (*Acis*) (Hong Kong).
pulcherrimum BALY, 1864, Descrip. n. gen. & spec. Phytoph., p. 15 (India).
metallicum CLARK, 1865, Ann. Mag. nat. Hist. (3), **15**, p. 142 (Pulopenang).
mutabile BALY, 1867, Trans. ent. Soc. London, (3), **4** (2), p. 273 (Java, Borneo, Moluccas, Timor).
capitatum JACOBY, 1889, Ann. Mus. civ. Stor. nat. Genova, **27**, p. 178 (Burma).
affine LEFÈVRE, 1890, Nouv. Arch. Mus. Paris, (3), **2**, p. 195 (Laos, Luang-Prabang).
oberthuri JACOBY, 1896, Entomolog., **29**, p. 6 (Amami-Oshima).
oberthuri ab. *aurita*, *azurea*, *azureolineata*, *bryanti*, *chujoi*, *sauteri*, *violacea* PAPP,
 1946, Additam. Fauna Col., **4** (1), p. 7 ("locality not cited").
 Distr.: China, Taiwan, Vietnam, Laos, Burma, India, Malaya, Andaman Is., Ryukyu Is. (Tokara, Amami-Oshima, Okinawa, Miyako, Ishigaki, Iriomote).

C. dauricum MANNERHEIM, 1849 イモサルハムシMANNERHEIM, 1849, Bull. Soc. Imp. Nat. Moscou, **1**, p. 247 (E. Siberia: Dauria).*cyanea* MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 177, pl. 11, fig. 2 (E. Siberia: Dauria, Amur).

Distr.: E. Siberia, N. China, Korea, Japan (Honshu, Awa-shima, Sado, Hachijō, Shikoku, Kyushu, Tsushima).

Genus **Phytorus** JACOBY セスジサルハムシ属*Phytorus* JACOBY, 1884, Ann. Mus. civ. Stor. nat. Genova, **20**, p. 226 (type-species: *Phytorus dilatatus* Jacoby, 1884).**P. lineatus** WEISE, 1913 セスジサルハムシ

WEISE, 1913, Philip. J. Sci., D8 (3), p. 220 (Batan).

Distr.: Philippines, S. Mariana, Ogasawara Is. (Naka-iôtō).

Genus **Rhyparida** BALY, 1861 モモプトサルハムシ属*Rhyparida* BALY, 1861, J. Ent. **1**, p. 286 (type-species: *Rhyparida dimidata* BALY, 1861).*Marsaeus* CLARK, 1864, *ibid.*, **2**, p. 252.**R. sakisimensis** YUASA, 1930 モモプトサルハムシYUASA, 1930, Proc. Imp. Acad. Tokyo, **6** (7), p. 293, fig. 9 (Is. Taketomi, Ishigaki, Miyako).

Distr.: Ryukyu Is. (Taketomi, Miyako, Ishigaki, Iriomote), Taiwan, Lan Hsu.

Genus **Scelodonta** WESTWOOD, 1837 ドウガネサルハムシ属*Scelodonta* WESTWOOD, 1837, Proc. zool. Soc. London, **5**, p. 129 (type-species: *Scelodonta curculionoides* WESTWOOD, 1837).*Heteraspis* CHEVROLAT, 1837 (nec LECONTE), in DEJEAN, Cat. Col. ed. 3, p. 437.*Odontionopa* MOTSCHULSKY, 1866, Bull. Soc. Hist. Nat. Moscou, **2**, **29**, p. 408 (nec Erichson, 1842, nec Chevrolat, 1837).**S. lewisii** BALY, 1874 ドウガネサルハムシBALY, 1874, Trans. ent. Soc. London, **1874**, p. 165 (Nagasaki).*orientalis* LEFÈVRE, 1887, Bull. Soc. ent. France, (6), **7**, p. 56 (Chekiang).

Distr.: China, Taiwan, Japan (Honshu, Shikoku, Kyushu, Tsushima, Yakushima).

S. sauteri CHÛJÔ, 1938 シワドウガネサルハムシCHÛJÔ, 1938, Arb. Morph. Tax. Ent. Berlin-Dahlem, **5** (1), p. 29 (Formosa).

Distr.: Taiwan, Ryukyu Is. (Tokara), Japan (Kyushu).

Genus **Hyperaxis** GEMMINGER & HAROLD, 1863 クロオビカサハラハムシ属*Metaxis* BALY, 1863, J. Ent. **2**, p. 157 (nec de CHAUDOIR, 1850) (type-species: *Metaxis sellata* BALY, 1863).*Hyperaxis* GEMMINGER & HAROLD, 1874, Cat. Col. **11**, p. 3377 (n. n. for *Metaxis* BALY, 1863, nec de CHAUDOIR, 1850).**H. fasciata** (BALY, 1874) クロオビカサハラハムシBALY, 1874, Trans. ent. Soc. London, **1874**, p. 162 (*Demotina*) (Nagasaki).*taiwana* CHÛJÔ, 1956, Philip. J. Sci. **85** (1), p. 94 (Formosa).

Distr.: S. China, Taiwan, Japan (Honshu, Shikoku, Kyushu, Tsushima, Yakushima).

Genus *Demotina* BALY, 1863 カサハラハムシ属

Demotina BALY, 1863, J. Ent., 2, p. 158 (type-species: *Demotina bowringii* Baly, 1863).

D. major CHÛJÔ, 1958 オオアラゲサルハムシ

CHÛJÔ, 1958, Mem. Fac. Lib. Arts & Educ. 2 (64), p. 6 (as subsp. of *decoratella* CHÛJÔ) (Urasoe in Okinawa).

sasakawai NAKANE & KIMOTO, 1959, Sci. Rept. Saikyo Univ., (A), 3 (1), p. 67 (Konia in Amami-Oshima, Mt. Yonaha in Okinawa, Nakanoshima in Tokara Is.).

Distr.: Ryukyu Is. (Tokara, Amami-Oshima, Okinawa).

D. tuberosa CHEN, 1935 コブアラゲサルハムシ

CHEN, 1935, Sinensia, 6 (3), p. 354, fig. 33 (China, Tonkin).

japana OHNO, 1960, Trans. Shikoku ent. Soc., 6, p. 65 (Mt. Amagi in Shizuoka Pref., Kuroson in Kochi Pref., Mt. Izugatake and Mt. Bônoore in Saitama Pref.).

Distr.: Japan (Honshu, Shikoku, Kyushu), S. China, Vietnam.

D. fasciculata BALY, 1874 マダラカサハラハムシ

BALY, 1874, Trans. ent. Soc. London, 1874, p. 162 (Nagasaki).

Distr.: S. China, Japan (Honshu, Hachijô, Hachijô-kojima, Shikoku, Kyushu, Tsushima, Tanegashima, Yakushima).

D. decorata BALY, 1874 チビカサハラハムシ

BALY, 1874, Trans. ent. Soc. London, 1874, p. 163 (Nagasaki).

Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima, Tanegashima), Ryukyu Is. (Amami-Oshima).

D. modesta BALY, 1874 カサハラハムシ

BALY, 1874, Trans. ent. Soc. London, 1874, p. 164 (Nagasaki).

bipunctata JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 204 (Kobe).

piceonotata PIC, 1927, Bull. Soc. zool. France, 54, p. 45 (Yunnan).

elegans CHÛJÔ & SHIRÔZU, 1955, Sieboldia, Fukuoka, 1 (3) p. 239 (Ambo in Yakushima).

decorarella CHÛJÔ, 1956, Philip. J. Sci. 85 (1), p. 96 (Formosa).

inornata NAKANE, 1958, Sci. Rept. Saikyo Univ., (A), 2 (5), p. 304 (Yakushima).

elegans var. *futamon* NAKANE, 1958, *ibid.* (Kosugidani in Yakushima).

aurosquama CHÛJÔ, 1961, Pub. ent. Lab. Univ. Osaka Pref., 6, p. 84 (Amami-Oshima, Hachijô-jima).

Distr.: Taiwan, China, Ryukyu Is. (Amami-Oshima, Tokunoshima, Okinawa, Ishigaki, Iriomote), Japan (Honshu, Hachijô, Shikoku, Kyushu, Tanegashima, Yakushima), Korea.

Genus *Trichochrysea* BALY, 1861 トビサルハムシ属

Trichochrysea BALY, 1861, J. Ent., 1, p. 195 (type-species: *Trichochrysea mouhoti* BALY, 1865).

Bromius BALY, 1865, *ibid.*, 2, p. 439 (nec CHEVROLAT) (type-species: *Eumolpus hirta* FABRICIUS, 1801).

Heteraspis CHAPUIS, 1874, Gen. Col. 10, p. 284 (nec CHEVROLAT; nec LECONTE).

T. japana (MOTSCHULSKY, 1857) トビサルハムシ

MOTSCHULSKY, 1857, Étud. ent., 6, p. 37 (*Heteraspis*) (Japan).

Distr.: China, Korea, Japan, Ryukyu Is., Taiwan.

subsp. **japana** (MOTSCHULSKY, 1857), s. str.

MOTSCHULSKY, 1857, Étud. ent., 6, p. 37 (*Heteraspis*) (Japan).

Distr.: China, Korea, Japan (Honshu, Shikoku, Kyushu, Tsushima, Yakushima).

subsp. **okinawana** NAKANE, 1956

NAKANE, 1956, Sci. Rept. Saikyo Univ., (A), 2 (3), p. 171, pl. 2, fig. 35 (Okinawa, Amami-Oshima, Taiwan).

Distr.: Ryukyu Is. (Amami-Oshima, Okinawa, Ishigaki, Iriomote), Taiwan.

Genus *Lypesthes* BALY, 1863 ケブカサルハムシ属

Lypesthes BALY, 1863, J. Ent., 2, p. 152 (type-species: *Fidia atra* MOTSCHULSKY, 1860).

Leptotes BALY, 1863, *ibid.*, p. 158 (type-species: *Adoxus gracilicornis* BALY, 1861).

Talmonus FAIRMAIRE, 1889, Ann. Soc. ent. France, (6), 9, p. 71.

L. ater (MOTSCHULSKY, 1860) コフキササルハムシ (リングコフキハムシ)

MOTSCHULSKY, 1860, Étud. ent. 9, p. 22 (*Fidia*) (Japan).

pulverulentus JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 203, pl. 11, fig. 9 (*Leptotes*) (Oyama, Kiga, Oguma, Nikko).

farinosus FAIRMAIRE, 1889, Ann. Soc. ent. France, (6), 9, p. 71 (*Talmous*) (Moupin).

testaceipes PIC, 1928, Mém. exot. Ent., 52, p. 26 (Kioto).

ater fulvipes CHÛJÔ, 1954, Ins. Matsumurana, 18, p. 106 (Numatacho in Gumma, Mt. Myoken-zan in Osaka, Mt. Komagatake in Nagano, Mt. Otaki-san in Kagawa).

ater f. tibialis OHNO, 1958, J. Toyo Univ. 12, pp. 174, 176 (Urayamadani and Kama-buse-toge in Saitama, Syosenkyo in Yamanashi, Mt. Amami in Shizuoka).

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu), Korea, N. China.

L. fulvus (BALY, 1878) セアカケブカサルハムシ

BALY, 1878, J. Linn. Soc. London, Zool., 14, p. 250 (*Leptotes*) (China).

anger CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, 25, p. 204 (Naze in Amami-Oshima).

taiwanus CHÛJÔ, 1956, Philip. J. Sci., 85(1), p. 105 (Formosa).

Distr.: Japan (Shikoku, Kyushu, Yakushima, Kuchinoerabu), Ryukyu Is. (Tokara, Amami-Oshima, Okinoerabu, Okinawa, Miyako, Ishigaki), S. China, Taiwan.

L. itoi CHÛJÔ, 1954 スギケブカサルハムシ

CHÛJÔ, 1954, Ins. Matsumurana, 18, p. 103, fig. 1 (Miyazaki).

Distr.: Japan (Kyushu, Yakushima), Ryukyu Is. (Tokara).

L. japonicus OHNO, 1958 ニホンケブカサルハムシ

OHNO, 1958, J. Toyo Univ., 12, pp. 174, 179, pl. 6, figs. 12, 18 (Urasa, Misasa-mura and Tajimagahara in Saitama).

babai CHÛJÔ, 1958, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., 2 (58), p. 4 (Sumo in Tsushima).

Distr.: Japan (Honshu, Tsushima).

L. kiienensis OHNO, 1958 セスジケブカサルハムシ

OHNO, 1958, J. Toyo Univ., 12, p. 177, pl. 6, figs. 4, 9, 15 (Cape Shionomisaki in Wakayama Pref.).

kiienensis ab. immaculata OHNO, 1958, *ibid.*, pp. 175, 179 (Cape Shionomisaki in Wakayama Pref.).

Distr.: Japan (Honshu).

L. lewisi (BALY, 1878) ケブカサルハムシBALY, 1878, J. Linn. Soc. London, Zool., **14**, p. 251 (*Leptotes*) (China, Japan).

Distr.: China, Japan (Shikoku, Kyushu).

Genus **Bromius** CHEVROLAT ブドウサルハムシ属**Bromius** CHEVROLAT, 1837 (between May 1 and July 5), in DEJEAN, Cat. Col. ed. 3, p. 412 (type-species: *Chrysomela obscurus* LINNAEUS, 1758).*Adoxus* KIRBY, 1837 (October), Fauna Bor.-Amer. **4**, p. 209.*Eumolpus* REDTENBACHER, 1858, Fauna Austr. ed. 2, p. 893.**B. obscurus** (LINNAEUS, 1758) ブドウサルハムシLINNAEUS, 1758, Syst. Nat. ed. 10, p. 375 (*Chrysomela*) (Europe).*cochlearius* SAY, 1859, Long's Exp., 2, p. 196 (*Eumolpus*) (N. America).*nigroquadratus* DE GEER, 1775, Mem. Ins., 5, p. 336 (*Chrysomela*) (Europe).*violaosus* var. *epilobii* WEISE, 1882, Naturg. Ins. Dtschl., **6**, p. 295 (*Adoxus*) (Europe).*obscurus* var. *concinuus* WEISE, 1898, Arch. Naturg., **64**, p. 190 (Japan).*obscurus* var. *lewisi* WEISE, 1898, *ibid.* ("Yokohama").*obscurus japonicus* OHNO, 1960, Zool. Mag., Tokyo, **69**, p. 241 (Honshu).

Distr.: Europe, Siberia, Korea, Sakhalin, Japan (Hokkaido, Honshu).

Genus **Aoria** BALY, 1863 ケマルサルハムシ属**Aoria** BALY, 1863, J. Ent., **2**, p. 149 (type-species: *Adoxus nigripes* BALY, 1860).*Pseudaroriana* PIC, 1930, Mél. exot. Ent., **56**, p. 3.Subgenus **Osnaparis** FAIRMAIRE, 1889 アカチャサルハムシ亜属**Osnaparis** FAIRMAIRE, 1889, Ann. Soc. ent. France, **1889**, p. 72 (type-species: *Osnaparis nucea* FAIRMAIRE, 1889).**A. (O.) nucea** (FAIRMAIRE, 1889) アカチャサルハムシFAIRMAIRE, 1889, Ann. Soc. ent. France, (6), **9**, p. 72 (*Osnaparis*) (China).*taliana* PIC, 1935, Échang, **51**, p. 8 (China).

Distr.: China, Taiwan, Japan (Shikoku, Kyushu, Yakushima).

Genus **Xanthonia** BALY, 1863 キカサハラハムシ属**Xanthonia** BALY, 1863, J. Ent., **2**, p. 151 (type-species: *Xanthonia stevensi* BALY, 1863).**X. placida** BALY, 1874 キカサハラハムシBALY, 1874, Trans. ent. Soc. London, **1874**, p. 161 (Nagasaki).*placida hachijoensis* OHNO, 1960, Mushi, Fukuoka, **33** (9), p. 66 (Hachijojima, Hachijo-kojima).*placida hachijoensis* f. *obscura* OHNO, 1960, *ibid.* (Hachijo-jima).*placida hachijoensis* f. *scutellata* OHNO, 1960, *ibid.* (Hachijo-jima, Hachijo-kojima).

Distr.: Japan (Honshu, Sado, Hachijō, Hachijō-kojima, Shikoku, Kyushu, Tsushima), Ryukyu Is. (Tokara).

