

CHECK-LIST  
OF  
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 19

August 30, 1980

The Coleopterists' Association of Japan  
c/o Department of Zoology, National Science Museum (Nat. Hist.)  
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family CERAMBYCIDAE カミキリムシ科  
(LEPTURINAE)

By

Masao HAYASHI

Subfamily LEPTURINAE ハナカミキリ亜科

Tribe XYLOSTEINI ムカンハナカミキリ族

Genus *Caraphia* GAHAN, 1906 ケプトハナカミキリ属

*Caraphia* GAHAN, 1906, Fauna Brit. India Col., 1, Ceramb., p. 75 (type-species: *C. cribrata* GAHAN, 1906).

*Neosalpinia* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34, p. 303 (type-species: *N. lepturoides* MATSUSHITA, 1933, in Callidiopini, CERAMBYCINAE); HAYASHI, 1950, Ent. Rev. Japan, 5, p. 62 (LEPTURINAE).

*C. lepturoides* (MATSUSHITA, 1933) ケプトハナカミキリ

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34, p. 304, pl. 2, figs. 15, a, b (*Neosalpinia*) (Okinawa).

Distr.: Japan (SC & W. Honshu, Shikoku, Kyushu, Tsushima, Izu-Oshima, Nii-jima, Miyakejima, Hachijôjima, Tanegashima, Yakushima, Tokara-Nakanoshima, Suwanosejima), Ryukyu Is. (Amami-Oshima, Tokunoshima, Okinawa).

Tribe ENCYCLOPINI テツイロハナカミキリ族

Genus *Encyclops* NEWMAN, 1838 テツイロハナカミキリ属

*Encyclops* NEWMAN, 1838, Ent. Mag., 5, p. 392 (type-species: *E. pallipes* NEWMAN, 1838 = *Leptura caerulea* SAY, 1826).

*Microrhabdium* KRAATZ, 1879, Berl. ent. Z., 23, p. 118 (type-species: *M. macilentum* KRAATZ, 1879).

- E. olivacea* BATES, 1884 テツイロハナカミキリ (ウスアオホソカミキリ)  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 211, pl. 1, fig. 7 (*olivaceus*) (Chuzenji, Tochigi, Ohmine, Nara, Honshu).  
*jozanense* MATSUSHITA, 1933, J. Agr. Hokkaido Univ., **34**, p. 169, pl. 1, fig. 8 (*Micro-rhabdium*) (Jozankei, Hokkaido) (ミドリホソカミキリ).  
 Distr.: Japan (Hokkaido to Kyushu).

Tribe RHAGIINI ハイイロハナカミキリ族

Genus *Rhagium* FABRICIUS, 1775 ハイイロハナカミキリ属

- Rhagium* FABRICIUS, 1775, Syst. Ent., p. 182 (type-species: *Cerambyx inquisitor* LINNÉ, 1758).  
*Hargium* SAMOUELLE, 1819, Ent. useful Comp., p. 210 (type-species: *Cerambyx inquisitor* LINNÉ, 1758).  
*Allorhagium* KOLBE, 1884, Ent. Nachr., **10**, p. 270 (type-species: *Cerambyx inquisitor* LINNÉ, 1758).  
*R. japonicum* BATES, 1884 ハイイロハナカミキリ  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 209 (*inquisitor* var.) (Nyohozan, Nikko, Tochigi, Honshu) (part).  
 Distr.: Japan (Hokkaido, Honshu, Kurile Is.?).  
*R. heyrovskyi* PODANY, 1964 エゾハイイロハナカミキリ  
 PODANY, 1964, Acta Zool. Mex., **7** (2), p. 28, pl. 2, fig. 23; pl. 6, fig. 4 (Jozankei, Hokkaido).  
 Distr.: Japan (Hokkaido, Honshu).  
*R. pseudojaponicum* PODANY, 1964 ニセハイイロハナカミキリ  
 PODANY, 1964, Acta Zool. Mex., **7** (2), p. 26, pl. 6, fig. 3 (Tsushima).  
*japonicum* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 209 (*inquisitor* var.) (Oyayama, Kumamoto, Kyushu) (part).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima).

Genus *Enoploderes* FALDERMANN, 1837 ヒラヤマコブハナカミキリ属

- Enoploderes* FALDERMANN, 1837, Fauna Transcaucasus, **2**, p. 309 (type-species: *E. sanguineum* FALDERMANN, 1837).  
*Pyrotrichus* LECONTE, 1862, Proc. Acad. nat. Sci. Phila., **14**, p. 41 (type-species: *P. viticollis* LECONTE, 1862).  
*Pyrenoploderes* HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 12 (type-species: *Enoploderes bicolor* OHBAYASHI, 1941).  
*E. bicolor* OHBAYASHI, 1941 ヒラヤマコブハナカミキリ  
 OHBAYASHI, 1941, Mushi no Sekai, Tokyo, **4** (1/2), p. 1, pl. 4, fig. 1 (Ubuchi, Gumma, Honshu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Xenophyrama* BATES, 1884 ムナコブハナカミキリ属

- Xenophyrama* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 210 (type-species: *X. purpureum* BATES, 1884).  
*X. purpureum* BATES, 1884 ムナコブハナカミキリ  
 BATES, 1884, *ibid.*, p. 210, pl. 1, fig. 1 (Yuyama, Kumamoto, Kyushu).

Distr.: Japan (W. Honshu, Kyushu).

Tribe STENOCORINI カタビロハナカミキリ族

Genus *Stenocorus* FABRICIUS, 1775 ナガカタビロハナカミキリ属

*Stenocorus* FABRICIUS, 1775, Syst. Ent., p. 178 (type-species: *Leptura meridiana* LINNÉ, 1758) (part).

*Toxotus* DEJEAN, 1821, Cat. Col., ed. 1, p. 112 (type-species: *Leptura meridiana* LINNÉ, 1758).

*S. caeruleipennis* (BATES, 1873) フタコブルリハナカミキリ

BATES, 1873, Ann. Mag. nat. Hist., (4), 12, p. 193 (*Toxotus*) (Japan ?).

*galloisi* OHBAYASHI, 1961, Ent. Rev. Japan, 13, p. 16 (*T. caeruleipennis* BATES subsp.) (Yokohama).

Distr.: Japan (Hokkaido to Kyushu), Sakhalin.

Genus *Toxotinus* BATES, 1884 モモグロハナカミキリ属

*Toxotinus* BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 213 (type-species: *T. longicornis* BATES, 1884=*Toxotus minutus* var. *Reini* HEYDEN, 1879).

*T. reini* (HEYDEN, 1879) モモグロハナカミキリ

HEYDEN, 1879, Dtsch. ent. Z., p. 359 (*Toxotus minutus* var. *Reini*) (Osaka, Honshu).

*longicornis* BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 213, pl. 1, fig. 6 (Oyama, Kanagawa, Honshu).

Distr.: Japan (Hokkaido to Kyushu).

Genus *Pachyta* DEJEAN, 1821 カタビロハナカミキリ属

*Pachyta* DEJEAN, 1821, Cat. Col., ed. 1, p. 112 (type-species: *Leptura 8-maculata* FABRICIUS 1792=*L. 8-maculata* LINNÉ, 1758).

*P. lamed* (LINNÉ, 1758) カタキカタビロハナカミキリ

LINNÉ, 1758, Syst. Nat., ed. 10, p. 391 (*Cerambyx*) (Europe).

Distr.: Japan (High montane Hokkaido), Sakhalin, Korea, N. E. China, Mongolia, Siberia, N. & C. Europe, Alaska, Canada.

*P. erebia* BATES, 1884 キベリカタビロハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 212 (Chuzenji, Tochigi, Honshu).

Distr.: Japan.

subsp. *erebia* BATES, 1884, s. str.

BATES, 1884, *ibid.*, p. 212 (Chuzenji).

forma *tamanukii* HAYASHI et IGA, 1945

HAYASHI & IGA, 1945, Ins. World, 49 (569), p. 30 (*erebia* ab.) (Nikko).

Distr.: Japan (High montane Central Honshu).

subsp. *kusamai* HAYASHI, 1968

HAYASHI, 1955, Col. Ill. Ins. Japan, ed. 1, 1, p. 22 (*erebia* var.) (Ooi-Nikengoya, Shizuoka, Honshu); HAYASHI, 1968, Ent. Rev. Japan, 21, p. 11 (subsp.).

Distr.: Japan (S. Central Honshu, Kyushu).

Genus *Sachalinobia* JACOBSON, 1899 アラメハナカミキリ属

*Sachalinobia* JACOBSON, 1899, Ann. Mus. Zool. Acad. Imp. Sci. St. Pétersb., 4, p. 39 (type-species: *S. retata* JACOBSON, 1899=*Brachyta koltzei* HEYDEN, 1887).

*Pseudopachyta* SWAINE et HOPPING, 1928, Bull. nat. Mus. Canada, **52**, p. 16 (type-species: *Pachyta rugipennis* NEWMAN, 1844).

**S. rugipennis** (NEWMAN, 1844)

NEWMAN, 1844, Zool., p. 476 (*Pachyta*) (Canada); SWAINE et HOPPING, 1928, Bull. nat. Mus. Canada, **52**, pp. 14, 16 (*Pseudopachyta*); AURIVILLIUS, 1912, Col. Cat., **39**, p. 188 (*Evodinus*); GRESSITT, 1953, Pan-Pacific Ent., **29**, p. 207 (*Sachalinobia r. rugipennis*).

Distr.: Japan, Sakhalin, Korea, NE. China, E. Siberia, E. Canada, N. E. United States.

subsp. **koltzei** (HEYDEN, 1887) アラメハナカミキリ

HEYDEN, 1887, Dtsch. ent. Z., **31**, p. 304 (*Barachyta Koltzei*) (Nikolajevsk, Amur). *retata* JACOBSON, 1899, Ann. Mus. Zool. Acad. Imp. Sci. St. Pétersb., **4**, p. 40 (*Sachalinobia*) (Sakhalin).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Korea, NE. China, E. Siberia.

Genus **Brachyta** FAIRMAIRE, 1864 トホシハナカミキリ属

*Brachyta* FAIRMAIRE, 1864, in JACQUELIN DU VAL, Genera de Col. d'Europe, **4**, p. 185 (type-species: *Leptura interrogationis* LINNÉ, 1758).

**B. interrogationis** (LINNÉ, 1758) トホシハナカミキリ

LINNÉ, 1758, Syst. Nat., ed. 10, p. 398 (*Leptura*) (Sweden); MULSANT, 1863, Col. France Longic., ed. 2, p. 484 (*Pachyta*); AURIVILLIUS, 1912, Col. Cat., **39**, p. 186 (*Evodinus*).

Distr.: N. & High Montane Europe, Siberia, Mongolia, N. China, Korea, Sakhalin, Japan.

subsp. **punctata** (FALDERMANN, 1833) (トホシハナカミキリ)

FALDERMANN, 1833, Bull. Soc. Nat. Mosc., **6**, p. 67 (*Pachyta*) (E. Siberia); AURIVILLIUS, 1912, Col. Cat., **39**, p. 187 (*Evodinus punctatus*).

Distr.: Japan (Montane Hokkaido, High Montane Honshu, Rishirito), Kurile Is., Korea, NE. China, E. Siberia.

subsp. **sachalinensis** MATSUMURA, 1911 (カラフトトホシハナカミキリ)

MATSUMURA, 1911, J. Coll. Agr. Tôhoku Imp. Univ., **4** (1), p. 135 (var.) (Korsakoff, Sakhalin).

Distr.: Japan (Hokkaido), Sakhalin.

**B. bifasciata** (OLIVIER, 1792) フタスジカタビロハナカミキリ

OLIVIER, 1792, Encycl. Meth., **7**, p. 520 (*Leptura*) (E. Siberia) *daurica* GEBLER, 1817, Mem. Mosc., **5**, p. 329 (*Pachyta*).

Distr.: Japan, Siberia, NE. China, Mongolia, Korea, Sakhalin.

subsp. **japonica** (MATSUSHITA, 1933) (フタスジカタビロハナカミキリ)

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 179 (*Evodinus bifasciatus* var.) (Kamikochi, Nagano, Honshu).

Distr.: Japan (Honshu, Shikoku).

Genus **Evodinus** LECONTE, 1850 (クモマハナカミキリ属)

*Evodinus* LECONTE, 1850, J. Acad. nat. Sci. Phila., (2), **1**, p. 325 (type-species: *Leptura monticola* RANDALL, 1838).

*Evodinellus* PAVILSTSHIKOV, 1915, Rev. russe Ent., **15**, p. 355 (type-species: *Evodinus*

*borealis* GYLLENHAL, 1827) (*Evodinus* subg.) (ヒメクモマハナカミキリ亜属).

**E. borealis** (GYLLENHAL, 1827) クモマハナカミキリ

GYLLENHAL, 1827, Ins. Suec., **1** (4), p. 36 (*Leptura* ?) (Sweden).

Distr.: Japan (Hokkaido, Montane Honshu), Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.

Genus **Gaurotes** LECONTE, 1850 カラカネハナカミキリ属

*Gaurotes* LECONTE, 1850, J. Acad. nat. Sci. Phila., (2), **1**, p. 324 (type-species: *Rhagium cyanipennis* SAY, 1824).

*Paragaurotes* PLAVILSTSHIKOV, 1921, Rev. russe Ent., **17**, p. 1 (type-species: *Gaurotes?* (*Acmaeops*) *ussuriensis* BLESSIG, 1873) (*Gaurotes* subg.).

**G. doris** BATES, 1884 カラカネハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 212 (Chuzenji, Nyohozan, Tochigi, Honshu).

forma *svorovi* SEMENOV-TJAN-SHANSKIJ, 1914 (スボロフハナカミキリ)

SEMENOV-TJAN-SHANSKIJ, 1914, Rev. russe Ent., **14**, p. 16, 17 (*G. svorovi*) (Nemuro, Hokkaido).

forma *amagisana* OHBAYASHI, 1955 (アマギカラカネハナカミキリ)

OHBAYASHI, 1955, Akitu, Kyoto, **5**, p. 7 (*G. amagisana*) (Mt. Amagi, Shizuoka, Honshu).

Distr.: Japan (Hokkaido to Kyushu, Sado), Kurile Is.

Genus **Carillia** MULSANT, 1863 クビアカハナカミキリ属

*Carillia* MULSANT, 1863, Col. France Longic., ed. 2, p. 489 (type-species: *Leptura virginea* LINNÉ, 1758).

*Neogaurotes* PODANY, 1962, Mitt. Münch. ent. Ges., **52**, p. 225 (type-species: *Gaurotes virginea* LINNÉ, 1758) (*Gaurotes* subg.).

**C. atripennis** (MATSUSHITA, 1933) クビアカドウガネハナカミキリ

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 176 (Tokyo, Honshu).

forma *kurosawai* (HAYASHI, 1951) (赤紫色型)

HAYASHI, 1951, Ent. Rev. Japan, **5**, p. 75 (*Gaurotes* (*Gaurotes*) *kozhenikovi* PLAVILSTSHIKOV var.) (Mt. Azuma, Yamagata, Odayama, Fukushima, N. Honshu).

forma *tamaukii* (OHBAYASHI, 1949) (青藍色型)

OHBAYASHI, 1949, Shin Konchû, **2** (3), pl. no. 4 (*G. k.* PLAVILSTSHIKOV var.) (Hirugano, Gifu, C. Honshu).

forma *hayashii* (PODANY, 1962) (青紫色型)

PODANY, 1962, Mitt. Münch. ent. Ges., **52**, pp. 225, 230, pl. 4, fig. 5; pl. 5, fig. 12 (*G. (Neogaurotes) hayashii*) (Osaka, Honshu).

forma *viridipennis* (YOKOYAMA, 1969) (金緑色型)

YOKOYAMA, 1969, Ent. Rev. Japan, **21**, p. 58 (*Gaurotes*) (Mt. Kisokoma, Nagano).  
Distr.: Japan (Honshu, Shikoku, Kyushu, Sado).

**C. aureopurpurea** (HAYASHI, 1955) クビアカハナカミキリ

HAYASHI, 1955, Col. Ill. Ins. Japan, ed. 1, **1**, p. 24 (*Gaurotes*); HAYASHI, 1960, Nipponius, **1** (6), p. 5.

*virginea* LINNÉ subsp. *violacea* KANO, 1933 (nec DEGEER, 1775), Kontyû, Tokyo, **6**, p. 263 (*Gaurotes*) (Kamikochi, Nagano, Honshu).

Distr.: Japan (Central Honshu).

- C. otome* (OHBAYASHI, 1959) ヒメクビアカハナカミキリ (オトメクビアカハナカミキリ)  
 OHBAYASHI, 1959, Ent. Rev. Japan, **10**, p. 1, fig. 1 (*Gaurotus* (s. str.)) (Hirayu, Gifu,  
 Central Honshu).  
 Distr.: Japan (Honshu).

Genus *Lemula* BATES, 1884 ニセハムシハナカミキリ属

*Lemula* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 211 (type-species: *L. decipiens*  
 BATES, 1884).

- L. decipiens* BATES, 1884 キバネニセハムシハナカミキリ  
 BATES, 1884, *ibid.*, p. 212, pl. 1, fig. 5 (Miyanoshita, Kiga, Kanagawa, Honshu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Oki), E. Siberia.
- L. japonica* TAMANUKI, 1938 ニセハムシハナカミキリ  
 TAMANUKI, 1938, Ins. Matsum., **12**, p. 166, fig. 1 a (Kamikochi, Nagano, Honshu).  
 Distr.: Japan (High montane Central Honshu).
- L. rufithorax* PIC, 1901 ビックニセハムシハナカミキリ  
 PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 57 (Central Honshu).  
 forma *nigroguttata* HAYASHI, 1944  
 HAYASHI, 1944, Trans. Kansai ent. Soc., **14** (1), p. 27, fig. 2 (*L. decipiens* ab.) (Mt.  
 Kammuri, Hiroshima, W. Honshu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).
- L. nishimurai* SEKI, 1944 アカイロニセハムシハナカミキリ  
 SEKI, 1944, Ins. World, Gifu, **48** (558), p. 25 (Mt. Kifune, Kyoto, Honshu).  
*hiratai* HAYASHI, 1944, Trans. Kansai ent. Soc., **14** (1), pp. 23, 27, fig. 3 (Mt. Kammuri  
 Hiroshima, W. Honshu).  
 Distr.: Japan (Honshu, Shikoku).

Genus *Gnathacmaeops* LINSLEY et CHEMSAK, 1972 キタクニハナカミキリ属

- Gnathacmaeops* LINSLEY et CHEMSAK, 1972, Univ. Calif. Publ. ent., **69**, pp. 28, 135 (type-  
 species: *Leptura pratensis* LAICHTING, 1784).
- G. septentrionis* (C. G. THOMSON, 1866) キタクニハナカミキリ  
 THOMSON, C. G., 1866, Skand. Col., **8**, p. 61 (*Acmaeops*) (Scandinavia).  
 Distr.: Japan (Montane Hokkaido), Sakhalin, Korea, NE. China, N. Mongolia,  
 Siberia, N. & Montane Europe.

Genus *Dinoptera* MULSANT, 1863 ヒナルリハナカミキリ属

- Dinoptera* MULSANT, 1863, Col. France Longic., ed. 2, p. 494 (type-species: *Leptura*  
*collaris* LINNÉ, 1758) (*Acmaeops* subg.)
- D. minuta* (GEBLER, 1832) ヒナルリハナカミキリ  
 GEBLER, 1832, Mem. Soc. nat. Mosc., **8**, p. 69 (*Pachyta*) (Nerchinsk, E. Siberia).  
*criocerina* BATES, 1873, Ann. Mag. nat. Hist., (4), **12**, p. 19 (Aomori, N. Honshu).  
*japonica* PIC, 1907, Mat. Longic., **6** (2), p. 20 (*Acmaeops* (*Dinoptera*)) (Sado, Japan)  
 (サドヒメハナカミキリ).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Sado, Oki, Tsushima), Sakhalin, Korea,  
 NE. China, E. Siberia.

Genus *Macropidonia* PIC, 1901 アカムネハナカミキリ属

*Sieversia* GANGLBAUER, 1886, Horae Soc. ent. Ross., **20**, p. 134 (type-species: *S. bicolor* GANGLBAUER, 1886) (シーベルス属)

*Macropidonia* PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 340 (type-species: *M. ruficollis* PIC, 1901).

*Sivana* STRAND, 1942, Fol. Zool. Hydrob., **11**, p. 391 (new name for *Sieversia* GANGLBAUER, 1866, nec KOBERT, 1880 for Mollusca).

*M. ruficollis* PIC, 1901 アカムネハナカミキリ

PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 340 (Chuzenji, Tochigi, Honshu).

Distr.: Japan (Central Honshu).

Genus *Pseudosieversia* PIC, 1902 ヒメコブハナカミキリ属

*Pseudosieversia* PIC, 1902, Échange, **18**, p. 19 (type-species: *Pidonia rufa* KRAATZ, 1879).

*P. japonica* (OHYAYASHI, 1937) チャイロヒメコブハナカミキリ

OHYAYASHI, 1937, Nippon no Kôchû, **1**, p. 5 (*Microrhabdium japonicum*) (Tokugotoge Nagano, Honshu).

Distr.: Japan (Montane Central Honshu).

*P. shikokensis* HAYASHI, 1959 シコクヒメコブハナカミキリ

HAYASHI, 1959, Ent. Rev. Japan, **10**, p. 55 (Kotsuzan, Mt. Tsurugi, Tokushima, Shikoku).

Distr.: Japan (Honshu?, Shikoku).

*P. amanoi* HAYASHI, 1971 キュウシュウヒメコブハナカミキリ

HAYASHI, 1971, Ent. Rev. Japan, **23**, p. 2 (Mt. Sobu, Ooita, Kyushu).

Distr.: Japan (Kyushu).

Genus *Pidonia* MULSANT, 1863 ヒメハナカミキリ属

*Pidonia* MULSANT, 1863, Col. France Longic., ed. 2, p. 570 (type-species: *Leptura lurida* FABRICIUS, 1792).

*Pseudopidonia* PIC, 1900, Échange, **16**, p. 81 (type-species: *P. amurensis* PIC, 1900) (トウアヒメハナカミキリ属).

Subgenus *Pidonia* s. str.

*P. (P.) yamato* HAYASHI et MIZUNO, 1953 ヤマトヒメハナカミキリ

HAYASHI & MIZUNO, 1953, Ent. Rev. Japan, **6**, p. 38, pl. 8, figs. 1, 2 (Mt. Kasuga, Nara; Ohsugidani, Mie, Honshu; Mt. Asahimaru, Tokushima, Shikoku).

Distr.: Japan.

subsp. *yamato* HAYASHI et MIZUNO, 1953, s. str.

HAYASHI & MIZUNO, 1953, *ibid.*, p. 38, pl. 8, figs. 1, 2 (Mt. Kasuga, Nara; Ohsugidani Mie, Honshu; Mt. Asahimaru, Tokushima, Shikoku).

Dist.: Japan (Honshu, Shikoku).

subsp. *higeshiro* YAMAWAKI, 1977

YAMAWAKI, 1977, Ent. Rev. Japan, **30**, p. 61, fig. (Mt. Shiraiwa, Motoyashiki, Miyazaki; Fukakura, Mt. Hiko, Fukuoka, Kyushu).

Distr.: Japan (Kyushu).

*P. (P.) nagayamai* HAYASHI, 1968 ナガヤマヒメハナカミキリ

HAYASHI, 1968, Bull. Osaka Jonan Women's Jr. Coll., **3**, pp. 35, 37, pl. 1, figs. 9, 10;

pl. 4, fig. 7; pl. 7, fig. 5; pl. 9, fig. 4 (Mt. Ohshiomori, Kassi pass, Fukushima, Honshu).

Distr.: Japan (NC. Honshu).

**P. (P.) obscurior** PIC, 1901 シラネヒメハナカミキリ

PIC, 1901, Bull. Mus. Hist. nat. Paris, 7, p. 338 (*P. (Pseudopidonia) semiobscura* var.) (near Nikko).

*shiranensis* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34 (2), p. 191 (*Pseudopidonia ohbayashii* ab.) (Mt. Shirane, Tochigi, Honshu).

*bisignata* PIC, 1935, Échange, 51, p. 12 (*Pseudopidonia*) (Japan) (フタモンヒメハナカミキリ).

Distr.: Japan.

subsp. *obscurior* PIC, 1901, s. str.

PIC, 1901, *ibid.*, p. 338 (*P. (Pseudopidonia) semiobscura* var.) (near Nikko).

Distr.: Japan (NC. Honshu).

subsp. *hakusana* OHBAYASHI et HAYASHI, 1960

OHBAYASHI & HAYASHI, 1960, Ent. Rev. Japan, 11, p. 14, pl. 2, figs. 9, 10; pl. 3, figs. 8, 16 (Hakusan, Ishikawa, Toyama, Honshu).

Distr.: Japan (NW. Central Honshu).

**P. (P.) yokoyamai** HAYASHI, 1968 ヨコヤマヒメハナカミキリ

HAYASHI, 1968, Bull. Osaka Jonan Women's Jr. Coll., 3, pp. 34, 40, pl. 1, figs. 16, 17; pl. 4, fig. 10; pl. 7, fig. 8; pl. 9, fig. 9 (Mt. Yatsugatake, Nagano, Oze, Gumma, Honshu).

Distr.: Japan (N. Central Honshu).

**P. (P.) semiobscura** PIC, 1901 ホソガタヒメハナカミキリ

PIC, 1901, Bull. Mus. Hist. nat. Paris, 7, p. 338 (*P. (Pseudopidonia)*) (near Nikko) ♀. *nipponensis* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34, p. 190 (*Pseudopidonia*) (Kamikochi, Nagano, Honshu) ♂ (ホソガタヒメハナカミキリ).

*rufoscuteolata* MATSUSHITA, 1933, *ibid.*, p. 192, pl. 2, fig. 9 (*Pseudopidonia*) (Tokugotoge, Nagano, Honshu) ♀.

Distr.: Japan (Central Honshu).

**P. (P.) matsushitai** (OHBAYASHI, 1958) ヘリモンヒメハナカミキリ

OHBAYASHI, 1958, Ent. Rev. Japan, 9, p. 62 (new name for *ruficollis* MATSUSHITA, 1933).

*ruficollis* MATSUSHITA, 1933 (nec SAY, 1824), J. Fac. Agr. Hokkaido Univ., 34, p. 193 (*Pseudopidonia*) (Kamikochi, Nagano, Honshu) (part).

Distr.: Japan (C. Honshu).

**P. (P.) limbaticollis** (PIC, 1902) ニッコウヒメハナカミキリ

PIC, 1902, Mat. Longic., 4 (1), p. 27 (*Pseudopidonia harmandi* var.).

*nikkoensis* OHBAYASHI et HAYASHI, 1957, Ent. Rev. Japan, 8, p. 7 (Okunikko, Tochigi, Honshu).

Distr.: Japan.

subsp. *limbaticollis* (PIC, 1902), s. str.

PIC, 1902, *ibid.*, p. 27 (*Pseudopidonia harmandi* var.).

Distr.: Japan (C. Honshu).

subsp. *ohbayashii* (MATSUSHITA, 1933) (オオバヤシヒメハナカミキリ)

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34, p. 191 (*Pseudopidonia*) (Tokugo-



- toge, Nagano, Honshu) ♀.  
*tokugoana* MATSUSHITA, 1933, *ibid.*, p. 194 (*Pseudopidonia*) (Tokugotoge, Nagano, Honshu) ♂ (トクゴウヒメハナカミキリ).  
 Distr.: Japan (C. Honshu).
- subsp. *breuningi* OHBAYASHI et HAYASHI, 1957 (ブロイニングヒメハナカミキリ)  
 OHBAYASHI & HAYASHI, 1957, *Ent. Rev. Japan*, **8**, p. 5, pl. 2, figs. 1-4 (Mt. Ishizuchi, Mt. Akaishi, Ehime; Mt. Tsurugi, Tokushima, Shikoku).  
 Distr.: Japan (Montane Shikoku).
- P. (P.) leucanthophila* KUBOKI, 1978 マホロバヒメハナカミキリ  
 KUBOKI, 1978, *Elytra*, Tokyo, **6** (1), p. 33, photo., figs. 1-4 (Mt. Higashiakaishi, Ehime, Shikoku).  
 Distr.: Japan (Shikoku).
- P. (P.) signifera* (BATES, 1884) ナガバヒメハナカミキリ  
 BATES, 1884, *J. Linn. Soc. London, Zool.*, **18**, p. 215 (*Grammoptera*) (Nikko, Tochigi, Oyama, Kanagawa, Honshu; Hitoyoshi, Kumamoto, Kyushu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Sado, Oki, Tsushima).
- P. (P.) japonica* OHBAYASHI et HAYASHI, 1960 ニッポンヒメハナカミキリ (トウホクヒメハナカミキリ)  
 OHBAYASHI & HAYASHI, 1960, *Ent. Rev. Japan*, **11**, p. 1, pl. 2, fig. 11; pl. 3, figs. 9, 17 (Mt. Hakkoda, Aomori, Akita, Iwate, Yamagata, N. Honshu).  
 Distr.: Japan (N. Honshu).
- P. (P.) shikokuana* HAYASHI, 1968 シコクヒメハナカミキリ  
 HAYASHI, 1968, *Bull. Osaka Jonan Women's Jr. Coll.*, **3**, p. 53 (new name for *shikokensis* OHBAYASHI et HAYASHI, 1957)  
*shikokensis* OHBAYASHI et HAYASHI, 1957 (nec *P. discoidalis* subsp. *shikokensis*, CHÛJÔ et HAYASHI, 1951), *Ent. Rev. Japan*, **8**, p. 6, pl. 10, fig. 14 (Matsubagawa, Kochi; Mt. Tsurugi, Mt. Kotsuzan, Tokushima; Amogo Valley, Ehime, Shikoku).  
 Distr.: Japan (Shikoku, Kyushu).
- P. (P.) kyushuensis* YAMAWAKI, 1959 キュウシュウヒメハナカミキリ (ナガホソヒメハナカミキリ)  
 YAMAWAKI, 1959, *Kitakyushu no Konchu*, **6**, p. 85, figs. (Mt. Sobo, Mt. Kujuu, Ooita Kyushu).  
 Distr.: Japan (Kyushu).
- P. (P.) mutata* (BATES, 1884) ヒメハナカミキリ  
 BATES, 1884, *J. Linn. Soc. London, Zool.*, **18**, p. 215 (*Grammoptera signifera* var.) (Nikko, Oyama, Hitoyoshi).  
*chairo* TAMANUKI, 1942, *Fauna Nipponica Ceramb. Lept.* **2**, p. 38, fig. 80 (*P. (Pseudopidonia)*) (Mt. Ohmine, Nara, Honshu) (ヤノヒメハナカミキリ).  
*yanoi*: HAYASHI, 1968 (nec TAMANUKI), *Bull. Osaka Jonan Women's Jr. Coll.*, **3**, p. 54 (nom. nud.).  
*alpina* HAYASHI, 1968, *ibid.*, p. 54, pl. 5, fig. 19 (Mt. Ontake, Nagano, Honshu).  
 Distr.: Japan.
- subsp. *mutata* BATES, 1884, s. str.  
 BATES, 1884, *J. Linn. Soc. London, Zool.*, **18**, p. 215 (*Grammoptera signifera* var.).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).

- subsp. *pallida* OHBAYASHI et HAYASHI, 1960  
 OHBAYASHI & HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 14, pl. 3, figs. 7, 15 (Mt. Warusawadake, Mt. Hijiri, Nagano; Marunuma, Gumma, Honshu).  
 Distr.: Japan (High Montane Central Honshu).
- P. (P.) maculithorax* PIC, 1901 カクムネヒメハナカミキリ  
 PIC, 1901, Bull., Mus. Hist. nat. Paris, **7**, p. 338 (Central Japan).  
*bouvieri* PIC, 1901, *ibid.*, p. 339 (Central Japan) (ブービーヒメハナカミキリ).  
*orientalis* MATSUSHITA, 1933, Ins. Matsum. **7**, p. 103, figs. 1, a, b (Kamikochi, Nagano Honshu).  
*rufithorax* PIC, 1956, Longicornia **3**, p. 647, fig. 1 (*Pseudopidonia*) (Nakabusaonsen, Nagano, Honshu).  
*tohokensis* HAYASHI, 1968, Bull. Osaka Jonan Women's Jr. Coll., **3**, p. 56, pl. 2, fig. 35 (forma) (Tanemakiyama, Mt. Azuma, Fukushima, Honshu).  
 Distr.: Japan (Honshu).
- P. (P.) sylvicola* KUBOKI, 1977 ミヤマヒメハナカミキリ  
 KUBOKI, 1977, Kontyû, Tokyo, **45**, p. 264, figs. 1, 3, 4, 8, 9 (Shimashima, Nagano; Shizuoka, Yamanashi, Gumma, Kanagawa, Honshu).  
 Distr.: Japan (Central Honshu).
- P. (P.) discoidalis* (PIC, 1901) キベリクロヒメハナカミキリ  
 PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 58 (*Grammoptera grallatrix* var.) (near Nikko, Tochigi, Honshu).  
*yatsugataensis* HAYASHI, 1969, Bull. Osaka Jonan Women's Jr. Coll., **4**, p. 74 (var.) (Mt. Yatsugatake, Nagano, Honshu).  
 Distr.: Japan.  
 subsp. *discoidalis* (PIC, 1901), s. str.  
 PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 58 (*Grammoptera grallatrix* var.) (near Nikko, Tochigi, Honshu).  
 Distr.: Japan (Honshu).
- subsp. *shikokensis* CHÛJÔ et HAYASHI, 1951  
 CHÛJÔ & HAYASHI, 1951, Bull. Takarazuka Ins., (78), p. 4 (*P. (Pseudopidonia) discoidalis* forma) (Mt. Tsurugi, Tokushima, Shikoku).  
 Distr.: Japan (Kii Peninsula, Honshu, Shikoku, Kyushu?).
- subsp. *amanoi* HAYASHI, 1969  
 HAYASHI, 1969, Bull. Osaka Jonan Women's Jr. Coll., **4**, pp. 71, 75, pl. 2, fig. 45; pl. 7, fig. 29 (Mt. Hiko Fukuoka, Kyushu).  
 Distr.: Japan (Kyushu).
- P. (P.) muneaka* TAMANUKI, 1942 ムネアカヒメハナカミキリ  
 TAMANUKI, 1942, Fauna Nipponica Ceramb. **2**, Lept., p. 41 (*P. (Pseudopidonia) discoidalis* ab.).  
*ruficollis* MATSUSHITA, 1938 (nec PIC, 1901), Ins. Matsum., **12**, p. 99 (*Macropidonia*) (Kochi Shikoku).  
*tristiculoides* OHBAYASHI, 1959, Ent. Rev. Japan, **10**, p. 1 (Mt. Hakusan, Ishikawa; Mt. Daisen, Tottori, Honshu).  
*tamanukii* HAYASHI, 1969, Bull. Osaka Jonan Women's Jr. Coll., **4**, pp. 71, 77, pl. 2, figs. 48, 49; pl. 8, figs. 35, 36 (Kamikochi, Nagano, Gifu, Honshu).  
 Distr.: Japan.

- subsp. *muneaka* TAMANUKI, 1942, s. str.  
 TAMANUKI, 1942, *ibid.*, p. 41 (*P. (Pseudopidonia) discoidalis* ab.).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Oki).
- subsp. *amagisana* HAYASHI, 1953 (アマギキベリクロヒメハナカミキリ)  
 HAYASHI, 1953, *Ent. Rev. Japan*, **6**, p. 39 (*P. (Pseudopidonia) discoidalis* f.) (Mt. Amagi, Shizuoka, Honshu).  
 Distr.: Japan (S. Central Honshu).
- P. (P.) grallatrix* (BATES, 1884) オオヒメハナカミキリ  
 BATES, 1884, *J. Linn. Soc. London, Zool.*, **18**, p. 214 (*Grammoptera*) (Nikko, Tochigi, Honshu).  
*harmandi* PIC, 1901, *Bull. Mus. Hist. nat. Paris*, **7**, p. 58 (*P. (? Pseudopidonia)*) (Central Japan).  
*pallidisuturalis* OHBAYASHI, 1956, *Akitsu, Kyoto*, **5**, p. 7 (*Pseudopidonia grallatrix* morph.) (Mt. Hooman, Fukuoka; Mt. Sobo, Ooita, Kyushu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).
- P. (P.) gibbicollis* (BLESSIG, 1873) ミスジヒメハナカミキリ  
 BLESSIG, 1873, *Horae Soc. ent. Ross.*, **9**, p. 258 (*Leptura (Anoplodera)*) (Amur, Ussuri).  
 Distr.: Japan (Tsushima), Korea, NE. China, Ussuri, Amur.
- P. (P.) signata* MATSUSHITA, 1933 フイリヒメハナカミキリ  
 MATSUSHITA, 1933, *J. Fac. Agr. Hokkaido Univ.*, **34**, p. 185, pl. 2, fig. 4 (Tokugotoge, Nagano, Honshu).  
*uenoi* HAYASHI, 1969, *Bull. Osaka Jonan Women's Jr. Coll.*, **4**, p. 85, pl. 3, fig. 60 (ab.) (Mt. Ontake, Nagano, Honshu).  
 Distr.: Japan (High Montane Honshu).
- P. (P.) tsukamotoi* MIZUNO, 1978 タカネヒメハナカミキリ  
 MIZUNO, 1978, *Ent. Rev. Japan*, **32**, p. 39, pl. 1, figs. 1-9 (Mt. Senjo, Nagano, Honshu).  
 Distr.: Japan (High Montane Honshu).
- P. (P.) insuturata* PIC, 1901 ヨコモンヒメハナカミキリ  
 PIC, 1901, *Bull. Mus. Hist. nat. Paris*, **7**, p. 58 (*P. (Pseudopidonia)*) (near Nikko, Tochigi, Honshu).  
 Distr.: Japan.
- subsp. *insuturata* PIC, 1901, s. str.  
 PIC, 1901, *ibid.*, p. 58 (*P. (Pseudopidonia)*) (near Nikko, Tochigi, Honshu).  
 Distr.: Japan (Montane Honshu).
- subsp. *hylophila* KUBOKI, 1977 (イヨヒメハナカミキリ)  
 KUBOKI, 1977, *Kontyû, Tokyo*, **45**, p. 67, fig. 2 (*P. hylophila*) (Mt. Ishizuchi, Ehime, Shikoku; Ooita, Fukuoka, Kumamoto, Kyushu).  
 Distr.: Japan (Shikoku, Kyushu).
- forma *ohdaisana* MIZUNO, 1978  
 MIZUNO, 1978, *Ent. Rev. Japan*, **32**, p. 39, fig. 1 (*hylophila* subsp.) (Mt. Ohdaigahara, Nara, Honshu).  
 Distr.: Japan (Kii Peninsula, Honshu).
- subsp. *hamadryas* KUBOKI, 1977 (ミチノクヒメハナカミキリ)  
 KUBOKI, 1977, *Kontyû, Tokyo*, **45**, p. 268, fig. 11 (*P. hamadryas*) (Mt. Zao, Yamagata, Honshu).  
 Distr.: Japan (N. Honshu).

- subsp. *hayashii* KOIKE, 1971 (クロヨコモモンヒメハナカミキリ)  
 KOIKE, 1971, Ent. Rev. Japan, **23**, p. 25, pl. 1, figs. 1-7 (Mt. Naeba, Niigata, Honshu).  
 Distr.: Japan (N. Central Honshu).
- P. (P.) masakii* TAMANUKI, 1942 ムネアカヨコモモンヒメハナカミキリ  
 TAMANUKI, 1942, Fauna Nipponica Ceramb. **2** Lept., p. 50 (*P. (Pseudopidonia) insuturata* ab.).  
*ruficollis* MATSUSHITA, 1933 (nec SAY, 1824), J. Fac. Agr. Hokkaido Univ., **34**, p. 193 (*Pseudopidonia*) (part).  
*hiurai* OHBAYASHI et HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 14, pl. 2, figs. 3, 4; pl. 3, figs. 4, 11, 19 (forma) (Mt. Hakkoda, Towada, Aomori; Mt. Chokai, Akita, Honshu).  
*lepesmei* PIC, 1956, Longicornia **3**, p. 647 (*Pseudopidonia insuturata* var.) (Nakabusa-onsen, Nagano, Honshu).  
 Distr.: Japan (Central & Northern Honshu).
- P. (P.) simillima* OHBAYASHI et HAYASHI, 1960 ニセヨコモモンヒメハナカミキリ  
 OHBAYASHI & HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 14, pl. 2, figs. 5, 6; pl. 3, figs. 5, 12, 19 (Honshu, Shikoku, Kyushu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).
- P. (P.) amentata* (BATES, 1884) セスジヒメハナカミキリ  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 215 (*Grammoptera*) (Miyanoshta, Oyama, Kanagawa, Suyama, Shizuoka, Chuzenji, Tochigi, Honshu).  
*staudingeri* PIC, 1901, Échange, **17** (194), p. 12.  
 Distr.: Japan.
- subsp. *amentata* (BATES, 1884), s. str.  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 215 (*Grammoptera*) (Miyanoshta, Oyama, Kanagawa, Suyama, Shizuoka, Chuzenji, Tochigi, Honshu).  
 Distr.: Japan (C. & S. Honshu, Shikoku, Kyushu, Sado, Oki).
- subsp. *kurosawai* OHBAYASHI et HAYASHI, 1960  
 OHBAYASHI & HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 13, pl. 2, figs. 7, 8; pl. 3, figs. 2, 3 (Rausu, Hokkaido, Aomori, Iwate, Akita, Yamagata, Fukushima).  
 Distr.: Japan (Hokkaido, N. Honshu), Kurile Is.
- subsp. *lyra* KUBOKI et SUZUKI, 1978 (コトヒメハナカミキリ)  
 KUBOKI & SUZUKI, 1978, Kontyû, Tokyo, **46**, p. 297 figs. 1-4 (*P. lyra*) (Mt. Tanzawa, Kanagawa, Tokyo, Shizuoka, Yamanashi, S. Central Honshu).  
 Distr.: Japan (S. C. Honshu).
- P. (P.) miwai* (MATSUSHITA, 1933) ミワヒメハナカミキリ  
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 190 (*Pseudopidonia*) (Towada, Aomori, Honshu).  
 Distr.: Japan.
- subsp. *miwai* (MATSUSHITA, 1933), s. str.  
 MATSUSHITA, 1933, *ibid.*, p. 190 (*Pseudopidonia*) (Towada, Aomori, Honshu).  
 Distr.: Japan (N. Honshu).
- subsp. *fujisana* OBIKA et KUSAMA, 1971 (フジヒメハナカミキリ)  
 OBIKA & KUSAMA, 1971, Annot. Zool. Japonensis, **44**, p. 233, figs. 1, a, b (*P. fujisana*) (Mt. Fuji, Shizuoka, Honshu).  
 Distr.: Japan (S. C. Honshu).

- subsp. *approximata* KUBOKI, 1977 (トサヒメハナカミキリ) (オモテミワヒメハナカミキリ)  
 KUBOKI, 1977, Kontyû, Tokyo, **45**, p. 65, fig. 1 (Mt. Irazu, Kochi, Ehime, Shikoku;  
 Kagoshima, Kumamoto, Fukuoka, Kyushu; Wakayama, Mie, Yamanashi,  
 Honshu).  
 Distr.: Japan (SW. Honshu, Shikoku, Kyushu).
- P. (P.) oyamae* (OYAMA, 1908) オヤマヒメハナカミキリ  
 OYAMA, 1908, Ins. World, Gifu, **12** (135), p. 19, fig. (*Grammoptera*).  
*ruficollis* MATSUSHITA (nec SAY, 1824), 1933, J. Fac. Agr. Hokkaido Univ., **34**, p.  
 193 (*Pseudopidonia*) (Kamikochi, Nagano, Honshu) (part.).  
*quadrinaculata* MATSUSHITA, 1933, *ibid.*, p. 192, pl. 1, fig. 12 (*Pseudopidonia*) (Kami-  
 kochi, Nagano, Honshu) (マツシタノミハナカミキリ).  
 Distr.: Japan (Montane C. Honshu).
- P. (P.) chujoi* OHBAYASHI et HAYASHI, 1960 チュウジョウヒメハナカミキリ  
 OHBAYASHI & HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 13, pl. 2, figs. 1, 2; pl. 3, figs. 1,  
 13, 20 (Mt. Tsurugi, Tokushima, Mt. Ishizuchi, Ehime, Shikoku).  
 Distr.: Japan (Montane Shikoku).

Subgenus *Omphalodera* SOLSKY, 1873

- Omphalodera* SOLSKY, 1873, Horae Soc. ent. Ross., **9**, p. 244 (type-species: *O. puziloi*  
 SOLSKY, 1873).
- P. (O.) puziloi* (SOLSKY, 1873) フタオビノミハナカミキリ  
 SOLSKY, 1873, *ibid.*, p. 245, pl. 8, fig. 5 (Suyfun, Amur).  
*flaviventris* BATES, 1884, J. Linn. Soc. London, Zool., **18**, 0. 212 (var.) (Oyama, Miya-  
 noshita, Nikko, Suyama, Honshu).  
 Distr.: Japan (Hokkaido to Kyushu, Oki, Tsushima), Sakhalin, Korea, E. Siberia.
- P. (O.) testacea* (MATSUSHITA, 1933) ニセフタオビノミハナカミキリ  
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 184 (*O. puziloi* var.) (Tokugo-  
 toge, Kamikochi, Nagano, Honshu).  
 Distr.: Japan (Montane C. Honshu).

Subgenus *Mumon* HAYASHI, 1968

- Mumon* HAYASHI, 1968, Bull. Osaka Jonan Women's Jr. Coll., **3**, p. 15 (type-species:  
*Grammoptera debilis* KRAATZ, 1879) (*Pidonia* subg.).
- P. (M.) debilis* (KRAATZ, 1879) チャイロヒメハナカミキリ  
 KRAATZ, 1879, Dtsch. ent. Z., **23**, p. 104 (*Grammoptera*) (Amur).  
*aegrota* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 214 (Nikko, Tochigi, Hon-  
 shu; Hitoyoshi, Kumamoto, Kyushu, etc.) (*Grammoptera*).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Sado, Oki, Tsushima), Sakhalin, E. &  
 NE. China, Korea, Ussuri, Amur.

Tribe LEPTURINI ハナカミキリ族

Genus *Pyrrhona* BATES, 1884 ヘリウスハナカミキリ属

- Pyrrhona* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 224 (type-species: *P. laeticolor*  
 BATES, 1884).
- P. laeticolor* BATES, 1884 ヘリウスハナカミキリ  
 BATES, 1884, *ibid.*, p. 224, pl. 1, fig. 8 (Yuyama, Kumamoto, Kyushu).

- forma *takakuwai* KUSAMA, 1971 (黒化型)  
 KUSAMA, 1971, Ent. Rev. Japan, **23**, p. 24 (*P. laeticolor* subsp.) (Kuromidake, Yakushima).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Koshikijima, Yakushima).  
 Genus *Ohbayashia* HAYASHI, 1958 ヘリグロホソハナカミキリ属
- Ohbayashia* HAYASHI, 1958, Ent. Rev. Japan, **9**, p. 4 (type-species: *Strangalomorpha nigromarginata* HAYASHI, 1953).
- O. nigromarginata** (HAYASHI, 1953) ヘリグロホソハナカミキリ  
 HAYASHI, 1953, Ent. Rev. Japan, **6**, p. 40, pl. 8, fig. 3 (*Strangalomorpha*) (Ohsugidani, Mie, Honshu; Mt. Asahimaru, Tokushima, Shikoku).  
 Distr.: Japan.  
 subsp. *nigromarginata* (HAYASHI, 1953), s. str.  
 HAYASHI, 1953, *ibid.*, p. 40, pl. 8, fig. 3 (*Strangalomorpha*) (Ohsugidani, Mie, Honshu, Mt. Asahimaru, Tokushima, Shikoku).  
 Distr.: Japan (Honshu, Shikoku).  
 subsp. *rufoflava* HAYASHI, 1968  
 HAYASHI, 1968, Ent. Rev. Japan, **21**, p. 12 (Mt. Sobo, Ooita, Kyushu).  
 Distr.: Japan (Kyushu).  
 Genus *Alosterna* MULSANT, 1863 ホクチチビハナカミキリ属
- Alosterna* MULSANT, 1863, Col. France Longic., ed. 2, p. 576 (type-species: *Leptura tabacicolor* DEGEER, 1775).
- A. tabacicolor** (DEGEER, 1775) ホクチチビハナカミキリ  
 DEGEER, 1775, Mem. Ins., **5**, p. 139 (*Leptura*) (Europe).  
 Distr.: Japan, Kurile Is., Sakhalin, Korea, NE. China, Siberia, Europe.  
 subsp. *bivittis* (MOTSCHULSKY, 1860)  
 MOTSCHULSKY, 1860, in SCHRENCK's Reisen im Amurlande, **2**, p. 146, pl. 9, fig. 22 (*Grammoptera bivittis*) (Amur).  
*fusca* MATSUSHITA, 1930, Zool. Mag., Tokyo, **42** (495), p. 24 (*tabacicolor* var.) (Kurodake, Hokkaido); PODANY, 1963, Ent. Ges. Basel, **3**, p. 49 (*A. plavilstshikovi* PODANY ab.).  
*nigra* MATSUSHITA et TAMANUKI, 1940, Ins. Matsum., **15**, p. 6 (*tabacicolor* ab.) (Etorofu, Sikotan, Kurile Is., Maruyama, Hokkaido); PODANY, 1963, Ent. Ges. Basel, **3**, p. 49 (*A. plavilstshikovi* PODANY ab.).  
 Distr.: Japan (Hokkaido, Honshu, Rishirito, Sado), Kurile Is., Sakhalin, Korea, NE. China, E. Siberia.  
 Genus *Grammoptera* SERVILE, 1835 チビハナカミキリ属
- Grammoptera* SERVILE, 1835, Ann. Soc. ent. Fr., **4**, p. 215 (type-species: *Leptura praeusta* FABRICIUS, 1787).
- G. chalybeella** BATES, 1884 チビハナカミキリ  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 216 (Nikko, Tochigi, Honshu).  
*debilis* TAMANUKI, 1933, Ins. Matsum., **8**, p. 73 (*Allosterna elegantula* var.) (Konuma, Sakhalin) (クロヒメハナカミキリ).  
*japonica* PIC, 1935, Echange, **41**, p. 11 (Japan) (キアシクロヒメハナカミキリ).  
 Distr.: Japan (Hokkaido to Kyushu, Sado, Tsushima), Kurile Is., Sakhalin, E. Siberia.

Genus *Pachypidonia* GRESSITT, 1935 ヒゲブトハナカミキリ属

*Pachypidonia* GRESSITT, 1935, Kontyû, Tokyo, 9, p. 166 (type-species: *P. crassicornis* GRESSITT, 1935=*Leptura bodemeyeri* PIC, 1934).

*P. bodemeyeri* (PIC, 1934) ヒゲブトハナカミキリ

PIC, 1934, Échange, 50, p. 17 (*Leptura*) (Kobe ?, Honshu).

*crassicornis* GRESSITT, 1935, Kontyû, Tokyo, p. 168 (Kamikochi, Nagano, Honshu).

*kochiana* MATSUSHITA, 1935, Trans. nat. Hist. Soc. Formosa, 25, p. 308 (*Leptura* (*Vadonia*)) (Kochiken, Shikoku).

Distr.: Japan (Hokkaido to Kyushu).

Genus *Nivellia* MULSANT, 1863 クビボソハナカミキリ属

*Nivellia* MULSANT, Col. France Longic., ed. 2, p. 564 (type-species: *Leptura sanguinosa* GYLLENHAL, 1827).

*N. sanguinosa* (GYLLENHAL, 1827) クビボソハナカミキリ

GYLLENHAL, 1827, Ins. Suec., 1, p. 21 (*Leptura*) (Sweden).

Distr.: Japan (Hokkaido), Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.

*N. extensa* GEBLER, 1841 スミイロハナカミキリ

GEBLER, 1841, Bull. Soc. nat. Mosc., 14, p. 613 (Nerchinsk, E. Siberia) (スミイロクビボソハナカミキリ)

Distr.: Japan (Hokkaido, Montane Honshu), Sakhalin, Siberia, N. Europe.

Genus *Munamizoa* MATSUSHITA et TAMANUKI, 1940 ムナミゾハナカミキリ属

*Munamizoa* MATSUSHITA et TAMANUKI, 1940, Ins. Matsum., 15, p. 4, fig. 2 b (type-species: *Nivellia maculata* MATSUSHITA et TAMANUKI, 1935) (*Nivellia* subg.).

*M. maculata* (MATSUSHITA et TAMANUKI, 1935) ムナミゾハナカミキリ

MATSUSHITA & TAMANUKI, 1935, Ins. Matsum., 10, p. 1, fig. 1 (*Nivellia* (*Munamizoa*)) (Mt. Ohdaigahara, Nara, Honshu).

Distr.: Japan (Honshu, Shikoku).

Genus *Pseudalosterna* PLAVILSTSHIKOV, 1934 チャボハナカミキリ属

*Pseudalosterna* PLAVILSTSHIKOV, 1934, Ent. Blätt., 30, p. 132 (type-species; *P. orientalis* PLAVILSTSHIKOV, 1934=*Leptura* (*Leptura*) *misella* BATES, 1884).

*P. misella* (BATES, 1884) チャボハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 216 (*Leptura* (s. str.)) (Kashiwagi, Nara, Wadatoge, Nagano, Honshu).

*orientalis* PLAVILSTSHIKOV, 1934, Ent. Blätt., p. 132 (Raddevka, E. Siberia).

Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima, Koshikijima), Korea, Amur, Ussuri.

*P. takagii* HAYASHI, 1961 クロチャボハナカミキリ

HAYASHI, 1961, Ent. Rev. Japan, 13, p. 38, pl. 9, fig. 4 (Amami-Oshima, N. Ryukyu).

Distr.: Ryukyu Is. (Amami-Oshima).

*P. aritai* OHBAYASHI et OHBAYASHI, 1965 オキナワチャボハナカミキリ

OHBAYASHI & OHBAYASHI, 1965, Bull. Japan ent. Acad., 2, p. 1 (*binotata* GRESSITT subsp.) (Okinawa, C. Ryukyu).

Distr.: Ryukyu Is. (Okinawa).

Genus *Anoploderomorpha* PIC, 1901 ルリハナカミキリ属

*Anoploderomorpha* PIC, 1901, Bull. Mus. Hist. nat. Paris, 7, p. 59 (type-species: *Leptura excavata* BATES, 1884) (*Leptura* subg.).

*A. cyanea* (GEBLER, 1832) ルリハナカミキリ

GEBLER, 1832, Nouv. Mem. Soc. nat. Moscou, 2, p. 70 (*Leptura* (s. str.)) (Siberia).

Distr.: Japan (Hokkaido to Kyushu, Sado), S. Kurile Is., Sakhalin, Korea, NE, China, Mongolia, Siberia, Transbaikal.

*A. excavata* (BATES, 1884) ミヤマクロハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 217 (*Leptura* (s. str.)) (Wadatoge, Nagano, Nyohozan, Tochigi, Honshu).

Distr.: Japan (Honshu, Shikoku, Kyushu, Oki, Yakushima, Koshikijima), E. China.

*A. monticola* NAKANE, 1955 クロルリハナカミキリ

NAKANE, 1955, Akitu, Kyoto, 4, p. 11 (Honshu, Shikoku, Sado).

Distr.: Japan (Honshu, Shikoku, Sado).

*A. kishiii* NAKANE, 1955 エゾクロルリハナカミキリ

NAKANE, 1955, Akitu, Kyoto, 4, p. 11 (Sounkyo, Hokkaido).

Distr.: Japan (Hokkaido).

Genus *Kanekoa* MATSUSHITA et TAMANUKI, 1942 ミヤママルリハナカミキリ属

*Kanekoa* MATSUSHITA et TAMANUKI, 1942, Zool. Mag., Tokyo, 54, p. 79 (type-species: *Leptura (Kanekoa) azumensis* MATSUSHITA et TAMANUKI, 1942) (*Leptura* subg.).

*K. azumensis* (MATSUSHITA et TAMANUKI, 1942) ミヤママルリハナカミキリ

MATSUSHITA & TAMANUKI, 1942, ibid., p. 79, fig. 1 (*Leptura (Kanekoa)*) (Azumasan, Fukushima, Honshu).

Distr.: Japan (Hokkaido to Kyushu, Sado).

Genus *Brachyleptura* CASEY, 1913 ヒメアカハナカミキリ属

*Brachyleptura* CASEY, Mem. Col., 4, p. 247 (type-species: *Leptura vagans* OLIVIER, 1795).

*B. pyrha* (BATES, 1884) ヒメアカハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 216 (*Leptura*) (Wadatoge, Nagano, Nikko, Tochigi, Honshu).

*bouvieri* PIC, 1901, Bull. Mus. Hist. nat. Paris, 7, p. 59 (*Leptura*) (C. Honshu).

Distr.: Japan (Honshu, Shikoku).

Genus *Anastrangalia* CASEY, 1924 ツヤケンハナカミキリ属

*Anastrangalia* CASEY, 1924, Mem. Col., 11, p. 279 (type-species: *Leptura sanguinea* LECONTE, 1859) (*Strangalia* subg.).

*Marthaleptura* OHBAYASHI, 1963, Frag. Col., (2), p. 9 (type-species: *Leptura scotodes* BATES, 1873).

*A. scotodes* (BATES, 1873) ツヤケンハナカミキリ

BATES, 1873, Ann. Mag. nat. Hist., (4), 12, p. 194 (*Leptura*) (Nagasaki, Kyushu).

*niponensis* PIC, 1901, Échange, 17, p. 12 (*Leptura*) (Japan).

*kongoensis* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34, p. 201 (*Leptura*) (Mt. Kongo, Osaka, Honshu).

Distr.: Japan (Hokkaido to Kyushu, Rishirito, Sado, Oki, Tsushima, Yakushima, Kamikoshikijima), Sakhalin, Korea, NE. China, E. Siberia.



- A. sequensi* (REITTER, 1898) ヨコグロハナカミキリ  
 REITTER, 1898, Wien. ent. Z., **17**, p. 194 (*Leptura*) (E. Siberia).  
*sachalinensis* MATSUSHITA, 1933, Ins. Matsum., **8**, p. 104, figs. 2, a, b (*Leptura*) (Sakhalin) (キバネハナカミキリ).  
 Distr.: Japan (Hokkaido), Kurile Is., Sakhalin, Korea, NE. China, Mongolia, E. Siberia.
- A. dissimilis* (FAIRMAIRE, 1900) ニイタカハナカミキリ  
 FAIRMAIRE, 1900, Ann. Soc. ent. France, **68**, p. 639 (*Leptura*) (Foochoo, Fukien, E. China).  
*niitakana* KANO, 1933, Kontyû, Tokyo, **6**, p. 264 (*Leptura*) (Tataka-ambu, Mt. Niitaka, Taiwan) (ニイタカハナカミキリ).  
 Distr.: Ryukyu Is. (Amami-Oshima, Tokunoshima, Ishigakijima), Taiwan, E. China.
- A. hirayamai* (MATSUSHITA et TAMANUKI, 1942) タテスジハナカミキリ  
 MATSUSHITA & TAMANUKI, 1942, Zool. Mag. Tokyo, **54**, p. 639, fig. 2 (*Leptura*) (Japan, questionable locality) (correctly Taiwan) ♀.  
*atofusca* HAYASHI, 1961, Ent. Rev. Japan, **13**, p. 39 (*Aredolpona hirayamai* f.) (Amami-Oshima).  
 Distr.: Ryukyu Is. (Amami-Oshima), Taiwan.

Genus *Corymbia* DES GOZIS, 1886 アカハナカミキリ属

- Corymbia* DES GOZIS, 1886, L'Espèce Typique, p. 33 (type-species: *Leptura rubra* LINNÉ, 1758).  
*Aredolpona* NAKANE et OHBAYASHI, 1957, Sci. Rept. Saikyo Univ., **2** (4), p. 50 (type-species: *Leptura rubra* LINNÉ, 1758).
- C. succedanea* (LEWIS, 1879) アカハナカミキリ  
 LEWIS, 1879, Ann. Mag. nat. Hist., (5), **4**, p. 464 (*Leptura*) (North Japan, apparently Hokkaido).  
 Distr.: Japan (Hokkaido to Kyushu, Okushirito, Sado), S. Kurile Is., Sakhalin, Korea, NE. China, E. Siberia.
- C. variicornis* (DALMAN, 1817) ブチヒゲハナカミキリ  
 DALMAN, 1817, in SCHÖNHERR, Synon. Ins., **1** (3), p. 482, note 1 (*Leptura*) (N. Europe).  
 Distr.: Japan (Hokkaido, Honshu), S. Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.
- C. igai* (TAMANUKI, 1942) イガブチヒゲハナカミキリ  
 TAMANUKI, 1942, Fauna Nipponica, Ceramb., Lept., **2**, p. 93, f. 119 (*Leptura variicornis* subsp.) (Mt. Koya, Wakayama, Honshu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Konoa* MATSUSHITA, 1933 オオハナカミキリ属

- Konoa* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 196 (type-species: *Nivellia (Konoa) valida* MATSUSHITA, 1933 = *Leptura granulata* BATES, 1884) (*Nivellia* subg.).
- K. granulata* (BATES, 1884) オオハナカミキリ  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 217 (*Leptura* (s. str.)) (Sapporo, Yani, Hokkaido).  
*valida* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 196 (*Nivellia (Konoa)*)

(Okuteine, Otaru, Hokkaido).  
Distr.: Japan (Hokkaido, Honshu).

Genus *Lepturobosca* REITTER, 1912 ケブカハナカミキリ属(アオハナカミキリ属)

*Lepturobosca* REITTER, 1912, Fauna Germanica, 4, p. 17 (type-species: *Leptura virens* LINNÉ, 1758) (*Leptura* subg.).

*L. virens* (LINNÉ, 1758) アオハナカミキリ

LINNÉ, 1758, Syst. Nat., ed 10, p. 397 (*Leptura*) (Europe).

Distr.: Japan (Hokkaido), Kurile Is., Sakhalin, NE. China, Mongolia, Siberia, Europe.

Genus *Judolia* MULSANT, 1863 ヤマトキモンハナカミキリ属

*Judolia* MULSANT, 1863, Col. France Longic., ed. 2, p. 496 (type-species: *Leptura sexmaculata* LINNÉ, 1758).

*J. sexmaculata* (LINNÉ, 1758) シララカハナカミキリ

LINNÉ, 1758, Syst. Nat., ed. 10 p. 398 (*Leptura*) (Europe).

*parallelopipeda* MOTSCHULSKY, 1860, in SCHRENCK'S Reisen im Amurlande, 2, p. 146, pl. 9. fig. 23 (Amur).

*shirarakensis* MATSUMURA, 1911, J. Coll. Agr. Tôhoku Imp. Univ., 4(1), p. 137 (*Leptura* (*Pidonia*)) (Shiraraka, Chijisani, Kushunnai, Sakhalin).

Distr.: Japan (Hokkaido, Honshu, Rishirito), Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe, N. America.

*J. japonica* (TAMANUKI, 1942) ヤマトキモンハナカミキリ

TAMANUKI, 1942, Fauna Nipponica, Ceramb., 2, Lept., p. 164, figs. 179, 180 (*Strangalia* (s. str.)) (Mt. Ohdaigahara, Nara, Honshu; Mt. Kirishima, Miyazaki, Kyushu).

Distr.: Japan (Hokkaido to Kyushu).

Genus *Pachytodes* PIC, 1891 マルガタハナカミキリ属

*Pachytodes* PIC, 1891, Mat. Longic., 1, p. 65 (type-species: *Leptura cerambyciformis* SCHRANK, 1781) (*Judolia* subg.).

*P. cometes* (BATES, 1884) マルガタハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., 18, p. 218 (*Leptura* (*Judolia*)) (Chuzenji, Nyohozan, Tochigi, Honshu; Sapporo, Hokkaido).

Distr.: Japan (Hokkaido to Kyushu), S. Kurile Is., Sakhalin.

Genus *Judolidia* PLAVILSTSHIKOV, 1936 スバタマハナカミキリ属

*Judolidia* PLAVILSTSHIKOV, 1936, Faune de l'URSS, 21 (1), pp. 399, 400 (type-species: *J. znojkoii* PLAVILSTSHIKOV, 1936).

*J. bangi* (PIC, 1901) スバタマハナカミキリ

PIC, 1901, Bull. Mus. Hist. nat. Paris, 7, p. 340 (*Leptura*) (Central Japan).

*akitensis* MATSUSHITA, 1931, Trans. Sapporo nat. Hist. Soc. 12, p. 42 (*Leptura* (*Vadonia*)) (Akita, Honshu) (ヤマトハナカミキリ).

*stygica* GRESSITT, 1935, Kontyû, Tokyo, 9, p. 168 (*Leptura*). (Chuzenji, Tochigi, Honshu) (チュウゼンジハナカミキリ).

*znojkoii* PLAVILSTSHIKOV, 1936, Faune de l'URSS, 21 (1), pp. 400, 573, fig. 202 (Os-

sinovka, Ozernaja, Spasskoje, Tigrobaja, Romanovka, Ussuri).  
Distr.: Japan (Hokkaido, Honshu, Kyushu), Korea, Ussuri, N. China.

Genus *Oedecnema* THOMSON, 1857 モモブトハナカミキリ属

*Oedecnema* THOMSON, 1857, Arch. ent., **1**, p. 319 (type-species: *Leptura dubia* FABRICIUS, 1781).

*O. dubia* (FABRICIUS, 1781) モモブトハナカミキリ

FABRICIUS, 1781, Spec. Ins., **1**, p. 249 (*Leptura*) (Siberia).

*10-maculata* MATSUMURA, 1911, J. Coll. Agr. Tôhoku Imp. Univ., **4** (1), p. 136 (*Leptura* (*Strangalia*)) (Kiminai, Shiraraka, Sakhalin; Sapporo, Hokkaido).

Distr.: Japan (Hokkaido, Honshu), S. Kurile Is., Sakhalin, Korea, NE, China, Mongolia, Siberia, Europe.

Genus *Leptura* LINNÉ, 1758 ヨツスジハナカミキリ属

*Leptura* LINNÉ, 1758, Syst. Nat., ed. 10, p. 397 (type-species: *L. quadrifasciata* LINNÉ, 1758).

*L. arcuata* PANZER, 1793 ヤツボンハナカミキリ

PANZER, 1793, Fauna Ins. Germ., **8**, p. 524, fig. 12 (Germany).

Distr.: Japan, Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.

subsp. *mimica* BATES, 1884 (ヤツボンハナカミキリ)

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 219 (*Leptura* (*Stenura*)) (Sapporo, Junsai, Hokkaido; Nikko, Honshu).

*harmandi* PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 61 (*Leptura* (*Strangalia*)) (Central Honshu).

*kinhokuana* MATSUSHITA, 1932, Zool. Mag., Tokyo, **44**, (*Strangalia* (s. str.)) (Sado) (キンホクハナカミキリ).

Distr.: Japan (Hokkaido, Honshu, Kyushu, Rishirito, Sado), Kurile Is.

subsp. *tsumagurohana* OHBAYASHI, 1957 (ツマグロハナカミキリ)

OHBAYASHI, 1957, Ent. Rev. Japan, **6**, p. 61.

*tenicornis* BATES, 1873 (nec MOTSCHULSKY), Ann. Mag. nat. Hist., (4), **12**, p. 11 (*Leptura*) (Nagasaki, Kyushu).

Distr.: Japan (Honshu, Shikoku, Kyushu, Yakushima).

*L. aethiops* PODA, 1761 クロハナカミキリ (コクロハナカミキリ)

PODA, 1761, Ins. Mus. Graecensis, p. 38 (Greece).

*aterrima* MOTSCHULSKY, 1860, in SCHRENCK'S Reisen im Amurlande, **2**, p. 147, pl. 9, fig. 24 (*Stenura*) (Amur).

*adustipennis* SOLSKY, 1870, Horae Soc. ent. Ross. **7**, p. 404 (*Strangalia*) (Siberia) (カネガタホソハナカミキリ).

*coreana* MATSUSHITA, 1933, Ins. Matsum. **7**, p. 104, fig. 3 (*Strangalia*) (Korea) (チャバネハナカミキリ).

*doii* MATSUSHITA, 1933, ibid., p. 105, fig. 4 (*Strangalia*) (Etorofu, S. Kurile Is.) (チンマハナカミキリ, ドイハナカミキリ).

Distr.: Japan, Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.

subsp. *aethiops* PODA, 1761, s. str.

PODA, 1761, Ins. Mus. Graecensis, p. 38 (Greece).

Distr.: Japan (Hokkaido, Montane Honshu), Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.

- subsp. *dimorpha* BATES, 1873 (ムネアカクロハナカミキリ)  
 BATES, 1873, Ann. Mag. nat. Hist., (4), **12**, p. 195 (*L. dimorpha*) (Japan).  
 Distr.: Japan (Hokkaido to Kyushu, Sado), Kurile Is., Sakhalin, Korea, E. Siberia.
- L. latipennis* (MATSUSHITA, 1933) ハネビロハナカミキリ  
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 214, pl. 1, fig. 10 (*Strangalia* (s. str.)) (Kamiotoineppu, Sapporo, Tomakomai, Hidaka, Hokkaido; Chichibu, Tokyo, Honshu).  
 Distr.: Japan (Hokkaido to Kyushu), Kurile Is.
- L. duodecimguttata* FABRICIUS, 1801 キモンハナカミキリ  
 FABRICIUS, 1801, Syst. Eleuth., **2**, p. 363 (Siberia).  
 Distr.: Japan (Hokkaido, Honshu, Tsushima), Sakhalin, Korea, NE. China, Mongolia, Siberia.
- L. quadrifasciata* LINNÉ, 1758 カラフトヨツスジハナカミキリ  
 LINNÉ, 1758, Syst. Nat., ed. 10, p. 398 (Europe).  
 Distr.: Japan (Hokkaido), Kurile Is., Sakhalin, Korea, NE. China, Siberia, Caucasia, Europe.
- L. ochraceofasciata* MOTSCHULSKY, 1861 ヨツスジハナカミキリ  
 MOTSCHULSKY, 1861, Étud. ent., **10**, p. 21 (*Stenura*) (Japan).  
 Distr.: Japan, Kurile Is., Sakhalin, Korea, NE. China.
- subsp. *ochraceofasciata* (MOTSCHULSKY, 1861), s. str.  
 MOTSCHULSKY, 1861, *ibid.*, p. 21 (*Stenura*) (Japan).  
 Distr.: Japan (Hokkaido, Honshu, Okushirito, Rishirito, Izu-Oshima, Nijijima, Kozujima), Kurile Is., Sakhalin, Korea, NE. China.
- subsp. *ochrotela* BATES, 1873  
 BATES, 1873, Ann. Mag. nat. Hist., (4), **12**, p. 196 (var.) (Nagasaki, Kyushu).  
 Distr.: Japan (Shikoku, Kyushu).
- subsp. *chujoi* HAYASHI, 1961  
 HAYASHI, 1961, Ent. Rev. Japan, **13**, p. 21, figs. 4, 6, 10 (Tsushima).  
 Distr.: Japan (Tsushima).
- subsp. *yokoyamai* HAYASHI, 1961  
 HAYASHI, 1961, *ibid.*, p. 21, figs. 3, 5, 9 (Yakushima).  
 Distr.: Japan (Tanegashima, Yakushima).
- subsp. *amamiana* HAYASHI, 1960 (アマミヨツスジハナカミキリ)  
 HAYASHI, 1960, Ent. Rev. Japan, **11**, p. 22, text fig. 2 (*L. auratopilosa* MATSUSHITA subsp.) (Amami-Oshima).  
 Distr.: Ryukyu Is. (Amami-Oshima, Tokunoshima).
- subsp. *watanabei* HAYASHI, 1962  
 HAYASHI, 1962, Ent. Rev. Japan, **15**, p. 2, pl. 1, fig. 2 (Okinawa).  
 Distr.: Ryukyu Is. (Okinawa).
- L. kusamai* OHBAYASHI et NAKANE, 1955 ヒメヨツスジハナカミキリ  
 OHBAYASHI & NAKANE, 1955, Akitu, Kyoto, **4**, p. 17, figs. 1-3 (Ooi-Sawarajima, Shizuoka, Honshu).  
 Distr.: Japan (Honshu, Kyushu).
- L. subtilis* BATES, 1884. コヨツスジハナカミキリ (ヤマトヨツスジハナカミキリ)  
 BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 219 (*L. (Stenura)*) (Chuzenji, Tochigi, Honshu) ♂.

*oishii* MATSUSHITA et TAMANUKI, 1935, Ins. Matsum., **10**, p. 147 (*Strangalia* (s. str.) *subtilis* BATES ab.) (Mt. Sobo, Ooita, Kyushu) ♀ (オオイシヨツスジハナカミキリ).

Distr.: Japan, Kurile Is.

subsp. *subtilis* BATES, 1884, s. str.

BATES 1884., *ibid.*, p. 219 (L. (*Stenura*)) (Chuzenji).

Distr.: Japan (Honshu, Shikoku, Kyushu), Kurile Is.

subsp. *aureopubescens* HAYASHI, 1974

HAYASHI, 1974, Ent. Rev. Japan, **26**, p. 12 (Yakushima).

Distr.: Japan (Yakushima).

*L. yakushimana* (TAMANUKI, 1942) ヤクシマヨツスジハナカミキリ

TAMANUKI, 1942, Fauna Nipponica Ceramb., **2**, Lept., p. 153, fig. 168 (*Strangalia* (s. str.)) (Kosugidani, Yakushima) ♂; N. OHBAYASHI, 1970, Bull. Japan ent. Acad., **5**, p. 1 figs. 1, 2, 7, 8; ♂, ♀.

Distr.: Japan (Yakushima).

Genus *Megaleptura* CASEY, 1913 オオヨツスジハナカミキリ属

*Megaleptura* CASEY, 1913, Mem. Col., **4**, p. 267 (type-species: *Leptura emarginata* FABRICIUS, 1787) (*Leptura* subg.).

*Stenura* DEJEAN, 1837, Cat. Col., p. 381 (nec CUVIER, 1829 for Aves).

*Macroleptura* NAKANE et OHBAYASHI, 1957, Sci. Rept. Saikyo Univ., **2** (4), p. 47 (type-species: *Leptura thoracica* CREUTZER, 1799).

*M. thoracica* (CREUTZER, 1799) クロオオハナカミキリ (セアカハナカミキリ)

CREUTZER, 1799, Ent. Versuche, p. 125, pl. 3, fig. 28 (*Leptura*) (Karlsbad, Czechoslovakia).

Distr.: Japan (Hokkaido, Honshu, Shikoku), S. Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Europe.

*M. regalis* (BATES, 1884) オオヨツスジハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 223 (*Strangalia*) (Sapporo, Hokkaido; Iga, Mie, Honshu).

*maindroni* PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 61 (*Leptura* (*Strangalia*)) (Central Honshu).

Distr.: Japan (Hokkaido to Kyushu, Yakushima), S. Kurile Is., Sakhalin, Korea.

Genus *Nakanea* OHBAYASHI, 1963 フタスジハナカミキリ属

*Nakanea* OHBAYASHI, 1963, Fragm. Col., (2) p. 9 (type-species: *Leptura* (*Stenura*) *vicaria* BATES, 1884).

*N. vicaria* (BATES, 1884) フタスジハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 218 (*Leptura* (*Stenura*)) (Nyohozan, Tochigi, Honshu; Sapporo, Hokkaido).

Distr.: Japan, S. Kurile Is., Sakhalin.

subsp. *vicaria* (BATES, 1884), s. str.

BATES, 1884, *ibid.*, p. 218 (*Leptura* (*Stenura*)) (Nyohozan, Tochigi, Honshu; Sapporo, Hokkaido).

Distr.: Japan (Hokkaido to Kyushu, Yakushima), S. Kurile Is., Sakhalin.

subsp. *adumbrata* (BATES, 1884) (ハススジハナカミキリ)

BATES, 1884, *ibid.*, p. 220 (*Leptura* (*Stenura*)) (Tokyo ?, Honshu).

*lajoyei* PIC, 1943, Opusc. Mart., **10**, p. 1 (*Strangalia*) (Japan).  
*biluteomaculata* PIC, 1943, *ibid.*, p. 1 (*Strangalia*) (Japan).  
 Distr.: Japan (S. Central Honshu).

Genus *Pedostrangalia* SOKOLOW, 1896 カタキハナカミキリ属

*Pedostrangalia* SOKOLOW, 1896, Horae ent. Ross., **30**, p. 461 (type-species: *Pedostrangalia kassajanowi* SOKOLOW, 1896) (*Strangalia* subg.).

*Sphenalia* DANIEL, 1904, München. Kol. Z., **2**, p. 355 (type-species: *Sphenalia ariadne* DANIEL, 1904).

*Etorofus* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, pp. 183, 204 (type-species: *E. variicornis* MATSUSHITA, 1933).

*P. femoralis* (MOTSCHULSKY, 1860) カタキハナカミキリ (チョウセンクロハナカミキリ)  
 MOTSCHULSKY, 1860, Étud. ent., **9**, p. 40 (*Stenura*) (Amur).

*xanthoma* BATES, 1873, Ann. Mag. nat. Hist., (4), **12**, p. 195 (*Leptura*) (カタキハナカミキリ).

Distr.: Japan (Hokkaido, Honshu, Sado), Kurile Is., Sakhalin, Korea, NE. China, E. Siberia.

*P. variicornis* (MATSUSHITA, 1933) エトロフハナカミキリ

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 205 (*Etorofus*) (Etorofu Is., Kurile Is.).

*circaocularis* PIC, 1934, Échange, **50**, p. 18 (*Strangalia*) (Sakhalin).

Distr.: Japan (Hokkaido), S. Kurile Is., Sakhalin.

Genus *Eustrangalis* BATES, 1884 クロスジハナカミキリ属

*Eustrangalis* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 222 (type-species: *E. distenoides* BATES, 1884).

*E. distenoides* BATES, 1884 クロスジハナカミキリ (カエデノヘリグロハナカミキリ)

BATES, 1884, *ibid.*, p. 222, pl. 1, fig. 4 (Nikko, Tochigi, Honshu; Sapporo, Hokkaido).  
 Distr.: Japan (Hokkaido to Kyushu), S. Kurile Is.

*E. anticereductus* (HAYASHI, 1955) クロサワヘリグロハナカミキリ

HAYASHI, 1955, Col. Illust. Ins. Japan, ed. 1, **1**, p. 31 (*E. distenoides* BATES ab.) (Jozankei, Hokkaido).

Distr.: Japan (Hokkaido to Kyushu).

Genus *Japanostrangalia* NAKANE et OHBAYASHI, 1957 ヒゲジロハナカミキリ属

*Japanostrangalia* NAKANE et OHBAYASHI, 1957, Sci. Rept. Saikyo Univ., **2** (4), p. 50 (type-species: *Leptura dentatipennis* PIC, 1901).

*J. dentatipennis* (PIC, 1901) ヒゲジロハナカミキリ

PIC, 1901, Bull. Mus. Hist. nat. Paris, **6**, p. 60 (*Leptura*) (Central Honshu).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Strangalomorpha* SOLSKY, 1873 アオバホソハナカミキリ属

*Strangalomorpha* SOLSKY, 1873, Horae Soc. ent. Ross., **9**, p. 253 (type-species: *S. tenuis* SOLSKY, 1873).

*S. tenuis* SOLSKY, 1873 アオバホソハナカミキリ

SOLSKY, 1873, *ibid.*, p. 253 (Suyfun, Amur).

*aenescens* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 221 (Chuzenji, Nyohozan, Tochigi; Wada-toge, Nagano, Honshu).

*aenescens atricolor* KANO, 1933, Kontyû, Tokyo, **7**, p. 130 (Kamikochi, Nagano, Honshu) (クロホソハナカミキリ).

Distr.: Japan (Honshu, Shikoku, Kyushu, Sado, Oki, Tsushima), Sakhalin, Korea, NE. China, Mongolia, Siberia.

Genus *Parastrangalis* GANGLBAUER, 1889 ニンフハナカミキリ属

*Parastrangalis* GANGLBAUER, 1889, Horae Soc. ent. Ross., **24**, p. 57 (type-species: *Leptura (Parastrangalis) potanini* GANGLBAUER, 1889) (*Leptura* subg.).

*P. nymphula* (BATES, 1884) ニンフハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 220 (*Leptura (Stenura)*) (Chuzenji, Nyohozan, Tochigi; Wada-toge, Nagano, Honshu).

Distr.: Japan (Hokkaido to Kyushu, Sado, Oki).

*P. lesnei* (PIC, 1901) ニヨウホウハナカミキリ

PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 60 (*Leptura nymphula* BATES ? var.) (Central Honshu).

*betesi* TAMANUKI, 1942, Fauna Nipponica, Ceramb., **2**, Lept., p. 134, fig. 153 (*Strangalomorpha*) (Honshu, Shikoku, Kyushu).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Strangaliella* HAYASHI, 1976 タテジマハナカミキリ属

*Strangaliella* HAYASHI, 1976, Bull. Osaka Jonan Women's Jr. Coll., **11**, p. 3 (type-species: *Strangalia shikokensis* MATSUSHITA, 1935).

*S. shikokensis* (MATSUSHITA, 1935) タテジマハナカミキリ

MATSUSHITA, 1935, Trans. nat. Hist. Soc. Formosa, **25**, p. 309 (*Strangalia* (s. str.)) (Kochiken, Shikoku).

Distr.: Japan (Honshu, Shikoku, Kyushu).

*S. ishigakiensis* (HAYASHI, 1971) ヤエヤマクロスジホソハナカミキリ

HAYASHI, 1971, Ent. Rev. Japan, **23**, p. 5 (*Parastrangalis*) (Mt. Omotodake, Ishigaki).

Distr.: Ryukyu Is. (Ishigaki).

Genus *Strangalia* SERVILLE, 1835 ホソハナカミキリ属

*Strangalia* SERVILLE, 1835, Ann. Soc. ent. France, **4**, p. 220 (type-species: *Leptura luteicornis* FABRICIUS, 1775).

*Ophistomis* THOMSON, 1857, Arch. ent., **1**, p. 319; THOMSON, 1860, Essai Classif. Ceramb., p. 154 (type-species: *O. flavocinctus* THOMSON, 1860).

*Strangalina* AURIVILLIUS, 1912, Col. Cat., **39**, p. 245 (new name for *Strangalia*).

*Sulcastostrangalia* OHBAYASHI, 1961, Ent. Rev. Japan, **13**, p. 17 (type-species: *Strangalia gracilis* GRESSITT, 1934) (*Strangalia* subg.).

*S. attenuata* LINNÉ, 1758 ヨツスジホソハナカミキリ

LINNÉ, 1758, Syst. Nat., ed. 10, p. 398 (Europe).

Distr.: Japan (Hokkaido), Kurile Is., Sakhalin, Korea, NE. China, Mongolia, Siberia, Caucasia, Europe.

*S. takeuchii* MATSUSHITA et TAMANUKI, 1935 タケウチホソハナカミキリ

MATSUSHITA & TAMANUKI, 1935, Ins. Matsum., **10**, p. 2, fig. 2 (*S. (Strangalina)*)

(Mt. Odaigahara, Nara, Honshu).

Distr.: Japan (Hokkaido to Kyushu).

**S. gracilis** GRESSITT, 1934 オオシマホソハナカミキリ

GRESSITT, 1934, Philip. J. Sci., **55**, p. 381 (*S. (Strangalina)*) (Amami Oshima).

Distr.: Ryukyu Is. (Amami-Oshima).

**S. koyaensis** MATSUSHITA, 1933 コウヤホソハナカミキリ

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 219, pl. 1, fig. 7 (*S. (Strangalina)*) (Mt. Koya, Wakayama, Honshu).

Distr.: Japan.

subsp. *koyaensis* (MATSUSHITA, 1933), s. str.

MATSUSHITA, 1933, *ibid.*, p. 219, pl. 1, fig. 7 (*S. Strangalina*) (Mt. Koya, Wakayama, Honshu).

Distr.: Japan (Honshu, Shikoku, Kyushu).

subsp. *haraguro* OHBAYASHI, 1958

OHBAYASHI, 1958, Ent. Rev. Japan, **9**, p. 18 (Shimashima, Nagano, Honshu).

Distr.: Japan (Central Honshu).

Genus *Mimostrangalia* NAKANE et OHBAYASHI, 1957 クロソソホソハナカミキリ属

*Mimostrangalia* NAKANE et OHBAYASHI, 1957, Sci. Rept. Saikyo Univ., **2** (4), p. 49 (type-species: *Strangalia kurosonensis* OHBAYASHI, 1936).

*Insularestrangalia* HAYASHI, 1961, Ent. Rev. Japan, **13**, p. 40 (type-species: *Strangalia longicornis* GRESSITT, 1934) (*Strangalia* subg.).

**M. kurosonensis** (OHBAYASHI, 1936) クロソソホソハナカミキリ

OHBAYASHI, 1936, Trans. Kansai ent. Soc., (7), p. 11, pl. 2, fig. 1 (*Strangalia*) (Kuroson, Kochi, Shikoku).

Distr.: Japan (W. Honshu, Shikoku, Kyushu, Tsushima, Yakushima, Izu-Mikurajima), Ryukyu Is. (Amami-Oshima).

**M. dulcis** (BATES, 1884) ジャコウホソハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 222 (*Strangalia*) (Wada-toge, Nagano, Honshu; Yuyama, Kumamoto, Kyushu).

*kinoshitai* KANO, 1933, Kontyû, Tokyo, **6**, p. 265 (*Typocerus*) (Aburazaka, Gifu, Honshu) (キノシタハナカミキリ).

Distr.: Japan (Honshu, Shikoku, Kyushu, Yakushima).

**M. longicornis** (GRESSITT, 1934) ヒゲナガホソハナカミキリ

GRESSITT, 1934, Philip. J. Sci., **55**, p. 382 (*Strangalia (Strangalina)*) (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima, Tokunoshima, Okinawa).

Genus *Idiostrangalia* NAKANE et OHBAYASHI, 1957 ミヤマホソハナカミキリ属

*Idiostrangalia* NAKANE et OHBAYASHI, 1957, Sci. Rept. Saikyo Univ., **2** (4), p. 49 (type-species: *Strangalia contracta* BATES, 1884).

*Pygostrangalia* HAYASHI, 1960 (nec PIC, 1957), Niponius, **1** (6), p. 18 (part).

**I. contracta** (BATES, 1884) ミヤマホソハナカミキリ

BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 223 (*Strangalia*) (Kashiwagi, Nara; Nyohozan, Tochigi; Wada-toge, Nagano, Honshu).

*ohbayashii* MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 220, pl. 1, fig. 11



(*Strangalia (Strangalina)*) (Shimashima, Nagano, Honshu).  
*lineatocolle* GRESSITT, 1937, Kontyû, Tokyo, **11**, p. 319, fig. 3 (*Strangalina*) (Mt. Yatsugatake, Nagano, Honshu) (スジクビホソハナカミキリ).  
 Distr.: Japan (Honshu, Shikoku, Kyushu), E. China?.

**I. hakonensis** (MATSUSHITA, 1933) ハコネホソハナカミキリ  
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., **34**, p. 218 (*Strangalia (Strangalina)*) (Hakone, Kanagawa; Shimashima, Nagano, Honshu; Awa, Tokushima, Shikoku).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).

**I. maruokai** (HAYASHI, 1963) マルオカホソハナカミキリ  
 HAYASHI, 1963, Ent. Rev. Japan, **15**, p. 52, pl. 7, fig. 2 (*Pygostrangalia (Idiostrangalia)*) (Hatsuno, Amami-Oshima).  
 Distr.: Ryukyu Is. (Amami-Oshima).

Genus *Leptostrangalia* NAKANE et OHBAYASHI, 1959 ヒメホソハナカミキリ属

*Leptostangalia* NAKANE et OHBAYASHI, 1959, Sci. Rept. Kyoto Pref. Univ., **3** (1), p. 64 (type-species: *Strangalia hosohana* OHBAYASHI, 1952).

**L. hosohana** (OHBAYASHI, 1952) ホソハナカミキリ  
 OHBAYASHI, 1952, Ent. Arb. Mus. Frey, **3**, p. 414 (*Strangalia*) (Nidoage, Gumma; Mt. Koya, Wakayama, Honshu; Mt. Saragamine, Ehime, Shikoku; Mt. Fukuchi, Fukuoka, Kyushu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Paranaspia* MATSUSHITA et TAMANUKI, 1940 ベニバハナカミキリ属

*Paranaspia* MATSUSHITA et TAMANUKI, 1940, Ins. Matsum., **15**, p. 5 (type-species: *Strangalia anaspidooides* BATES, 1873) (*Strangalia* subg.).

**P. anaspidooides** (BATES, 1873) ベニバハナカミキリ  
 BATES, 1873, Ann. Mag. nat. Hist., (4), **12**, p. 196 (*Leptura*) (Japan).  
*mikadoi* PIC, 1906, Mat. Longic., **6** (1), p. 16 (*Leptura*) (Kyoto, Honshu).  
 Distr.: Japan (Honshu, Shikoku, Kyushu, Kammurijima, Izu-Mikurajima, Tsushima, Yakushima).

**P. yayeyamensis** HAYASHI et YOKOYAMA, 1974 ヤエヤマヒオドシハナカミキリ  
 HAYASHI & YOKOYAMA, 1974, Ent. Rev. Japan, **26**, p. 12 (Iriomotejima).  
 Distr.: Ryukyu Is. (Iriomote, Ishigaki).

Genus *Ephies* PASCOE, 1866 モウセンハナカミキリ属

*Ephies* PASCOE, 1866, Proc. Zool. Soc. London, p. 506 (type-species: *E. cruentus* PASCOE, 1866).

**E. japonicus** NAKANE et OHBAYASHI, 1961 モウセンハナカミキリ  
 NAKANE & OHBAYASHI, 1961, Fragm. Col., (1), p. 6 (Miyanoura-genba, Yakushima; Mt. Inunaki, Mt. Fukuchi, Fukuoka, Kyushu).  
 Distr.: Japan (W. Honshu, Shikoku?, Kyushu, Yakushima, Tanegashima).

**E. okinawanus** HAYASHI, 1968 リュウキュウモウセンハナカミキリ  
 HAYASHI, 1968, Ent. Rev. Japan, **20**, p. 21, pl. 1, fig. 2 (Okinawa).  
 Distr.: Ryukyu Is. (Okinawa).

## Tribe EROSCHEMINI キスツヤハナカミキリ族

Genus *Corennys* BATES, 1884 キスツヤハナカミキリ属

*Corennys* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 224 (type-species: *C. sericata* BATES, 1884).

*C. sericata* BATES, 1884 キスツヤハナカミキリ

BATES, 1884, *ibid.*, p. 225, pl. 1, fig. 2 (Nanai, Hokkaido; Chuzenji, Tochigi; Ontake, Nagano, Honshu).

Distr.: Japan (Hokkaido to Kyushu), N. China.

Genus *Formosopyrrhona* HAYASHI, 1957 アカハネハナカミキリ属

*Formosopyrrhona* HAYASHI, 1957, Akitu, Kyoto, **6**, p. 37 (type-species: *Corennys cinabarina*: HAYASHI (nec GRESSITT, 1951, 1957 = *Pyrocalymma hozanensis* MATSUSHITA, 1931); HAYASHI, 1960, Niponius, **1** (6), p. 22; HAYASHI, 1974, Bull. Osaka Jonan Women's Jr. Coll., **9**, p. 14.

*F. satoi* (HAYASHI, 1957) アマミアカハネハナカミキリ

HAYASHI, 1957, Ent. Rev. Japan, **8**, p. 45, fig. 1 (*Pyrocalymma*) (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima, Tokunoshima, Okinawa).

## Tribe NECYDALINI ホソコバネカミキリ族

Genus *Necydalis* LINNÉ, 1758 ホソコバネカミキリ属

*Necydalis* LINNÉ, 1758, Syst. nat., ed. 10, p. 421 (type-species: *N. major* LINNÉ, 1758 ?).

Subgenus *Necydalis* s. str.*N. (N.) gigantea* KANO, 1933 オニホソコバネカミキリ

KANO, 1933, Kontyû, Tokyo, **6**, p. 271 (Shiratori, Gifu, Honshu).

*marginalis* OHBAYASHI, 1948, Ent. Rev. Japan, **1**, p. 14 (Yonezawa, Yamagata, Honshu).

Distr.: Japan, Kurile Is.

subsp. *gigantea* KANO, 1933, s. str.

KANO, 1933, Kontyû, Tokyo, **6**, p. 271 (Shiratori, Gifu, Honshu).

Distr.: Japan (Hokkaido to Kyushu), Kurile Is.

subsp. *akiyamai* HAYASHI, 1978

HAYASHI, 1978, Ent. Rev. Japan, **31**, p. 87 (Shiratani, Yakushima).

Distr.: Japan (Yakushima).

*N. (N.) major* LINNÉ, 1758 チョウセンホソコバネカミキリ

LINNÉ, 1758, Syst. nat. ed. 10, p. 421 (Europe)

Distr.: Japan, Sakhalin, Korea, Siberia, Caucasia, Europe.

subsp. *aino* KUSAMA, 1974 (アイヌホソコバネカミキリ)

KUSAMA, 1974, Rept. Fac. Sci., Shizuoka Univ., **9**, p. 54 (Shumarinai, Sorachi, Hokkaido).

Distr.: Japan (Hokkaido).

*N. (N.) yakushimensis* KUSAMA, 1975 ヤクシマホソコバネカミキリ

KUSAMA, 1975, Elytra, Tokyo, **2** (2), p. 22 (new name for *N. brevicornis* YOKOYAMA, 1971).

*brevicornis* YOKOYAMA, 1971, Ent. Rev. Japan, **23**, p. 93, pl. 6, fig. 1 (Yakushima) (nec LINNÉ, 1758).

Distr.: Japan (Yakushima).

- N. (N.) moriyai* KUSAMA, 1970 アマミホソコバネカミキリ  
KUSAMA, 1970, Ent. Rev. Japan, **22**, p. 73, pl. 7, fig. 1-6 (Amami-Oshima).  
Distr.: Ryukyu Is. (Amami-Oshima).

Subgenus *Necydalisca* PLAVILSTSHIKOV, 1936

- Necydalisca* PLAVILSTSHIKOV, 1936, Faune d'URSS, **21**, p. 464 (type-species: *Necydalis ebenina* BATES, 1884) (*Necydalis* subg.).
- N. (Necydalisca) sachalinensis* MATSUMURA et TAMANUKI, 1927 カラフトホソコバネカミキリ  
MATSUMURA & TAMANUKI, 1927, Ins. Matsum., **1**, p. 176 (Ichinosawa, Furumaki, Higashisakotan, Sakhalin).  
Distr.: Japan (Honshu), Sakhalin, Korea, E. Siberia.
- N. (Necydalisca) pennata* LEWIS, 1879 ホソコバネカミキリ  
LEWIS, 1879, Ann. Mag. nat. Hist., (5), **4**, p. 464 (Japan).  
*ebenina* BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 225 (Junsai, Hokkaido) (ツヤホソコバネカミキリ).  
*eoae* PLAVILSTSHIKOV, 1936, Faune d'URSS, **21**, p. 470, 591 (Vladivostok, Ussuri).  
Distr.: Japan (Hokkaido, Tsushima), S. Kurile Is., Quelpart Is., Sakhalin, E. Siberia.
- N. (Necydalisca) morio* KRAATZ, 1879 ウスリーホソコバネカミキリ  
KRAATZ, 1879, Dtsch. ent. Z., **23**, p. 106 (Ussuri).  
*pacifica* PLAVILSTSHIKOV, 1936, Faune d'URSS, **21**, p. 471 (Vladivostok, Ussuri).  
*ussuriensis* PLAVILSTSHIKOV, 1936, *ibid.*, p. 473 (Vladivostok, Sedanka, Osinovka, Ascord Is., Ussuri).  
*semenovi* PLAVILSTSHIKOV, 1936, *ibid.*, p. 475 (Chabarovsk, E. Siberia; Osinovka, Ussuri).  
Distr.: Japan (Hokkaido), NE. China, E. Siberia.
- N. (Necydalisca) solida* BATES, 1884 オオホソコバネカミキリ.  
BATES, 1884, J. Linn. Soc. London, Zool., **18**, p. 225 (Chuzenji, Tochigi, Honshu).  
*nikkoensis* MATSUMURA et TAMANUKI, 1927, Ins. Matsum., **1**, p. 175, fig. 1 (Chuzenji, Tochigi, Honshu) (ニッコウホソコバネカミキリ).  
*towadensis* MATSUMURA et TAMANUKI, 1927, *ibid.*, p. 175 (Towada, Aomori, Honshu) (トワダホソコバネカミキリ).  
*galloisi* MATSUMURA et TAMANUKI, 1927, *ibid.*, p. 176, fig. 2 (Chuzenji, Tochigi, Honshu) (ガロアホソコバネカミキリ).  
Distr.: Japan (Honshu, Shikoku, Kyushu).
- N. (Necydalisca) harmandi* PIC, 1901 クロホソコバネカミキリ  
PIC, 1901, Bull. Mus. Hist. nat. Paris, **7**, p. 340 (Central Honshu).  
Distr.: Japan (Honshu, Shikoku, Kyushu).
- N. (Necydalisca) odai* HAYASHI, 1951 ヒゲジロホソコバネカミキリ  
HAYASHI, 1951, Ent. Rev. Japan, **5**, p. 76 (Mt. Gomanodan, Wakayama, Honshu; Mt. Takachiho, Miyazaki, Kyushu).  
Distr.: Japan (Hokkaido to Kyushu).

Subgenus *Eonecydalis* OHBAYASHI, 1961

- Eonecydalis* OHBAYASHI, 1961, Ent. Rev. Japan, **13**, p. 17 (type-species: *N. formosana* KANO, 1933) (*Necydalis* subg.).

- N. (Eonecydalis) formosana* KANO, 1933 タイワンホソコバネカミキリ  
KANO, 1933, Kontyû, Tokyo, 6, p. 270 (Kyuhabon, Formosa).  
Distr.: Japan, Taiwan.
- subsp. *niimurai* HAYASHI, 1949 (トガリバホソコバネカミキリ)  
HAYASHI, 1949, Trans. Kansai ent. Soc., 14 (2), p. 14, fig. (Mt. Ashitaka, Shizuoka,  
Honshu).  
Distr.: Japan (Honshu, Shikoku).
- subsp. *matsudai* HAYASHI, 1949  
HAYASHI, 1949, ibid., p. 16 (Mt. Sobo, Ooita, Kyushu).  
Distr.: Japan (Kyushu, Yakushima).

