

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 21

March 30, 1983

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family ENDOMYCHIDAE テントウムシダマシ科

By

Hiroyuki SASAJI

Subfamily MYCHOTHENINAE マルテントウダマシ亜科

Genus *Mychothenus* STROHECKER, 1953 ダエンテントウダマシ属

Mychothenus STROHECKER, 1953, Gen. Ins., **210**, p. 24 (type-species: *Mychophilus minutus* FRIVALDSZKY, 1877).

Mychophilus FRIVALDSZKY, 1877, Termész. Füzetek, **1**, p. 19 (nec HESSE, 1865).

M. asiaticus SASAJI, 1978 ダエンテントウダマシ

SASAJI, 1978, Mem. Fac. Educ. Fukui Univ., Sect. II (Nat. Sci.), (28), p. 2 (Mt. Hikosan, Fukuoka Pref.).

Distr.: Japan (Honshu, Kyushu).

Genus *Dialexia* GORHAM, 1891 ベニモンマルガタテントウダマシ属

Dialexia GORHAM, 1891, Biol. Cent. Amer. Coleopt., **7**, p. 146 (type-species: *Dialexia setulosa* GORHAM, 1891).

D. hisanoi SASAJI, 1978 ベニモンマルガタテントウダマシ

SASAJI, 1978, Mem. Fac. Educ. Fukui Univ., Sect. II (Nat. Sci.), (28), p. 12 (Iwaya, Fukui Pref.).

Distr.: Japan (Honshu).

Genus *Dexialia* SASAJI, 1970 ヒメマルガタテントウダマシ属

Dexialia SASAJI, 1970, Etizenia, Fukui, (43), p. 4 (type-species: *Dexialia ovalia* SASAJI, 1970).

D. minor (CHÛJÔ, 1941) ヒメマルガタテントウダマシ

CHÛJÔ, 1941, Mushi, Fukuoka, **13**, p. 82 (*Symbiotes*) (Noziri, Nagano-ken).
Distr.: Japan (Hokkaido, Honshu, Kyushu).

Genus *Bystodes* STROHECKER, 1953 マルガタテントウダマン属

Bystodes STROHECKER, 1953, Gen. Ins., **210**, p. 19 (type-species: *Bystodes paulus* STROHECKER, 1953).

B. orbicularis (GORHAM, 1887) マルガタテントウダマン

GORHAM, 1887, Proc. zool. Soc. London, **1887**, p. 652 (*Symbiotes*?) (Kashiwagi, Kurigahara).

Distr.: Japan (Honshu).

Genus *Idiophyes* BLACKBURN, 1895 コマルガタテントウダマン属

Idiophyes BLACKBURN, 1895, Trans. roy. Soc. Austral., **19**, p. 234 (type-species: *Idiophyes brevis* BLACKBURN, 1895).

I. niponensis (GORHAM, 1874) コマルガタテントウダマン

GORHAM, 1874, Ent. month. Mag., **10**, p. 225 (*Symbiotes*) (Nagasaki).

Distr.: Japan (Honshu, Shikoku, Kyushu), China?

I. boninensis SASAJI, 1978 オガサワラマルテントウダマン

SASAJI, 1978, Mem. Fac. Educ. Fukui Univ., Sect. II (Nat. Sci.), (28), p. 19 (Chichijima).

Distr.: Bonin Is. (Chichijima).

Genus *Clemmus* HAMPE, 1850 ヒメテントウダマン属

Clemmus HAMPE, 1850, Stett. ent. Ztg., **11**, p. 353 (type-species: *Clemmus troglodytes* HAMPE, 1850).

C.? *abbreviatis* REITTER, 1889 ヒメテントウダマン

REITTER, 1889, Dtsch. ent. Z., **33**, p. 277 (*Clemmus*) (Japan).

Distr.: Japan.

Genus *Asymbius* GORHAM, 1896 セスジテントウダマン属

Asymbius GORHAM, 1896, Ann. Mus. Stor. nat. Genova, **36**, p. 299 (type-species: *Asymbius crinipes* GORHAM, 1896).

A. foveicollis CHÛJÔ et KIUCHI, 1974 トカラセスジテントウダマン

CHÛJÔ & KIUCHI, 1974, Bull. Jap. ent. Acad., **8**, p. 1 (Is. Nakanoshima, Tokara).

Distr.: Ryukyu Is. (Nakanoshima).

Subfamily MYCETAEINAE ホソテントウダマン亜科

Tribe LEIESTINI ホソテントウダマン族

Genus *Leiestes* REDTENBACHER, 1845 イツホソテントウダマン属

Leiestes REDTENBACHER, 1845, Gattung Dtsch. Käfer-Fauna, p. 121 (type-species: *Cryptophagus seminiger* GYLLENHAL, 1808).

L. decoratus (GORHAM, 1887) イツホソテントウダマン

GORHAM, 1887, Proc. zool. Soc. London, **1887**, p. 648 (*Panamomus*) (Oyayama, Kashiwagi).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Panamomus* GORHAM, 1873 ホソテントウダマン属

Panamomus GORHAM, 1873, Ent. month. Mag., 9, p. 207 (type-species: *Panamomus lewisi* GORHAM, 1873).

P. brevicornis GORHAM, 1887 ホソテントウダマン

GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 649 (Miyanoshiba).
Distr.: Japan (Honshu, Shikoku).

P. lewisi GORHAM, 1873 ルイステントウダマン

GORHAM, 1873, Ent. month. Mag., 9, p. 207 (Nagasaki).
Distr.: Japan (Shikoku, Kyushu).

P. yoshidai KIUCHI, 1972 キウチテントウダマン

KIUCHI, 1972, Kontyû, Tokyo, 40, p. 160 (Mt. Tsurugi, Tokushima Pref.).
Distr.: Japan (Shikoku).

Subfamily TROCHOIDEINAE ヒゲブテントウダマン亜科

Genus *Trochoideus* WESTWOOD, 1833 ヒゲブテントウダマン属

Trochoideus WESTWOOD, 1833, Trans. Linn. Soc. London, 16, p. 673 (type-species: *Paussus cruciatus* DALMAN, 1825).

Trochoides CHAPIUS, 1876, Gen. Coléopt., p. 147.

Pseudopaussus SCHULTZE, 1916, Philipp. J. Sci., 11, p. 291.

T. desjardinsi GUÉRIN-MÉNEVILLE, 1838 ヒゲブテントウダマン

GUÉRIN-MÉNEVILLE, 1838, Rev. Zool., p. 22 (Madagascar?).

termitophilus ROEPKE, 1919, Treubia, 1, p. 34.

monstrosus SCHULZE, 1916, Philipp. J. Sci., 11, p. 292 (*Pseudopaussus*).

rouyeri PIC, 1922, Mém. Exot.-Ent., (35), p. 8.

particularis PIC, 1922, l. c., p. 8

amphora CANTOR, 1844, J. Proc. ent. Soc. London, 1844, p. 118.

Distr.: Bonin Is. (Chichijima, Hahajima), Botel-Tobago Is. Philippines, Java, Borneo, Malay Penn., Burma, India, Ceylon, Andaman Is., Seychelles Is., Cocos-Keeling Is., Batchian, Rodrigues Is., E. Africa, Madagascar, Samoa, Mariana Is., Caroline Is., Trak Is., Fiji Is., Mauritius, New Guinea.

Subfamily STENOTARSINAE ムクゲテントウダマン亜科

Genus *Stenotarsus* PERTY, 1832 ムクゲテントウダマン属

Stenotarsus PERTY, 1832, Dilect. Animal Artic., p. 112 (type-species: *Stenotarsus brevicollis* PERTY, 1832).

Quirinus THOMSON, 1857, Archives Ent., 1, p. 157.

Systaechea GORHAM, 1890, Biol. Cent. Amer. Coleopt., 7, p. 132.

Stenotarsoides CSIKI, 1900, Termész. Füzetek, 23, p. 401.

S. chrysomelinus GORHAM, 1887 チャバネムクゲテントウダマン

GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 645 (Ichiuchi, Nara).

Distr.: Japan (Honshu, Shikoku, Kyushu).

S. internexus GORHAM, 1887 クロスジムクゲテントウダマン

GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 645 (Nagasaki, Kashiwagi).

Distr.: Japan (Honshu, Kyushu), Ryukyu Is. (Nakanoshima, Taketomi-jima).

S. nakanoshimensis NAKANE, 1959 トカラムクゲテントウダマン

NAKANE, 1959, Sci. Rept. Kyoto Pref. Univ., (Nat. Sci. & Liv. Sci.), 3A, p. 60 (Naka-

noshima, Tokara).

Distr.: Ryukyu Is. (Nakanoshima).

S. oshimanus NAKANE, 1981 アマミムクゲテントウダマン

NAKANE, 1981, Kita-Kyūshū no Konchū, **28**, p. 55, 57-58 (Amami-Ōshima).

Distr.: Ryukyu Is. (Amami-Ōshima).

S. ryukyuensis CHŪJŌ et KIUCHI, 1974 キイロムクゲテントウダマン

CHŪJŌ & KIUCHI, 1974, Bull. Jap. ent. Acad., **8**, p. 5 (Amami-Ōshima).

peguensis OHTA, 1931 (nec GORHAM, 1896), J. Fac. Agr. Hokkaido Univ., **30**, p. 228 (Okinawa).

Distr.: Ryukyu Is. (Nakanoshima, Amami-Ōshima, Okinawa).

S. kurosai NAKANE, 1981 コゲチャムクゲテントウダマン

NAKANE, 1981, Kita-Kyūshū no Konchū, **28**, p. 55, 56-57 (Ishigaki-jima).

Distr.: Ryukyu Is. (Ishigaki-jima).

Genus *Chondria* GORHAM, 1887 アンボソテントウダマン属

Chondria GORHAM, 1887, Proc. zool. Soc. London, **1887**, p. 651 (type-species: *Chondria lutea* GORHAM, 1887).

C. lutea GORHAM, 1887 キイロアンボソテントウダマン

GORHAM, 1887, Proc. zool. Soc. London, **1887**, p. 652 (Nagasaki).

Distr.: Japan (Honshu, Kyushu).

Genus *Ectomychus* GORHAM, 1887 ケブカテントウダマン属

Ectomychus GORHAM, 1887, Proc. zool. Soc. London, **1887**, p. 646 (type-species: *Ectomychus basalis* GORHAM, 1887).

E. basalis GORHAM, 1887 カタバニケブカテントウダマン

GORHAM, 1887, l. c., p. 646 (Kawachi, Miyanoshita, Kurigahara, Sapporo).

Distr.: Japan (Hokkaido, Honshu, Kyushu, Tsushima).

E. musculus (GORHAM, 1887) クロモンケブカテントウダマン

GORHAM, 1887, l. c., p. 645 (*Stenotarsus*) (Nagasaki, Kashiwagi).

nakanei CHŪJŌ, 1941, Mushi, Fukuoka, **13**, p. 81 (Oku-tama).

Distr.: Japan (Honshu, Shikoku, Kyushu).

E. nigriclavus (GORHAM, 1873) チャイロケブカテントウダマン

GORHAM, 1873, Ent. month. Mag., **9**, p. 206 (*Stenotarsus*) (Nagasaki).

Distr.: Japan (Honshu, Kyushu).

E. sakaii CHŪJŌ et KIUCHI, 1974 ムネアカケブカテントウダマン

CHŪJŌ & KIUCHI, 1974 Bull. Jap. ent. Acad., **8**, p. 4 (Amami-Ōshima).

Distr.: Ryukyu Is. (Nakanoshima, Amami-Ōshima, Tokunoshima).

Genus *Saula* GERSTAECKER, 1858 キイロテントウダマン属

Saula GERSTAECKER, 1858, Monogr. Endom., p. 223 (type-species: *Saula nigripes* GERSTAECKER, 1858).

S. japonica GORHAM, 1874 キイロテントウダマン

GORHAM, 1874, Ent. month. Mag., **10**, p. 224 (Nagasaki).

nigripes? : GORHAM, 1873, Ent. month. Mag., **9**, p. 206.

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Genus *Danae* REICHE, 1847 ダナエテントウダマン属

- Danaë* REICHE, 1847, in FERRET & GALINIER, Voyage en Abyssinie, 3, p. 408 (type-species: *Danaë rufula* REICHE, 1847).
- Oediarthrus* GERSTACKER, 1858, Monogr. Endom., p. 344.
- Rhabduchus* GORHAM, 1873, Ent. month. Mag., 9, p. 257.
- Coniopoda* GORHAM, 1873, 1. c., p. 257.
- D. castanea* SASAJI, 1978 カバイロダナエテントウダマン
SASAJI, 1978, Mem. Fac. Educ. Fukui Univ., Sect. II (Nat. Sci.), (28), p. 23 (Nagasaki).
Distr.: Japan (Kyushu).
- D. denticornis* (GORHAM, 1873) オオダナエテントウダマン
GORHAM, 1873, Ent. month. Mag., 9, p. 257 (*Rhabduchus*) (Nagasaki).
Distr.: Japan (Honshu, Kyushu, Tsushima).
- D. orientalis* (GORHAM, 1873) トウヨウダナエテントウダマン
GORHAM, 1873, Ent. month. Mag., 9, p. 205 (*Coniopoda*) (Hiogo).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).
- D. sibatai* NAKANE, 1958 オニダナエテントウダマン
NAKANE, 1958, Akitu, Kyoto, 7, p. 35 (Nara).
Distr.: Japan (Honshu).

Subfamily EUMORPHINAE オオテントウダマン亜科
Genus *Lycoperdina* LATREILLE, 1807 ツヤテントウダマン属

- Lycoperdina* LATREILLE, 1807, Gen. Crust. et Insect., 3, p. 73 (type-species: *Galleruca bovista* FABRICIUS, 1792).
- Lycoperdinodes* ARROW, 1923, Trans. ent. Soc. London, 1923, p. 485.
- Lycoperdinella* ARROW, 1920, 1. c., 1920, p. 23.
- Falsoylaia* PIC, 1945, Échange, 501, p. 11.

Subgenus *Gorgia* MULSANT, 1846

- Gorgia* MULSANT, 1846, Hist. nat. Coléopt. France, Sulcicolles, p. 17 (type-species: *Silpha succincta* LINNAEUS, 1758).
- L. (G.) castaneipennis* GORHAM, 1874 クリバネツヤテントウダマン
GORHAM, 1874, Ent. month. Mag., 11, p. 151 (Kobe).
Distr.: Japan (Honshu, Shikoku, Kyushu, Sado).
- L. (G.) dux* GORHAM, 1873 フチトリツヤテントウダマン
GORHAM, 1873, Ent. month. Mag., 9, p. 205 (Hiogo).
Distr.: Japan (Hokkaido, Honshu, Shikoku).
- L. (G.) mandarinae* GERSTAECKER, 1858 セグロツヤテントウダマン
GERSTAECKER, 1858, Monogr. Endom., p. 212 (Hongkong).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Ryukyu Is. (Nakanoshima),
Formosa, China, Hongkong, Tonkin.

Genus *Ancylopus* COSTA, 1854 ヨツボシテントウダマン属

- Ancylopus* COSTA, 1854, Fauna Regno Napoli, Coleopt. I. Endomichidei, p. 14 (type-species: *Endomychus melanocephalus* OLIVIER, 1808).

- A. pictus** WIEDEMANN, 1823 ヨツボシテントウダマン
 WIEDEMANN, 1823, Zool. Mag., 2, p. 78 (*melanocephalus* var.).
 Distr.: Japan, South Asia, Borneo, New Guinea.
- subsp. *asiaticus* STROHECKER, 1972 (ヨツボシテントウダマン)
 STROHECKER, 1972, Pacif. Ins., 14, p. 704 (Fukien).
villiersi DAJOZ, 1973, Bull. Mus. Hist. nat. Paris, (3), Zool., 64, p. 1046 (Nilgiri Hills).
melanocephalus: GORHAM, 1873 (nec OLIVIER, 1808), Ent. month. Mag., 9, p. 205 (Japan).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Formosa, China, India.
- A. punji** PIC, 1926 ベニヨツボシテントウダマン
 PIC, 1926, Mém. Exot.-Ent., (45), p. 10 (Tonkin).
 Distr.: Japan, China, Tonkin, Tibet.
- subsp. *punji* PIC, s. str.
 PIC, 1926, l. c., p. 10 (Tonkin).
 Distr.: Japan (Honshu), China, Tonkin, Tibet.

Genus *Mycetina* MULSANT, 1846 ムナビロテントウダマン属

- Mycetina* MULSANT, 1846, Hist. nat. Coléopt. France., Sulcicolles, p. 15 (type-species: *Chrysomela cruciata* SCHALLER, 1783).
Phaeomychus GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 649.
Mycetinina PIC, 1929, Rev. Sci. Bourbonnais, p. 35.
- M. amabilis** GORHAM, 1873 キボシテントウダマン
 GORHAM, 1873, Ent. month. Mag., 9, p. 206 (Nagasaki).
humeropecta NAKANE, 1958, Akitu, Kyoto, 7, p. 33 (as a forma).
takahashii NAKANE, 1958, l. c., p. 33 (as a forma).
 Distr.: Saghalien, Japan (Hokkaido, Honshu, Shikoku, Kyushu).
- M. ancoriger** GORHAM, 1873 イカリモンテントウダマン
 GORHAM, 1873, Ent. month. Mag., 9, p. 206 (Nagasaki).
 Distr.: Japan (Honshu, Shikoku, Kyushu).
- M. laticollis** GORHAM, 1887 ムナビロテントウダマン
 GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 644 (Kashiwagi, Nara, Maiyasan).
 Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima).
- M. rufipennis** (MOTSCHULSKY, 1860) ベニバネテントウダマン
 MOTSCHULSKY, 1860, Étud. ent., 1860, p. 18 (Japonia).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Genus *Indalmus* GERSTAECKER, 1858 ヒラタテントウダマン属

- Indalmus* GERSTAECKER, 1858, Monogr. Endom., p. 158 (type-species: *Eumorphus kirbyanus* LATREILLE, 1807).
Mycella CHAPIUS, 1876, in LACORDAIRE, Gen. Coléopt., 12, p. 424.
- I. quadripunctatus** (OHTA, 1931) ヨツボシヒラタテントウダマン
 OHTA, 1931, J. Fac. Agr. Hokkaido Imp. Univ., 30, p. 222 (*Encymon*) (Formosa: Baibara).
 Distr.: Ryukyu Is. (Ishigaki-jima), Formosa.

Genus *Eumorphus* WEBER, 1801 オオテントウダマン属

- Eumorphus* WEBER, 1801, *Observ. Ent.*, p. 31 (type-species: *Erotylus quadriguttatus* ILLIGER, 1800=*Eumorphus sumatrae* WEBER, 1801).
Eumorphoides GUÉRIN-MÉNEVILLE, 1858, *Rév. Mag. Zool.*, (2), **10**, p. 12.
Enaisimus GUÉRIN-MÉNEVILLE, 1858, 1. c., p. 16.
Haplomorphus GUÉRIN-MÉNEVILLE, 1858, 1. c., p. 18.
Heterandrus GUÉRIN-MÉNEVILLE, 1858, 1. c., p. 26.

- E. quadriguttatus* (ILLIGER, 1800) タイワンオオテントウダマン
 ILLIGER, 1800, *Arch. f. Zool.*, (1), **2**, p. 124, pl. 1, fig. 4 (*Erotylus*) (India?).
sumatrae WEBER, 1801, *Observ. Ent.*, p. 59.
immarginatus FABRICIUS, 1801, *Syst. Eleuth.*, **2**, p. 11.
laevis GORHAM, 1874, *Trans. ent. Soc. London*, **1874**, p. 305 (*Pedanus*).
 Distr.: Japan, Formosa, China, Philippines, Malaysia, Indonesia, Andamans, Burma, India, Ceylon.
- subsp. *pulchripes* GERSTAECKER, 1857 (タイワンオオテントウダマン)
 GERSTAECKER, 1857, *Archiv Naturg.*, **23**, p. 228 (full species) (Ceylon).
formosanus PIC, 1930, *Mél. Exot.-Ent.*, (55), p. 8 (subsp. of *quadriguttatus*).
baibaranus OHTA, 1931, *J. Fac. Agr. Hokkaido Univ.*, **30**, p. 219 (*Engonius*).
taitoensis OHTA, 1931, 1. c., p. 219 (*Engonius*).
 Distr.: Japan (Tsushima), Formosa, China, Hainan Is., India, Ceylon.

Subfamily ENDOMYCHINAE テントウダマン亜科

Genus *Endomychus* PANZER, 1795 テントウダマン属

- Endomychus* PANZER, 1795, *Ent. Taschenbuch*, p. 175 (type-species: *Chrysomela coccinea* LINNAEUS, 1758).
Cyanauges GORHAM, 1874, *Ent. month. Mag.* **11**, p. 54.
Caenomychus LEWIS, 1893, *Entomologist*, **26**, p. 153.
- E. gorhami* (LEWIS, 1874) ルリテントウダマン
 LEWIS, 1874, *Ent. month. Mag.*, **11**, p. 55 (*Cyanauges*) (Kawatchi).
violaceipennis MADER, 1941, *Ent. Bl.*, **37**, p. 251 (*Caenomychus*).
 Distr.: Japan.
- subsp. *gorhami* (LEWIS, 1874), s. str.
 LEWIS, 1874, 1. c., p. 55 (*Cyanauges*) (Kawatchi).
 Dist.: Japan (Hokkaido, Honshu, Shikoku).
- subsp. *kyushuensis* SASAJI, 1978
 SASAJI, 1978, *Mem. Fac. Fukui Univ., Sect. II (Nat. Sci.)*, (28), p. 25 (Mt. Hikosan, Fukuoka Pref.).
 Distr.: Japan (Kyushu).
- E. hiranoi* SASAJI, 1978 ヒラノクロテントウダマン
 SASAJI, 1978, *Mem. Fac. Fukui Univ. Sect. II (Nat. Sci.)*, (28), p. 27 (Miyanoshiba in Hakone).
 Distr.: Japan (Honshu).
- E. nigropiceus* (GORHAM, 1887) ウスグロテントウダマン
 GORHAM, 1887, *Proc. zool. Soc. London*, **1887**, p. 651 (Kashiwagi). (*Cyanauges*).
ohbayashii NAKANE, 1951, *Ins. Matsumurana*, **17**, p. 116.

shirahatai NAKANE, 1951, 1. c., p. 117 (subsp. of *ohbayashii*).

Distr.: Japan (Honshu, Shikoku, Kyushu).

E. plagiatus (GORHAM, 1887) キスジテントウダマシ

GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 650 (*Cyanauges*) (Yuyama, Hitoyoshi).

Distr.: Japan (Shikoku, Kyushu).

E. quadra (GORHAM, 1887) ハバビロテントウダマシ

GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 651 (*Cyanauges*) (Kashiwagi).

Distr.: Japan (Honshu).

Genus ***Bolbomorphus*** GORHAM, 1887 セダカテントウダマシ属

Bolbomorphus GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 647 (type-species:

Bolbomorphus gibbosus GORHAM, 1887).

B. gibbosus GORHAM, 1887 セダカテントウダマシ

GORHAM, 1887, Proc. zool. Soc. London, 1887, p. 647 (Kashiwagi).

Distr.: Japan (Honshu).