

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 22

March 30, 1983

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family CHRYSOMELIDAE ハムシ科
(CHLAMISINAE, LAMPROSOMATINAE, SYNETINAE)

By

Shinsaku KIMOTO

Subfamily CHLAMISINAE コブハムシ亜科
Genus *Chlamys* RAFINESQUE, 1815 コブハムシ属

- Chlamys* KNOCH, 1801 (nec RODING in BOLTEN, 1798), Neue Beytr. Insectenk., 1, p. 122.
Chlamys RAFINESQUE, 1815, Analyse Nat. Tab. l'Univ. Corps Organ, p. 16 (type-species:
Bruchus gibbosa FABRICIUS, 1777).
Exema LACORDAIRE, 1848, Monogr. Phytoph., 2, p. 634, 844.
Myochlamys IHERING, 1905 (nec FAIRMAIRE, 1876), Rev. Mus. Paulista, 6, p. 642 (n. n.
for *Chlamys* KNOCH, nec RODING, 1798).
Arthrochlamys IHERING, 1907, Ann. Mus. Nac. Buenos Aires, 14, p. 251 (n. n. for
Myochlamys IHERING, nec FAIRMAIRE, 1876).
Boloschensis JACOBSON, 1924, Rev. russe Ent., 18, p. 239 (n. n. for *Chlamys* KNOCH,
nec RODING, 1798).
C. consimilis (CHÛJÔ, 1942) カンワコブハムシ
CHÛJÔ, 1942, Trans. nat. Hist. Soc. Formosa, 32, p. 86, fig. 2 (*Exema*) (Handakogen
in Oita Pref., Nojiri in Nagano Pref., Hachinohe in Aomori Pref.)
Distr.: Japan (Honshu, Kyushu).
C. diminutus (GRESSITT, 1942) ヒメコブハムシ
GRESSITT, 1942, Lingnan Sci. J., 20, p. 359, pl. 21, fig. 2 (*Chlamys*) (China).
Distr.: Japan (Kyushu), Korea, China, Taiwan.
C. geniculatus (JACOBY, 1896) アマミコブハムシ
JACOBY, 1896, Entomolog., 29, p. 6. (*Chlamys*) (Amami-Ôshima).

Distr.: Japan (Yakushima, Tanegashima), Ryukyu Is. (Amami-Ōshima, Tokunoshima, Okinawa).

C. interjectus (BALY, 1873) ミズキコブハムシ

BALY, 1873, Trans. ent. Soc. London, 1873, p. 84 (*Chlamys*) (Nagasaki).

cirsicola CHŪJŌ, 1942, Trans. nat. Hist. Soc. Formosa, 32, p. 93, fig. 3 (*Exema*) (Mt. Inunaki-yama and Mt. Hiko in Fukuoka Pref.).

Distr.: Japan (Shikoku, Kyushu).

C. japonicus (JACOBY, 1885) ハバビロコブハムシ

JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 198, pl. 11, fig. 5 (*Chlamys*) (Kiga, Fukushima).

Distr.: Japan (Honshu, Kyushu, Yakushima), Ryukyu Is. (Amami-Ōshima, Ishigakijima, Iriomote-jima), China.

C. laticollis (CHŪJŌ, 1942) ツツジコブハムシ

CHŪJŌ, 1942, Trans. nat. Hist. Soc. Formosa, 32, p. 82, fig. 1 (*Exema*) (Mt. Hiko in Fukuoka Pref., Tsushima).

Distr.: Japan (Honshu, Kyushu, Tsushima).

C. lewisii (BALY, 1873) ツバキコブハムシ

BALY, 1873, Trans. ent. Soc. London, 1873, p. 83 (*Chlamys*) (Nagasaki).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan.

C. spilotus (BALY, 1873) ムシクソハムシ

BALY, 1873, Trans. ent. Soc. London, 1873, p. 85 (*Chlamys*) (Japan).

Distr.: Japan (Honshu, Sado, Shikoku, Kyushu).

C. yakushmanus OHNO, 1961 ヤクシマコブハムシ

OHNO, 1961, Ent. Rev. Japan, 13, p. 1 (Yakushima).

Distr.: Japan (Shikoku, Yakushima), Ryukyu Is. (Okinawa, Ishigakijima).

Subfamily LAMPROSOMATINAE ツヤハムシ亜科

Genus *Oomorplus* CURTIS, 1831 ヒメツヤハムシ属

Oomorplus CURTIS, 1831, Brit. Ent., 8, p. 374 (type-species: *Byrrhus concolor* STURM, 1807).

Subgenus *Oomorplus* CURTIS, 1831, s. str.

O. (Oomorplus) japanus JACOBY, 1885 ヒメツヤハムシ

JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 197 (Oyama, Ichiuchi).

japanus f. *chujoi* OHNO, 1961, Ent. Rev. Japan, 13, p. 48 (Honshu).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Oomorphoides* MONRÓS, 1956 ツヤハムシ属

Oomorphoides MONRÓS, 1956, Rev. Agron. Noro. Argent., 2, p. 54 (type-species: *Lamprosoma cupreatum* BALY, 1873).

O. cupreatus (BALY, 1873) ドウガネツヤハムシ

BALY, 1873, Trans. ent. Soc. London, 1873, p. 82 (*Lamprosoma*) (Nagasaki).

cupreatus f. *yusai* OHNO, 1961, Ent. Rev. Japan, 13, p. 49 (Honshu, Shikoku, Kyushu).

Distr.: Japan (Hokkaido, Honshu, Hachijo-jima, Shikoku, Kyushu), Korea.

- O. loochooensis* CHÛJÔ, 1958 リュウキュウツヤハムシ
 CHÛJÔ, 1958, Mem. Fac. Lib. Arts & Educ., Kagawa Univ., 2 (64), p. 4
 (Yurudji in Okinawa).
loochooensis yakushimensis OHNO, 1961, Ent. Rev. Japan, 13, p. 48, 50 (Yakushima).
 Distr.: Japan (Yakushima), Ryukyu Is. (Amami-Ôshima, Tokunoshima).
- O. nigrocaeruleus* (BALY, 1873) アオグロツヤハムシ
 BALY, 1873, Trans. ent. Soc. London, 1873, p. 83 (*Lamprosoma*) (Nagasaki).
 Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima).
- O. okinawensis* (CHÛJÔ, 1935) オキナワツヤハムシ
 CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, 25, p. 75 (*Lamprosoma*) (Naze, Amami-
 Ôshima).
cupreatus kurosawai NAKANE, 1958, Sci. Rept. Saikyo Univ., (A), 2, p. 304, figs. 2,
 4, 6 (Yakushima).
 Distr.: Japan (Yakushima), Ryukyu Is. (Amami-Ôshima, Tokunoshima).
- O. sakishimanus* KIMOTO et GRESSITT, 1966 サキシマツヤハムシ
 KIMOTO & GRESSITT, 1966, Pacif. Ins., 8, p. 472, 499 (Nakaragawa, Iriomote Is.).
 Distr.: Ryukyu Is. (Iriomote-jima).

Subfamily SYNETINAE ホソハムシ亜科

Genus *Syneta* ESCHSCHOLTZ, 1835 カバノキハムシ属

- Syneta* ESCHSCHOLTZ, 1835, in DEJEAN, Cat. Coléopt., 2 ed., fasc. 4, p. 359 (type-species:
Crioceris betulae FABRICIUS, 1792).
- S. adamsi* BALY, 1877 カバノキハムシ
 BALY, 1877, Ann. Mag. nat. Hist., (4), 20, p. 378 (Tsushima, Vladimir Bay,
 Manchuria).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima), Kuriles,
 Sakhalin, Manchuria, E Siberia.
- S. brevitibialis* KIMOTO, 1971 ヒロアシカバハムシ
 KIMOTO, 1971, Bull. Osaka Mus. nat. Hist., 25, p. 2 (Syonoiwaya—Mt. Daifugen,
 and Gomadanzan, Nara Pref.).
 Distr.: Japan (Honshu).
- S. major* NAKANE, 1963 オオカバハムシ
 NAKANE, 1963, Fragm. Coleopt., (5), p. 19 (Kamikochi, Nikko).
 Distr.: Japan (Honshu).