

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 28

April 30, 1985

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family CHRYSOMELIDAE ハムシ科
(GALERUCINAE ヒゲナガハムシ亜科)

By

Shinsaku KIMOTO

Subfamily GALERUCINAE ヒゲナガハムシ亜科

Genus *Galeruca* MÜLLER, 1764 オオアラメハムシ属

Galeruca MÜLLER, 1764, Fauna Insect. Fridrichsdalina, p. 14 (type-species: *Chrysomela tanacetii* LINNÉ, 1758)

Adimonia LAICHARTING, 1781, Verz. Tyrol. Ins., p. 190.

Emarhopa WEISE, 1886, Naturg. Ins. Dtschl., 6, p. 657.

Haptoscelis WEISE, 1886, *ibid.*, p. 658.

Galerima REITTER, 1903, Wien. ent. Ztg., 22, p. 133.

Galerotoma REITTER, 1903, *ibid.*, p. 139.

Fassatia HAVELKA, 1954, Acta Soc. Ent. Cechosloveniae, 31, p. 115.

G. extensa MOTSCHULSKY, 1861 ネギオオアラメハムシ

MOTSCHULSKY, 1861, Étud. ent., 10, p. 22 (Japan).

banghaasi WEISE, 1894, Dtsch. ent. Z., 1894, p. 168 (Yokohama).

Distr.: Japan (Hokkaido, Honshu), Korea, NE China, E. Siberia.

G. spectabilis FALDERMANN, 1837 スジグロオオハムシ

FALDERMANN, 1837, Nouv. Mem. Soc. imp. Natur. Moscou, 5, p. 326, pl. 12, fig. 4
(S. Russia).

Distr.: Japan (Honshu), China, C. Asia, S. Russia.

G. vicina SOLSKY, 1872 アザミオオハムシ

SOLSKY, 1872, Horae Soc. ent. Ross., 8, p. 252 (Vladivostock).

dahli var. *japonica* WEISE, 1894, Dtsch. ent. Z., 1894, p. 168 (Yokohama).

- KIMOTO, 1963, *Fragm. Col.*, Kyoto, (4), p. 17 (Tokyo, Nagano, Nara, Gifu, Osaka, Kyoto).
Distr.: Japan (Honshu).
- P. fuscipennis* (JACOBY, 1885) イタヤハムシ
JACOBY, 1885, *Proc. zool. Soc. London*, 1885, p. 746 (*Galerucella*) (Awomori).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yakushima), Korea, E. Siberia.
- P. humeralis* (CHEN, 1942) サンゴジュハムシ
CHEN, 1942, *Notes d'Ent. Chinoise*, 9, p. 17 (*Galerucella*) (China).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Ryukyu Is, (Okinawa), China.
- P. kawashimai* KIMOTO, 1964 カワシマケブカハムシ
KIMOTO, 1964, *J. Fac. Agr. Kyushu Univ.*, 13, p. 297 (Kanagawa).
Distr.: Japan (Honshu).
- P. konishii* KIMOTO, 1964 コニシケブカハムシ
KIMOTO, 1964, *J. Fac. Agr. Kyushu Univ.*, 13, p. 298 (Hokkaido).
Distr.: Japan (Hokkaido)
- P. lineola* (FABRICIUS, 1781) ハシバミハムシ
FABRICIUS, 1781, *Spec. Ins.*, 1, p. 149 (*Chrysomela*) (Europe).
Distr.: Japan (Hokkaido), NE China, Sakhalin, Siberia, Europe.
- P. maculicollis* (MOTSCHULSKY, 1853) ニレハムシ
MOTSCHULSKY, 1853, *Étud. ent.*, 2, p. 49 (*Galeruca*) (N. China).
Distr.: SE Siberia, N China, Korea, Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu).
- P. nigricornis* OHNO, 1961 クロヒゲケブカハムシ
OHNO, 1961, *Kontyû*, 30, p. 27 (Amami-Oshima).
Distr.: Ryukyu Is. (Amami-Oshima).
- P. nigromarginata* (JACOBY, 1885) クロヘリウスチャハムシ
JACOBY, 1885, *Proc. zool. Soc. London*, 1885, p. 743, pl. 46, fig. 10 (*Galerucella*) (Japan).
Distr.: Japan (Honshu), S. China, Hainan.
- P. semifulva* (JACOBY, 1885) アカタデハムシ
JACOBY, 1885, *Proc. zool. Soc. London*, 1885, p. 745, pl. 46, fig. 11 (*Galerucella*) (Kiga).
modesta JACOBY, 1885, *ibid.*, p. 745 (*Galerucella*) (Nikko).
japonica WEISE, 1922, *Tijdschr. ent.*, 65, p. 67 (*Lochmaea*) (Kioto).
Dist.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Yakushima), Taiwan, NE China, E. Siberia.
- P. seminigra* (JACOBY, 1885) カエデハムシ
JACOBY, 1885, *Proc. zool. Soc. London*, 1885, p. 744 (*Galerucella*) (Yagohara).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).
- P. takeii* (CHÛJÔ, 1950) オオサクラケブカハムシ
CHÛJÔ, 1950, *Kontyû*, 18, p. 114, 1 fig. (*Lochmaea*) (Gumma).
Distr.: Japan (Honshu).

P. tibialis (BALY, 1874) エノキハムシBALY, 1874, Trans. ent. Soc. London, 1874, p. 176 (*Galleruca*) (Nagasaki).*nigromembris* FAIRMAIRE, 1888, Ann. Soc. ent. Belg., 33, p. 41 (China).

Distr.: Japan (Honshu, Shikoku, Kyushu), C. China.

P. yasumatsui KIMOTO, 1964 ヤスマツケブカハムシ

KIMOTO, 1964, J. Fac. Agr. Kyushu Univ., 13, p. 301 (Okinawa, Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima, Okinawa).

P. yoshimotoi KIMOTO et GRESSITT, 1966 ヨシモトケブカハムシ

KIMOTO et GRESSITT, 1966, Pac. Ins., 8, p. 521 (Okinawa).

Distr.: Ryukyu Is. (Okinawa).

Genus *Issikia* CHÛJÔ, 1961 アマミウスバハムシ属*Issikia* CHÛJÔ, 1961, Pub. ent. Lab. Univ. Osaka Pref., (6), p. 87 (type-species: *Galeruca issikii* CHÛJÔ, 1961).**I. issikii** (CHÛJÔ, 1961) アマミウスバハムシCHÛJÔ, 1961, Pub. ent. Lab. Univ. Osaka Pref., (6), p. 86 (*Galeruca*) (Amami-Oshima).

Distr.: Ryukyu Is. (Kikaigashima, Amami-Oshima, Okinawa).

Genus *Aulacophora* CHEVROLAT, 1837 ウリハムシ属*Aulacophora* CHEVROLAT, 1837, in DEJEAN, Cat. Col., ed. 3, p. 402 (type-species: *Galeruca quadraria* OLIVIER, 1808).*Raphidopalpa* CHEVROLAT, 1837, in DEJEAN, Cat. Col., ed. 3, 402.*Acuteipalpa* ROSENHAUER, 1857, Tiere Andalusia, p. 327.*Ceratia* CHAPUIS, 1876, C. R. Soc. ent. Belg., 19, p. 100.*Triplatyps* FAIRMAIRE, 1877, Petit. Nouv. ent., 2 p. 186.*Pachypalpa* WEISE, 1892, Dtsch. ent. Z., 1892, p. 392.*Orthaulaca* WEISE, 1892, *ibid.*, p. 396.*Cerania* WEISE, 1892, *ibid.*, p. 396.*Spaerarthra* WEISE, 1892, *ibid.*, p. 396.*Paraulacophora* CSIKI, 1953, Ann. Hist. Natur. Mus. Hungarici, n. ser., 3, p. 131.**A. bicolor** (WEBER, 1801) フタイロウリハムシWEBER, 1801, Obs. Ent., p. 56 (*Galleruca*) (Sumatra).*bicolor* FABRICIUS, 1801, Syst. Eleut., 1, p. 482 (*Galleruca*) (Sumatra).*haemorrhoea* FABRICIUS, 1803, in ILLIGER, Mag. Ins., 2, p. 293 (n. n. for *bicolor* FABRICIUS, 1801).*sexnotata* OLIVIER, 1808, Ent., 6, p. 627, pl. 2, fig. 29 (*Galeruca*) (Timor).*hemichroa* GISTEL, 1857, Naturg. Tierreichs, ed. 2, p. 540.*sexnotata* CHAPUIS, 1876, C. R. Soc. ent. Belg., 19, p. 100 (Philippines).*semiopaca* JACOBY, 1886, Ann. Mus. Civ. Genova, 24, p. 51 (Sumatra).

Distr.: Ryukyu Is. (Amami-Oshima, Okinoerabu, Okinawa, Miyako, Ishigaki, Iriomote, Yonaguni), Taiwan, SE Asia.

A. femoralis (MOTSCHULSKY, 1857) ウリハムシMOTSCHULSKY, 1857, Étud. ent., 6, p. 37 (*Raphidopalpa*) (Japan).*similis*: BALY, 1886, J. Linn. Soc. London, 20, pp. 3, 5, 16 (Japan).*coffaeae*: ALLARD, 1888, Ann. Soc. ent. France, (6), 8, pp. 306, 319 (Japan).

chinensis WEISE, 1892, Dtsch. ent. Z., 1892, p. 395 (*Rhaphidopalpa*) (Shanghai).
Distr.: Japan (Honshu, Hachijo, Awashima, Sado, Oki, Shikoku, Kyushu, Tsushima, Yakushima, Kuchinoerabu), Ryukyu Is. (Tokara, Kikai, Amami-Oshima, Tokunoshima, Okinoerabu, Yoron, Okinawa, Miyako, Ishigaki, Iriomote, Yonaguni, Daito-jima), Korea, China, Taiwan.

A. lewisii BALY, 1886 ヒメクロウリハムシ

BALY, 1886, J. Linn. Soc. London, 20, pp. 5, 24 (China).

cattigarensis WEISE, 1892, Dtsch. ent. Z., 1892, p. 397 (*Orthaulaca*) (Japan, Shanghai).

intermedia JACOBY, 1892, Ann. Mus. Civ. Genova, 32, p. 942 (Burma).

Distr.: Japan (Yakushima), Ryukyu Is. (Amami-Oshima, Okinawa, Miyako, Ishigaki, Iriomote), Taiwan, SE Asia, India, Ceylon.

A. loochooensis CHÛJÔ, 1957 ルリバネウリハムシ

CHÛJÔ, 1957, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., 2 (52), p. 2 (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima).

A. nigripennis MOTSCHULSKY, 1857 クロウリハムシ

MOTSCHULSKY, 1857, Étud. ent., 6, p. 38 (Japan).

Distr.: Japan, Ryukyu Is., Taiwan, China, Korea, E. Siberia.

subsp. *nigripennis* MOTSCHULSKY, 1857, s. str.

MOTSCHULSKY, 1857, Étud. ent., 6, p. 38 (Japan).

Distr.: Japan (Honshu, Awashima, Sado, Oki, Shikoku, Kyushu, Tsushima, Gotô, Yakushima), Ryukyu Is. (Kikaigashima), Taiwan, China, Korea, E. Siberia.

subsp. *nitidipennis* CHÛJÔ, 1935

CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, 25, p. 83 (Okinawa, Amami-Oshima, Iriomote).

nigripennis insularis OHNO, 1963, Bull. Biogeograph. Soc. Japan, 22, p. 88 (Izu Is.).

Distr.: Japan (Miyake, Mikura), Ryukyu Is. (Amami-Oshima, Tokunoshima, Okinawa, Iriomote).

Genus **Oides** WEBER, 1801 キベリハムシ属

Oides WEBER, 1801, Obs. Ent., p. 26 (type-species: *Chrysomela bipunctata* FABRICIUS, 1781).

Adorium FABRICIUS, 1801, Syst. Ent., 1, p. 409.

Isosoma BILLBERG, 1820, Enum. Insect. Mus. Billberg, p. 56.

Ochralea CHEVROLAT, 1837, in DEJEAN, Cat. Col., ed. 3, p. 399.

Rhombopalpa CHEVROLAT, 1837, *ibid.*, p. 399.

Callipela DEJEAN, 1837, Cat. Col., ed. 3, p. 399.

Boisduvalia MONTROUZIER, 1855, Ann. Soc. Agr. Lyon, 7, p. 72.

Botanctona FAIRMAIRE, 1877, Petit. Nouv. ent., 2, p. 185.

O. bowringii (BALY, 1863) キベリハムシ

BALY, 1863, Trans. ent. Soc. London, (3), 1, p. 623 (*Adorium*) (Hongkong).

elegans LABOISSIÈRE, 1919, Bull. Soc. ent. France, 1919, p. 161 (Tonkin).

Distr.: Japan (Honshu), Korea, S. China, Vietnam.

Genus **Clerotilia** JACOBY, 1885 キアシヒゲナガアオハムシ属

Clerotilia JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 751 (type-species: *Clerotilia*

flavomarginata JACOBY, 1885).

- C. flavomarginata** JACOBY, 1885 キアシヒゲナガアオハムシ
 JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 751, pl. 46, fig. 12 (Tsumago, Nara).
 Distr.: Japan (Honshu, Shikoku, Kyushu), E. China.

Genus *Japonitata* STRAND, 1935 クロセスジハムシ属

Japonia WEISE, 1922, Tijdschr. ent., **65**, p. 70 (nec GOULD, 1859) (type-species: *Phyllobrotica nigrita* JACOBY, 1885)

Japonitata STRAND, 1935, Folia Zool. Hydrobiol., **7**, p. 294 (n. n. for *Japonia* WEISE).

- J. nigrita** (JACOBY, 1885) クロセスジハムシ
 JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 742, pl. 46, fig. 3 (*Phyllobrotica*)
 (Osaka).
 Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Euliroetis* OGLOBLIN, 1936 ハラグリハムシ属

Euliroetis OGLOBLIN, 1936, Fauna USSR, **26** (1), pp. 197, 371, 403 (type-species: *Aenidea ornata* BALY, 1874).

- E. abdominalis** (BALY, 1874) クロバハラグリハムシ
 BALY, 1874, Trans. ent. Soc. London, **1874**, p. 180 (*Aenidea*) (Nagasaki).
 Distr.: Japan (Honshu, Kyushu).

- E. nigripes** (BALY, 1874) クロアシハラグリハムシ
 BALY, 1874, Trans. ent. Soc. London, **1874**, p. 182 (*Aenidea*) (Nagasaki).
 Distr.: Japan (Kyushu).

- E. ornata** (BALY, 1874) シロモンハラグリハムシ
 BALY, 1874, Trans. ent. Soc. London, **1874**, p. 180 (*Aenidea*) (Nagasaki; China).
abdominalis LABOISSIÈRE, 1929, Ann. Soc. ent. France, **98**, p. 278 (*Liroetis*) (nec
 BALY, 1874) (China).
 Distr.: Japan (Kyushu), E. Siberia, Korea, E. China.

Genus *Cerophysella* LABOISSIÈRE, 1930 カクムネヒゲナガハムシ属

Cerophysella LABOISSIÈRE, 1930, Ann. Soc. ent. France, **99**, p. 352 (type-species: *Cerophysella tonkinensis* LABOISSIÈRE, 1930).

- C. basalis** (BALY, 1874) カクムネヒゲナガハムシ
 BALY, 1874, Trans. ent. Soc. London, **1874**, p. 181 (*Aenidea*) (Satsuma; Shanghai).
plagiata LABOISSIÈRE, 1930, Ann. Soc. ent. France, **99**, p. 354 (China, Tonkin).
 Distr.: Japan (Kyushu), SE China, Hainan, Vietnam.

Genus *Haplosomoides* DUVIVIER, 1890 セスジハムシ属

Haplosomoides DUVIVIER, 1890, C. R. Soc. ent. Belg., **34**, p. 34 (type-species: *Rhaphidopalpa serena* BOHEMAN, 1859).

Hoplasomedia MAULIK, 1936, Fauna India, Galeruc., p. 493.

- H. costata** (BALY, 1878) ホソセスジハムシ
 BALY, 1878, Ann. Mag. nat. Hist., (5), **2**, p. 415 (*Mimastra*) (China).
 Distr.: Ryukyu Is. (Amami-Oshima, Okinawa, Ishigaki, Iriomote, Yonaguni),
 S. China, Taiwan.

H. miyamotoi KIMOTO, 1965 キセスジハムシKIMOTO, 1965, J. Fac. Agr. Kyushu Univ., **13**, p. 371 (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima).

Genus *Taumacera* THUNBERG, 1814 クロバヒゲナガハムシ属*Taumacera* THUNBERG, 1814, Vet. Acad. Handl., p. 48 (type-species: *Taumacera dusta* THUNBERG, 1814).*Cerophyta* BLANCHARD, 1845, Hist. Insect., **2**, p. 190.*Neocharis* JACOBY, 1881, Proc. zool. Soc. London, **1881**, p. 448 (nec SHARP, 1877).*Metellus* JACOBY, 1886, Ann. Mus. Civ. Genova, **24**, p. 63 (n. n. for *Neocharis* JACOBY).*Necrea* BALY, 1886, Trans. ent. Soc. London, **1886**, p. 29.Subgenus *Cerophysa* CHEVROLAT, 1837*Cerophysa* CHEVROLAT, 1837, in DEJEAN, Cat. Col., ed. 3, p. 399 (type-species: *Galleruca nodicornis* WIEDEMANN, 1823).*Ozomena* CHEVROLAT, 1845, in D'ORBIGNY, Dict. Univ. Hist. Nat., **6**, p. 5.**T. (C.) tibialis** (JACOBY, 1885) クロバヒゲナガハムシJACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 750, pl. 46, fig. 8 (*Aenidea*) (Nara, Kobe Maiya-san, Nikko).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan.

Genus *Fleutiauxia* LABOISSIÈRE, 1933 クワハムシ属*Fleutiauxia* LABOISSIÈRE, 1933, Ann. Soc. ent. France, **102**, p. 53 (type-species: *Fleutiauxia cyanipennis* LABOISSIÈRE, 1933).**F. armata** (BALY, 1874) クワハムシBALY, 1874, Trans. ent. Soc. London, **1873**, p. 179 (Nagasaki, Hiogo, Tushima; Manchuria).

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima, Yakushima), Korea, E. Siberia, N. China.

Genus *Liroetis* WEISE, 1889 ルリバナナガハムシ属*Liroetis* WEISE, 1889, Horae Soc. ent. Ross., **23**, p. 507 (type-species: *Liroetis aeneipennis* WEISE, 1889).**L. coeruleipennis** WEISE, 1889 ルリバナナガハムシWEISE, 1889, Horae Soc. ent. Ross., **23**, p. 609 nota (Hagi).

Distr.: Japan (Honshu).

Genus *Cneorane* BALY, 1865 キムネアオハムシ属*Cneorane* BALY, 1865, Ent. monthly Mag., **2**, p. 97 (type-species: *Cneorane fulvicollis* BALY, 1865).**C. elegans** BALY, 1874 キムネアオハムシBALY, 1874, Trans. ent. Soc. London, **1874**, p. 182 (Nagasaki).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Paridea* BALY, 1886 アトボンハムシ属*Paridea* BALY, 1886, J. Linn. Soc. London, **20**, p. 26 (type-species: *Paridea thoracica* BALY, 1886).

Paraulaca BALY, 1888, J. Linn. Soc. London, **20**, p. 168.

Semacia FAIRMAIRE, 1889, Ann. Soc. ent. France, **58**, p. 92.

Aeropa WEISE, 1889, Horae Soc. ent. Ross., **23**, p. 621.

Carapaula CHÔJÔ, 1962, Philip. J. Sci., **91**, p. 198.

P. angulicollis (MOTSCHULSKY, 1853) アトボシハムシ

MOTSCHULSKY, 1853, Étud. ent., **2**, p. 50 (*Rhaphidopalpa*) (China).

nipponensis LABOISSIÈRE, 1930, Ann. Soc. ent. France, **99**, p. 335 (*Semacia*) (Japan).

Distr.: Japan (Hokkaido, Honshu, Hachijo, Sado, Shikoku, Kyushu, Tsushima, Danjo-gunto, Yakushima), E. China, Taiwan.

P. quadriplagiata (BALY, 1874) ヨツボシハムシ

BALY, 1874, Trans. ent. Soc. London, **1874**, p. 186 (*Aulacophora*) (Nagasaki).

verticallis LABOISSIÈRE, 1930, Ann. Soc. ent. France, **99**, p. 343 (Yunnan).

Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima), S. China.

Genus ***Morphosphaera*** BALY, 1861 イチモンジハムシ属

Morphosphaera BALY, 1861, J. Ent., **1** (4), p. 298 (type-species: *Chrysomela japonica* HORNSTEDT, 1788).

M. japonica (HORNSTEDT, 1788) イチモンジハムシ

HORNSTEDT, 1788, Schrift. Ges. Naturf. Freunde Berlin, **2**, p. 1, pl. 1, fig. 1 (*Chrysomela*) (Japan).

japonicum BALY, 1874, Trans. ent. Soc. London, **1874**, p. 176 (*Adorium*) (Nagasaki; China).

Distr.: Japan (Honshu, Shikoku, Kyushu, Tsushima), Ryukyu Is. (Tokara), E. Siberia, China, India.

M. coerulea (SCHÖNFELDT, 1890) オキナワイチモンジハムシ

SCHÖNFELDT, 1890, Ent. Nachr., **16**, (11), p. 173 (*Adorium japonicum* var.) (Amami-Oshima).

Distr.: Ryukyu Is. (Kikaigashima, Amami-Oshima, Okinoerabu, Okinawa, Miyako, Ishigaki, Iriomote).

Genus ***Agelastica*** CHEVROLAT, 1837 ハンノキハムシ属

Agelastica CHEVROLAT, 1837, in DEJEAN, Cat. Col., ed. 3, p. 405 (type-species: *Chrysomela alni* LINNÉ, 1758).

A. coerulea BALY, 1874 ハンノキハムシ

BALY, 1874, Trans. ent. Soc. London, **1874**, p. 188 (Yokohama).

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu), Korea, NE China, E. Siberia, N. America.

Genus ***Luperus*** MÜLLER, 1764 ウスバハムシ属

Luperus MÜLLER, 1764, Fauna Insect. Fridrichsdalina, p. 13 (type-species: *Chrysomela flavipes* LINNÉ, 1767; proposed, Z.N. (S.) 2240).

Lyperus BEDEL, 1892, Col. Bassin Seine, **5**, p. 160.

L. laricis MOTSCHULSKY, 1859 ムナキホソヒゲナガハムシ

MOTSCHULSKY, 1859, Mel. Biol. Acad. Sci. Petersb., **3**, p. 236 (Dauria).

viridipennis laricis: OGLOBLIN, 1936, Fauna USSR, **26** (1), pp. 297, 419, 428 (Siberia).

viridipennis murakamii KIMOTO, 1965, J. Fac. Agr. Kyushu Univ., **13**, p. 379 (Hokkaido).

Distr.: Japan, Siberia, Sakhalin, Kuril Is.

subsp. *laricis* MOTSCHULSKY, 1859, s. str.

MOTSCHULSKY, 1859, Mel. Biol. Acad. Sci. Petersb., **3**, p. 236 (Dauria).

Distr.: Japan (Hokkaido), Siberia, Sakhalin, Kuril Is.

subsp. *supurius* OGLOBLIN, 1936

OGLOBLIN, 1936, Fauna USSR, **26** (1), pp. 297, 416, 429 (as *viridipennis supurius*) ("Japan").

Distr.: Japan (Honshu).

L. moorii BALY, 1874 クロウスバハムシ

BALY, 1874, Trans. ent. Soc. London, **1874**, p. 188 (Yokohama).

Distr.: Japan (Honshu, Kyushu).

Genus *Exosoma* JACOBY, 1903 ヒメハムシ属

Malacosoma CHEVROLAT, 1845, Dist. Univ. d'Hist. Nat., **6**, p. 5 (nec HÜBNER, 1916) (type-species: *Chrysomela lusitanica* LINNÉ, 1769).

Exosoma JACOBY, 1903, Trans. ent. Soc. London, **1903**, p. 25 (n. n. for *Malacosoma* CHEVROLAT).

Malacodera BEDEL, 1905, Abeille, **30**, p. 236.

E. akkoeae (CHÛJÔ, 1954) ハラマダラヒメハムシ

CHÛJÔ, 1954, Trans. Shikoku ent. Soc., **4**, p. 53, fig. 1 (*Calomicrus*) (Yamagata, Fukuoka, Aichi, Osaka, Wakayama, Tokushima, Kagawa, Nagano).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan.

E. amamiense (NAKANE et KIMOTO, 1961) アマミキバラヒメハムシ

NAKANE et KIMOTO, 1961, Kontyû, **29**, p. 19 (*Calomicrus*) (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima, Okinawa, Ishigaki, Iriomote).

E. chujoii (NAKANE, 1958) ニセキバラヒメハムシ

NAKANE, 1958, Sci. Rep. Saikyo Univ., **2** (5), A, p. 309, fig. 20 (*Calomicrus*) (Yakushima, Kagoshima, Nagano).

Distr.: Japan (Honshu, Shikoku, Kyushu, Yakushima), C. China.

E. flaviventre (MOTSCHULSKY, 1860) キバラヒメハムシ

MOTSCHULSKY, 1860, Étud. ent., **9**, p. 26 (*Calomicrus*) (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, E. Siberia, N. China, Taiwan.

Genus *Calomicrus* STEPHENS, 1834 オオルリヒメハムシ属

Calomicrus STEPHENS, 1834, Brit. Ent. Mandb., **4**, p. 293 (type-species: *Crioceris circumfusus* MARSHALL, 1802).

C. cyaneus (JACOBY, 1885) ハラグロヒメハムシ

JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 748 (*Monolepta*) (Nikko, Yuyama, Miyanoshita, Suyama, Subashiri).

Distr.: Japan (Honshu, Shikoku, Kyushu).

C. iniquus (WEISE, 1889) ルリヒラタヒメハムシ

WEISE, 1889, Horae Soc. ent. Ross., **23**, pp. 568, 617 (*Luperus*) (Kansu).

iniquus aconticolus CHÛJÔ, 1959, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., **2**

- (81), p. 10 (Honshu).
Distr.: Japan (Honshu, Shikoku), China, Nepal.
- C. nobyi** CHÛJÔ, 1954 オオルリヒメハムシ
CHÛJÔ, 1954, Trans. Shikoku ent. Soc., 4, p. 55, fig. 2 (Tokushima).
Distr.: Japan (Honshu, Shikoku, Kyushu).
- Genus *Hesperomorpha* OGLOBLIN, 1936 ケブカクロナガハムシ属
Hesperomorpha OGLOBLIN, 1936, Fauna USSR, 26 (1), pp. 298, 375, 429 (type-species: *Luperus hirsutus* JACOBY, 1885).
- H. hirsuta** (JACOBY, 1885) ケブカクロナガハムシ
JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 472, pl. 46, fig. 4 (*Luperus*) (Nikko, Yuyama, Kiga, Hitoyoshi).
Distr.: Japan (Honshu, Shikoku, Kyushu), C. China.
- Genus *Stenoluperus* OGLOBLIN, 1936 ヒゲナガウスバハムシ属
Stenoluperus OGLOBLIN, 1936, Fauna USSR, 26 (1), pp. 247, 408, 419 (type-species: *Luperus potanini* WEISE, 1889).
- S. bicarinatus** (WEISE, 1889) チビウスバハムシ
WEISE, 1889, Horae Soc. ent. Ross., 23, p. 615 nota (*Luperus*) (Japan).
Distr.: Japan (Honshu).
- S. cyaneus** (BALY, 1874) ルリウスバハムシ
BALY, 1874, Trans. ent. Soc. London, 1874, p. 184 (*Arthrotus*) (Higo).
Distr.: Japan (Hokkaido, Honshu, Kyushu).
- S. nipponensis** (LABOISSIÈRE, 1913) ヒゲナガウスバハムシ
LABOISSIÈRE, 1913, Ann. Ass. Nat. Levallois-Perret, 1913, p. 67 nota (*Luperus*) (n. n. for *L. longicornis* JACOBY).
longicornis JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 742, pl. 46, fig. 5 (nec FABRICIUS, 1781) (*Luperus*) (Nikko, Kobe, Yuyama).
jacobyi WEISE, 1924, Col. Cat., in Junk, 78, p. 119 (n. n. for *Luperus longicornis* JACOBY).
Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu), Korea, E. Siberia, China.
- Genus *Medythia* JACOBY, 1887 フタスジヒメハムシ属
Medythia JACOBY, 1887, Notes Leyden Mus., 9, p. 242 (type-species: *Medythia quadrimaculata* JACOBY, 1887).
Paraluperodes OGLOBLIN, 1936, Fauna USSR, 26, (1), pp. 210, 431.
- M. nigrobilineata** (MOTSCHULSKY, 1860) フタスジヒメハムシ
MOTSCHULSKY, 1860, Étud. ent., 9, p. 26 (*Cnecodes*) (Japan).
suturalis ab. *abbreviatus* WEISE, 1922, Tijdschr. ent., 65, p. 81 (*Luperodes*).
suturalis nigrobilineata: OGLOBLIN, 1936, Fauna USSR, 26 (1), pp. 312, 433 (*Paraluperodes*) (Amur, Manchuria, China, Japan).
Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima), Ryukyu Is. (Okinoerabu), Korea, E. Siberia, N. China.
- M. suturalis** (MOTSCHULSKY, 1858) フタスジヒメハムシ
MOTSCHULSKY, 1858, Étud. ent., 7, p. 100 (*Cnecodes*) (Burma).
suturalis ab. *vittatus* WEISE, 1922, Tijdschr. ent., 65, p. 85 (*Luperodes*).

Distr.: Ryukyu Is. (Okinawa, Miyako, Ishigaki), S. China, Taiwan, Hainan, Vietnam, Cambodia, Thailand, Burma, India, Malaya, Philippines, Sumatra, Java, Celebes.

Genus *Atrachya* DEJEAN, 1837 ウリハムシモドキ属

Atrachya DEJEAN, 1837, Cat. Col., ed. 3, p. 401 (type-species: *Galeruca menetriesii* FALDERMANN, 1835).

Luperodes MOTSCHULSKY, 1858, Étud. ent., 7, p. 102.

Iphidea BALY, 1865, Ent. monthly Mag., 2, p. 127.

Dyolania LABOISSIÈRE, 1931, Wien. ent. Ztg., 48, p. 51.

A. flavomaculata (CHÛJÔ, 1935) モンキウリハムシモドキ

CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, 25, p. 206 (*Luperodes*) (Iriomote).

Distr.: Ryukyu Is. (Ishigaki, Iriomote).

A. menetriesii (FALDERMANN, 1835) ウリハムシモドキ

FALDERMANN, 1835, Mem. Acad. St. Petersburg, 2, p. 439, pl. 5, fig. 7 (*Galeruca*) (N. China).

nigripennis MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 232, pl. 11, fig. 18 (*Luperodes*) (Dauria, Amur).

praeustus MOTSCHULSKY, 1860, *ibid.*, p. 232, pl. 11, fig. 19 (*Luperodes*) (Dauria, Amur).

discrepans BALY, 1865, Ent. monthly Mag., 2, p. 127 (*Iphidea*) (Japan).

praeustus var. *insularis* WEISE, 1922, Tijdschr. ent., 65, p. 81 (Japan).

Distr.: Japan (Hokkaido, Honshu, Sado, Hachijo, Shikoku, Kyushu, Tsushima, Yakushima), Korea, E. Siberia, China, Sakhalin.

Genus *Monolepta* CHEVROLAT, 1837 アシナガハムシ属

Monolepta CHEVROLAT, 1837, in DEJEAN, Cat. Col., ed. 3, p. 407 (type-species: *Crioceris bioculata* FABRICIUS, 1781).

Cnecodes MOTSCHULSKY, 1858, Étud. ent., 7, p. 100.

Ochralea CLARK, 1865, Ann. Mag. nat. Hist., (3), 15, p. 144.

Candezea CHAPUIS, 1879, Ann. Mus. Civ. Genova, 15, p. 24.

Alaotra DUVIVIER, 1891, C. R. Soc. ent. Belg., (4), 21, p. 318.

Damais JACOBY, 1908, Ann. Soc. ent. Belg., 48, p. 118.

M. chujoi NAKANE et KIMOTO, 1961 コモンアシナガハムシ

NAKANE et KIMOTO, 1961, Kontyû, 29, p. 20 (Amami-Oshima).

longitarsoides: CHÛJÔ, 1957, Mem. Fac. Lib. Arts & Educ., 2 (5), p. 4 (Amami-Oshima).

M. dichroa HAROLD, 1877 ホタルハムシ

HAROLD, 1877, Dstch. ent. Z., 21, p. 366 (Hakodate).

dichroa var. *apicipennis* JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 749 (Sapporo).

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima).

M. fulvicollis JACOBY, 1885 アオバアシナガハムシ

JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 749 (Kashiwagi).

Distr.: Japan (Honshu).

M. hiurai KIMOTO, 1965 ヒウラアシナガハムシ

KIMOTO, 1965, J. Fac. Agr. Kyushu Univ., 13, p. 392 (Yajima-cho in Akita).

- Distr.: Japan (Honshu).
- M. kurosawai** CHÛJÔ et OHNO, 1961 ムネアカウスイロハムシ
CHÛJÔ et OHNO, 1961, Mem. Fac. Lib. Arts & Educ., **2** (106), p. 10 (Ishikawa, Gifu, Fukushima, Gumma, Saitama, Yamagata, Kochi, Niigata).
Distr.: Japan (Honshu, Shikoku).
- M. minor** CHÛJÔ, 1938 ヒメアシナガハムシ
CHÛJÔ, 1938, Arb. Morph. Taxon. Ent. Berlin-Dahlem, **2** (5), p. 149 (Formosa).
seminigra CHÛJÔ, 1962, Philip. J. Sci., **91**, p. 113 (Formosa).
Distr.: Japan (Yakushima), Ryukyu Is. (Amami-Oshima, Okinawa), Taiwan.
- M. miyamotoi** KIMOTO, 1965 ミヤモトアシナガハムシ
KIMOTO, 1965, J. Fac. Agr. Kyushu Univ., **13**, p. 393 (Amami-Oshima).
Distr.: Ryukyu Is. (Amami-Oshima, Okinoerabu).
- M. nojiriensis** NAKANE, 1963 ヒメウスイロハムシ
NAKANE, 1963, Fragm. Col., Kyoto, (5), p. 2 (Nojiri).
tsushimana KIMOTO, 1965, J. Fac. Agr. Kyushu Univ., **13** p. 393 (Tsushima).
Distr.: Japan (Honshu, Kyushu, Tsushima).
- M. pallidula** BALY, 1874 キイロクワハムシ
BALY, 1874, Trans. ent. Soc. London, **1874**, p. 187 (Nagasaki).
gracilipes CHÛJÔ, 1938, Arb. Morph. Taxon. Ent. Berlin-Dahlem, **2** (5), p. 145 (Formosa).
Distr.: Japan (Honshu, Shikoku, Kyushu, Tanegashima, Yakushima), Ryukyu Is. (Tokara, Okinawa), Korea, S. China, Taiwan.
- M. quadriguttata** (MOTSCHULSKY, 1860) モンキアシナガハムシ
MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 233, pl. 11, fig. 20 (*Luperodes*) (E. Siberia: Dauria, Amur).
hieroglyphica biarcuata: OGLOBLIN, 1936, Fauna USSR, **26** (1), pp. 315, 318, fig. 131 (Japan).
Distr.: Japan (Honshu, Shikoku, Kyushu), E. Siberia, NE China, Korea.
- M. sakishimana** KIMOTO et GRESSITT, 1966 サキシマアシナガハムシ
KIMOTO et GRESSITT, 1966, Pac. Ins., **8**, p. 529 (Iriomote, Ishigaki).
Distr.: Ryukyu Is. (Ishigaki, Iriomote).
- M. shirozui** KIMOTO, 1965 シロウズアシナガハムシ
KIMOTO, 1965, J. Fac. Agr. Kyushu Univ., **13**, p. 390 (Tsushima).
Distr.: Japan (Tsushima).

Genus *Epiluperodes* KIMOTO et GRESSITT, 1966 キイロヒメハムシ属

Epiluperodes KIMOTO et GRESSITT, 1966, Pac. Ins., **8**, p. 531 (type-species: *Epiluperodes ryukyuana* KIMOTO et GRESSITT, 1966).

E. ryukyuana KIMOTO et GRESSITT, 1966 キイロヒメハムシ
KIMOTO et GRESSITT, 1966, Pac. Ins., **8**, p. 531 (Ishigaki, Iriomote).
Distr.: Ryukyu Is. (Ishigaki, Iriomote).

Genus *Theopea* BALY, 1864 キンリョクヒゲナガハムシ属

Theopea BALY, 1864, Trans. ent. Soc. London, (3), **2**, p. 237 (type-species: *Crioceris impressa* FABRICIUS, 1801).

Ozomena ALLARD, 1889, C. R. Soc. ent. Belg., **33**, p. 111.

T. aureoviridis CHÛJÔ, 1935 キンリョクヒゲナガハムシ

CHÛJÔ, 1935, Trans. nat. Hist. Soc. Formosa, **25**, p. 85 (Iriomote).

Distr.: Ryukyu Is. (Kikaigashima, Okinawa, Ishigaki, Iriomote).

Genus *Hoplosaenidea* LABOISSIÈRE, 1933 オキナワヒメナガハムシ属

Hoplosaenidea LABOISSIÈRE, 1933, Ann. Soc. ent. France, **102**, p. 59 (type-species: *Hoplosaenidea touzalini* LABOISSIÈRE, 1933).

Diaphaenidea LABOISSIÈRE, 1933, *ibid.*, p. 61.

Micraenidea LABOISSIÈRE, 1933, *ibid.*, p. 64.

H. miyatakei KIMOTO et GRESSITT, 1966 オキナワヒメナガハムシ

KIMOTO et GRESSITT, 1966, Pac. Ins., **8**, p. 533 (Ishigaki, Iriomote).

Distr.: Ryukyu Is. (Ishigaki, Iriomote).

Genus *Epaenidea* GRESSITT et KIMOTO, 1963 アオバヒメハムシ属

Epaenidea GRESSITT et KIMOTO, 1963, Pac. Ins. Mon., **1B**, p. 684 (type-species: *Epaenidea subvirida* GRESSITT et KIMOTO, 1963).

E. elegans KIMOTO et GRESSITT, 1966 アオバヒメハムシ

KIMOTO et GRESSITT, 1966, Pac. Ins., **8**, p. 535 (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima).

Genus *Dercetina* GRESSITT et KIMOTO, 1963 アオバヒゲナガハムシ属

Dercetis CLARK, 1865, Ann. Mag. nat. Hist., (3), **15**, p. 146 (nec MÜNSTER et AGASSIZ, 1834) (type-species: *Dercetis depressa* CLARK, 1865).

Antipha BALY, 1865, Ann. Mag. nat. Hist., (3), **16**, p. 251 (nec WALKER, 1855).

Dercetina GRESSITT et KIMOTO, 1963, Pac. Ins. Mon., **1B**, p. 704 (n. n. for *Dercetis* CLARK, 1865).

D. azumai KIMOTO et GRESSITT, 1966 アオバヒゲナガハムシ

KIMOTO et GRESSITT, 1966, Pac. Ins., **8** p. 534 (Iriomote).

Distr.: Ryukyu Is. (Iriomote).

Genus *Arthrotus* MOTSCHULSKY, 1857 ムナグロツヤハムシ属

Arthrotus MOTSCHULSKY, 1857, Étud. ent., **6**, p. 38. (type-species: *Arthrotus niger* MOTSCHULSKY, 1857).

Taphinella JACOBY, 1889, Ann. Mus. Civ. Genova, **27**, p. 224.

Dercestra CHÛJÔ, 1962, Philip. J. Sci., **91**, p. 163.

A. niger MOTSCHULSKY, 1857 ムナグロツヤハムシ

MOTSCHULSKY, 1857, Étud. ent., **6**, p. 39 (Japan).

variabilis BALY, 1874, Trans. ent. Soc. London, **1874**, p. 183 (Nagasaki, Yokohama, Hiogo).

suzukii MATSUMURA, 1934, Ins. Matsumurana, **9**, p. 63 (*Luperodes*) (Kioto, Binomori).

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu).

Genus *Hamushia* CHÛJÔ, 1956 ヨツキボンハムシ属

Hamushia CHÛJÔ, 1959, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., **2** (31), p. 9 (type-

species: *Galerucida eburata* HAROLD, 1879).

H. eburata (HAROLD, 1879) ヨツキボシハムシ

HAROLD, 1879, Dtsch. ent. Z., **23** (2), p. 368 (*Galerucida*) (Hakone).

Distr.: Japan (Honshu).

H. konishii CHÛJÔ, 1956 ヨモギモンキハムシ

CHÛJÔ, 1956, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., **2** (31), p. 11 (Honshu).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Sphenoraia* CLARK, 1865 アオバノコヒゲハムシ属

Sphenoraia CLARK, 1865, Ann. Mag. nat. Hist., (3), **16**, p. 262 (type-species: *Galleruca bicolor* HOPE, 1831).

Sphenoraoides LABOISSIÈRE, 1934, Ann. Assoc. Nat. Levallois-Perret, **21**, p. 131.

Neosermylassa CHÛJÔ, 1956, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., **2** (31), p. 14.

S. intermedia JACOBY, 1885 アオバノコヒゲハムシ

JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 747 (Noheji).

japonica CHÛJÔ, 1957, Mem. Fac. Lib. Arts & Educ., **2** (31), p. 15, fig. 7 (*Sermylassa*) (Kyoto, Gumma, Yamagata).

Distr.: Japan (Honshu).

Genus *Agelasa* MOTSCHULSKY, 1860 キクビアオハムシ属

Agelasa MOTSCHULSKY, 1860, Étud. ent., **9**, p. 20 (type-species: *Agelasa nigriceps* MOTSCHULSKY, 1860).

A. nigriceps MOTSCHULSKY, 1860 キクビアオハムシ

MOTSCHULSKY, 1860, Étud. ent., **9**, p. 25 (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yakushima), Korea, NE China, Siberia.

Genus *Gallerucida* MOTSCHULSKY, 1860 イタドリハムシ属

Gallerucida MOTSCHULSKY, 1860, Étud. ent., **9**, p. 25 (type-species: *Gallerucida bifasciata* MOTSCHULSKY, 1860).

Eustetha BALY, 1861, J. Ent. **1**, p. 296.

Melosphila BALY, 1861, *ibid.*, p. 297.

Coptomesa WEISE, 1912, Arch. Naturg., (A) **78** (2), p. 91.

G. bifasciata MOTSCHULSKY, 1860 イタドリハムシ

MOTSCHULSKY, 1860, Étud. ent., **9**, p. 24 (Japan).

nigromaculata BALY, 1861, J. Ent., **1**, p. 297 (*Melospila*) (N. China).

consociata BALY, 1874, Trans. ent. Soc. London, **1874**, p. 185 (*Melospila*) (Hokkaido)

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima), E. Siberia, China, Korea, Taiwan.

G. flavipennis (SOLSKY, 1872) ズグロアカハムシ

SOLSKY, 1872, Horae Soc. ent. Ross., **8**, p. 257 (*Galerucida*) (E. Siberia).

melanocephala JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 748 (*Sphenoraia*) (Yuyama).

Distr.: Japan (Honshu, Shikoku, Kyushu), E. Siberia.

G. lewisi (JACOBY, 1885) ムツキボシハムシ

JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 747, pl. 46, fig. 6 (*Galerucida*)

(Oyama).

Distr.: Japan (Honshu, Shikoku).

G. oshimana KIMOTO et GRESSITT, 1966 アマミカバイロハムシ

KIMOTO et GRESSITT, 1966, Pac. Ins., 8, p. 536 (Amami-Oshima).

Distr.: Ryukyu Is. (Amami-Oshima).