

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 29

April 30, 1985

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.),
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family PLATYPODIDAE ナガキクイムシ科

By

Akira NOBUCHI

Subfamily DIAPORINAE トゲナガキクイムシ亜科

Genus *Diapus* CHAPUIS, 1865 トゲナガキクイムシ属

Diapus CHAPUIS, 1865, Monographie des Platypides, p. 329 (type-species: *D. quadri-spinatus* CHAPUIS, 1865).

D. aculeatus BLANDFORD, 1894 トゲナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 139, ♀ (lege ♂) (Higo).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan, India, Java.

Subfamily PLATYPODINAE ナガキクイムシ亜科

Genus *Crossotarsus* CHAPUIS, 1865 オオナガキクイムシ属

Crossotarsus CHAPUIS, 1865, Monographie des Platypides, p. 44 (type-species: *Platypus wallacei* THOMSON, 1858).

C. concinnus BLANDFORD, 1894 ヒゴナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 578, (Higo).

chapuisi BLANDFORD, 1894, *ibid.*, p. 129, ♀ (lege ♂) (nec DUVIVIER, 1891).

Distr.: Japan (Kyushu).

C. emancipatus MURAYAMA, 1934 マルオナガキクイムシ

MURAYAMA, 1934, J. Coll. Agr., Hokkaido imp. Univ., 35, p. 138, ♂, ♀. (Nagao).

terminatus MURAYAMA, 1928, *ibid.*, 19, p. 288 (nec CHAPUIS, 1865).

Distr.: Japan (Kyushu, Yakushima), Ryukyu Is. (Amami-Ôshima), Taiwan.

C. externedentatus (FAIRMAIRE, 1850) ソトハナガキクイムシ

FAIRMAIRE, 1850, Rev. Mag. Zool. (2), 2, p. 51, ♂ (*Platypus*) (Taiti).

saundersi CHAPUIS, 1865, Monographie des Platypides, p. 80, fig. 19, ♂, ♀ (lege ♀, ♂) (Celebes & Borneo).

Distr.: Japan (Kyushu), Ryukyu Is. (Amami-Ōshima, Okinawa, Ishigaki, Iriomote), Indochina, Caroline Is., Hawaii, Samoa, Fiji, Society Is., Madagascar.

C. fairmairei CHAPUIS, 1865, フェアメルナガキクイムシ

CHAPUIS, 1865, Monographie des Platypides, p. 79, fig. 18, ♀ (lege ♂) (India).

Distr.: Japan (Honshu), India.

C. flavomaculatus STROHMEYER, 1912 キバネナガキクイムシ

STROHMEYER, 1912, Ent. Mitt., **1**, p. 40, ♂, ♀ (Hoozan in Formosa, Philippines).

Distr.: Japan (Kyushu), Taiwan, Philippines, Malaya, Java.

C. niponicus BLANDFORD, 1894 ヤチダモノナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 130, ♂, ♀ (lege ♀, ♂) (Sapporo, Hakodate, Miyanoshta, Yuyama, Kyushu).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Ryukyu Is. (Amami-Ōshima), Taiwan.

C. simplex MURAYAMA, 1925 カシノコナガキクイムシ

MURAYAMA, 1925, J. Coll. Agr., Hokkaido imp. Univ., **15**, p. 231, ♂, ♀ (Ayakita).

Distr.: Japan (Honshu, Shikoku, Kyushu, Yakushima), Ryukyu Is. (Okinawa, Iriomote), Korea, Taiwan.

Genus *Platypus* HERBST, 1793 ナガキクイムシ属

Platypus HERBST, 1793, Natursystem aller bekannten in- und ausländischen Insekten, **5**, p. 128 (type-species: *Bostrichus cylindrus* FABRICIUS, 1792).

P. calamus BLANDFORD, 1894 ヨシブエナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 137, ♀ (lege ♂) (Miyanoshta, Oshima, Kiushu (Higo, Yuyama, etc.)).

Distr.: Japan (Hokkaido, Honshu, Miyake-jima, Shikoku, Kyushu, Yakushima), Ryukyu Is. (Amami-Ōshima, Okinawa, Ishigaki, Iriomote), Korea, Taiwan.

P. contaminatus (BLANDFORD, 1894) タブノナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 131, ♀ (*Crossotarsus*) (Higo).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan, Himalaya, China (Fukien).

P. hamatus BLANDFORD, 1894 カギナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 138, ♀ (lege ♂) (Yuyama, Miyanoshta).

tenuis MURAYAMA, 1925, J. Coll. Agr., Hokkaido imp. Univ., **15**, p. 234, ♀ (Sapporo).

Distr.: Japan (Hokkaido, Honshu, Kyushu).

P. kiushuensis MURAYAMA, 1936 キュウシュウナガキクイムシ

MURAYAMA, 1936, Tenthredo, **1**, pp. 139, 149, ♀ (Ishikochi, Nagao).

Distr.: Japan (Kyushu), Taiwan.

P. lewisi BLANDFORD, 1894 ルイスナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 134, ♂, ♀ (lege ♀, ♂) (Miyanoshta, Kiga, Yuyama).

uncacanthurus BEESON, 1941, Forest Insects of India, p. 347 (India).

Distr.: Japan (Hokkaido, Honshu, Kyushu), Querpart Is., Taiwan, China (Fukien), India.

P. modestus BLANDFORD, 1894 チュウガタナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 133, ♂, ♀ (lege ♀, ♂) (Nikko, Shimidzu Toge).

Distr.: Japan (Honshu, Kyushu), Taiwan.

P. quercivorus (MURAYAMA), 1925 カシノナガキクイムシ

MURAYAMA, 1925, J. Coll. Agr., Hokkaido imp. Univ., 15, p. 229, ♂, ♀ (*Crossotarsus*) (Echigo, Ayakita).

sexfenestratus BEESON, 1937, Ind. For. Rec., N.S., (3) 3, p. 94, ♂, ♀ (*Crossotarsus*).

Distr.: Japan (Honshu, Miyake-jima, Shikoku, Kyushu), Ryukyu Is. (Amami-Ōshima, Okinawa), Taiwan, India, Java, New Guinea.

P. severini BLANDFORD, 1894 シナノナガキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 136, ♀ (lege ♂) (Nikko, Chuzenji, Hakodate).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

P. solidus WALKER, 1858 トガリハネナガキクイムシ

WALKER, 1858, Ann. Mag. nat. Hist., (3), 2, p. 286, ♂ (Ceylon).

Distr.: Japan (Honshu, Shikoku), Ryukyu Is. (Okinawa), Korea, Taiwan, Philippines, Borneo, Celebes, Tonkin, Malaya, Sumatra, Java, Caroline Is., Mariana Is., Guam, India, Ceylon, Australia, New Guinea, Aroe, Solomon Is.

P. taiwansis SCHEDL, 1960 タイワンアカキクイムシ

SCHEDL, 1960, Ent. B1., 56, p. 111.

formosanus NIJIMA et MURAYAMA, 1925, J. Coll. Agr., Hokkaido imp. Univ., 15, p. 215, ♂, ♀ (Rengeti, Formosa) (nec STROHMEYER, 1912).

Distr.: Japan (Kyushu), Ryukyu Is. (Tokara Is., Amami-Ōshima, Okinawa), Taiwan.