

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 30

April 30, 1985

The Coleopterists' Association of Japan
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family SCOLYTIDAE キクイムシ科*

By

Akira NOBUCHI

Subfamily SCOLYTINAE ゾウキカワノキクイムシ亞科

Genus *Scolytus* GEOFFROY, 1762 ゾウキカワノキクイムシ属

Scolytus GEOFFROY, 1762, Historie abrégée insects, 1, p. 305 (type-species: *Bostrihus scolytus* FABRICIUS, 1775).

Ekkoptogaster HERBST, 1793, Natursystem aller bekannten in- und ausländischen Insekten, 5, p. 124 (type-species: *Bostrihus scolytus* FABRICIUS, 1775).

S. aomoriensis NOBUCHI, 1973 アオモリキクイムシ

NOBUCHI, 1973, Bull. Gov. For. Exp. Sta., (258), p. 16 (Aburakawa).

Distr.: Japan (Honshu).

S. aratus BLANDFORD, 1894, ウメノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 79. pl. 1, fig. 1 (Junsai).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Kuril Is., Siberia, Korea.

S. chikisanii NIIJIMA, 1905 ニレノキクイムシ

NIIJIMA, 1905, J. Sapporo Agr. Coll., 2, p. 69, pl. 2, figs. 4, 5 (Sapporo, Mt. Moiwa).

curviventralis NIIJIMA, 1905, ibid., p. 70, pl. 2, figs. 6, 7 (Mt. Moiwa).

betulae NIIJIMA, 1943, Trans. Sapporo nat. Hist. Soc., 17, p. 141 (Pompira).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Kuril Is., Siberia, China.

S. claviger BLANDFORD, 1894 サワシバノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 80 (Kiga).

Distr.: Japan (Hokkaido, Honshu, Kyushu), Siberia, Korea, NE. China.

* Studies on Scolytidae XXVI

- S. dahuricus* CHAPUIS, 1873 シラカバノキクイムシ (ダフリアキクイムシ)
 CHAPUIS, 1873, Mém. Soc. roy. Sci. Liège, 2 (3), p. 268 (Daouria).
agnatus BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 78 (Junsai).
 Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, NE. China.
- S. ellipticus* MURAYAMA, 1958 アサマヤマキクイムシ
 MURAYAMA, 1958, Bull. Fac. Agr. Yamaguti Univ., 9, p. 929 (Onino-oshidashi).
 Distr.: Japan (Hokkaido, Honshu).
- S. esuriens* BLANDFORD, 1894 ニレノオオキクイムシ (シラカバノキクイムシ)
 BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 77 (Junsai, Miyanoshita, Chuzenji).
 Distr.: Japan (Hokkaido, Honshu), Kuril Is., Siberia.
- S. frontalis* BLANDFORD, 1894 ニレカワノキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 79 (Fukushima).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan.
- S. jacobsoni* SPESSIVTZEV, 1919 ジャコブソンキクイムシ
 SPESSIVTZEV, 1919, Ent. month. Mag., 55, p. 246 (Vladivostok).
 Distr.: Japan(?), Sakhalin, Siberia.
- S. japonicus* CHAPUIS, 1875 ニホンキクイムシ
 CHAPUIS, 1875, Ann. Soc. ent. Belg., 18, p. 199 (Kyushu).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Siberia, Korea, China, Mongolia, Taiwan.
- S. ratzeburgi* JANSON, 1856 ラツェブルグキクイムシ
 JANSON, 1856, Ent. Ann., 1856, p. 87 (Rannoch, Perthshire, England).
amurensis EGGLERS, 1908, Wien. ent. Ztg., 27, p. 144 (*Eccoptogaster*) (Siberia orientalis, Amur sup.).
sibiricus EGGLERS, 1922, Ent. B1., 18, p. 14 (*Eccoptogaster*). (Werkhne-Udinsk).
 Distr.: Japan (Hokkaido), Sakhalin, Siberia, Korea, China, Europe.
- S. trispinosus* STROHMEYER, 1908 ミツハリキクイムシ
 STROHMEYER, 1908, Ent. Wochentbl., 1908, p. 69.
 Distr.: Japan (Hokkaido), Kuril Is., Sakhalin, Siberia.
- S. ussuriensis* KURENZOV, 1941 ウスリーキクイムシ
 KURENZOV, 1941, Koroedy dal'nego vostoka SSSR, pp. 102, 226 (River Suputinka).
 Distr.: Japan (Honshu), Siberia.
- Subfamily HYLESININAE ヒレジニ亞科
 Tribe SPHAEROTRYPINI マルキクイムシ族
 Genus *Sphaerotrypes* BLANDFORD, 1894 マルキクイムシ属
- Sphaerotrypes* BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 61 (type-species:
Sphaerotrypes pila BLANDFORD, 1894).
- S. carpini* EGGLERS, 1926 シデノマルキクイムシ
 EGGLERS, 1926, Ent. B1., 22, p. 134 (Sapporo, Maruyama).
 Distr.: Japan (Hokkaido, Honshu).
- S. imitans* EGGLERS, 1926 タカオマルキクイムシ
 EGGLERS, 1926, Ent. B1., 22, p. 134 (Takao Berg).

Distr.: Japan (Honshu), China.

S. pila BLANDFORD, 1894 ケブカマルキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 62 (Hitoyoshi).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Taiwan.

Tribe HYORRHYNCHINI オオキクイムシ族

Genus *Hyorrhynchus* BLANDFORD, 1894 オオキクイムシ属

Hyorrhynchus BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 60 (type-species:
Hyorrhynchus lewisi BLANDFORD, 1894).

H. lewisi BLANDFORD, 1894, ルイスオオキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London 1894, p. 60, ♂, ♀ (Sapporo).
 Distr.: Japan (Hokkaido, Honshu, Shikoku), Taiwan.

H. unicornis NOBUCHI, 1966 ツノオオキクイムシ
 NOBUCHI, 1966, Bull. Gov. For. Exp. Sta., (185), p. 51, ♂ (Gamushi).
 Distr.: Japan (Hokkaido).

Genus *Pseudohyorrhynchus* MURAYAMA, 1950 ニセオオキクイムシ属
Pseudohyorrhynchus MURAYAMA, 1950, Ins. Matsum., 17, p. 61 (type-species: *Pseudohyorrhynchus wadai* MURAYAMA, 1950).

P. wadai MURAYAMA, 1950 ワダオオキクイムシ
 MURAYAMA, 1950, Ins. Matsum., 17, p. 61 (Daidominamiyama).
 Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Sueus* MURAYAMA, 1951 マルキクイムシモドキ属
Sueus MURAYAMA, 1951, Bull. Fac. Agr. Yamaguti Univ., 2, p. 1 (type-species: *Sueus sphaerotropoides* MURAYAMA, 1951).
Neohyorrhynchus SCHEDL, 1962, Ent. B1., 58, p. 202 (type-species: *Hyorrhynchus niisimai* EGGERS, 1926).

S. niisimai (EGGERS, 1926) ニイシマキクイムシ
 EGGERS, 1926, Ent. B1., 22, p. 133, ♀ (*Hyorrhynchus*) (Urakawa).
pilosus EGGERS, 1936, Tijd. Ent., 79, p. 81, ♀ (*Hyorrhynchus*) (Java).
sphaerotropoides MURAYAMA, 1951, Bull. Fac. Agr. Yamaguti Univ., 2, p. 2 (Sue).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yakushima), Ryukyu Is. (Iriomote), Malaya, Java, Ceylon.

Genus *Parasphaerotypes* MURAYAMA, 1954 ニセマルキクイムシ属
Parasphaerotypes MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 9, p. 933 (type-species: *Sphaerotypes controversae* MURAYAMA, 1950).

P. controversae (MURAYAMA, 1950) ミズキマルキクイムシ
 MURAYAMA, 1950, Ins. Matsum., 17, p. 62 (*Sphaerotypes*) (Irazuyama).
 Distr.: Japan (Shikoku).

Tribe HYLASTINI ヒラステイ族

Genus *Hylastes* ERICHSON, 1836 クロキクイムシ属*Hylastes* ERICHSON, 1836, Arch. Naturg., 2, p. 47 (type-species: *Bostrichus ater* PAYKULL, 1800).*H. ambiguus* BLANDFORD, 1894 フジキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 57 (Fujisan).

Distr.: Japan (Honshu, Kyushu).

H. ater (PAYKULL, 1800) マツノクロキクイムシPAYKULL, 1800, Fauna Suecia, 3, p. 153 (*Bostrichus*) (Europe).

Distr.: Japan (Honshu), Siberia, Korea, China, Europe, New Zealand.

H. attenuatus ERICHSON, 1836 マツノホソキクイムシ

ERICHSON, 1836, Arch. Naturg., 2, p. 50.

Distr.: Japan (Honshu), Siberia, Korea, China, Europe.

H. cunicularius ERICHSON, 1836 クロナガキクイムシ

ERICHSON, 1836, Arch. Naturg., 2, p. 49.

Distr.: Japan (Hokkaido, Honshu), Sakhalin, China, Europe.

H. parallelus CHAPUIS, 1875 マツノホソスジキクイムシ

CHAPUIS, 1875, Ann. Soc. ent. Belg., 18, p. 196 (Nipon, Kiushiu).

Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Taiwan, China.

H. plumbeus BLANDFORD, 1894 マツノヒロスジキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 57.

obscurus CHAPUIS, 1875, Ann. Soc. ent. Belg., 18, p. 197 (nec MARSHAM, 1803) (Nagasaki, Hiogo).

Distr.: Japan (Honshu, Shikoku, Kyushu), Siberia, Korea, Taiwan, China, Europe.

Genus *Hylurgops* LECONTE, 1876 カバイロキクイムシ属*Hylurgops* LECONTE, 1876, Proc. Amer. Phil. Soc., 15, p. 389 (type-species: *Hylastes pinifex* FITCH, 1858).*H. glabratus* (ZETTERSTEDT, 1828) マツノカバイロキクイムシ (エゾチャイロキクイムシ)

ZETTERSTEDT, 1828, Fauna Insectorum Lapponica, p. 343 (Europe).

Distr.: Japan (Hokkaido, Honshu, Kyushu), Sakhalin, Siberia, Korea, Taiwan, China, Europe.

H. interstitialis (CHAPUIS, 1875) マツノスジキクイムシCHAPUIS, 1875, Ann. Soc. ent. Belg., 18, p. 196 (*Hylastes*) (Nagasaki, Kiushiu).*imitator* REITTER, 1900, Dtsch. ent. Z., 1900, p. 59 (Chabarowska).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Kamchatka, Siberia, Korea, Taiwan, China.

H. longipilus REITTER, 1894 ケナガエゾキクイムシ

REITTER 1894, Best.-tab. Borkenkäfer, p. 63 (Chabarowka).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, China.

H. niponicus MURAYAMA, 1936 ニホンカバイロキクイムシ

MURAYAMA, 1936, Tenthredo, 1, pp. 123, 142 (Kamikochi, Nagano).

Distr.: Japan (Honshu).

H. palliatus (GYLLENHAL, 1813) ウスイロキクイムシ

GYLENHAL, 1813, Insecta svec., 3, p. 340 (*Hylesinus*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, China, Europe.

H. spessiwzeffi (EGGERS, 1914) スペッシフツエフキイムシ

EGGERS, 1914, Ent. B1., 10, p. 187 (*Hylastes*) (Siberia).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, Taiwan, NE. China.

H. transbaicalicus EGGERS, 1941 トランスバイカルキイムシ

EGGERS, 1941, Stettin. ent. Ztg., 102, p. 199 (Wladiwostok, Tomakomai).

Distr.: Japan (Hokkaido, Honshu), Siberia.

Tribe HYLURGINI ヒルギ族

Genus *Pseudohylesinus* SWAINE, 1917 ニセヒレジスス属

Pseudohylesinus SWAINE, 1917, Dom. Canada Dept. Agric. Ent. Br., Tech. Bull., 14, p.

11 (type-species: *Pseudohylesinus grandis* SWAINE, 1917 = *Hylurgus sericeus* MANNERHEIM, 1843).

P. sericeus MANNERHEIM, 1843 ヤスマツキイムシ

MANNERHEIM, 1843, Bull. Soc. imp. Nat. Moscou, 16, p. 296 (Sitka, Alaska).

yasumatsui NOBUCHI, 1971, Bull. Gov. For. Expt. Sta., (236). p. 160 (Takanishi, Nagano).

Distr.: Japan (Honshu), Alaska, Canada, U.S.A.

Genus *Tomicus* LATREILLE, 1802 マツノキイムシ属

Tomicus LATREILLE, 1802, Hist. nat. générale et particulière des Crustacés et des Insectes, 3, p. 203 (type-species: *Dermestes piniperda* LINNÉ, 1758).

Blastophagus EICHHOFF, 1864, Berl. ent. Z., 8, p. 25 (nec GRAVENHORST, 1827) (type-species: *Dermestes piniperda* LINNÉ, 1758).

Myelophilus EICHHOFF, 1870, Stettin. ent. Ztg., 40, p. 400 (nec TREITSCHKE, 1855) (type-species: *Dermestes piniperda* LINNÉ, 1758).

T. brevipilosus (EGGERS, 1929) ケミジカキイムシ

EGGERS, 1929, Ent. B1., 25, p. 103 (*Blastophagus*) (China).

Distr.: Japan (Honshu), Korea, China.

T. minor (HARTIG, 1834) マツノコキイムシ

HARTIG, 1834, Forstliches und forst-naturwissenschaftliches Conversations-Lexikon, p. 413 (*Dendroctonus*) (Europe).

Distr.: Japan (Honshu, Shikoku, Kyushu), Sakhalin, Siberia, Korea, Taiwan, China, Europe, North Africa.

T. multisetosus (MURAYAMA, 1963)

MURAYAMA, 1963, Studies in the scolytid-fauna of the northern half of the Far East, 5, p. 37 (*Blastophagus*) (Mt. Manza, Gunma Pref.).

Distr.: Japan (Honshu).

T. piniperda (LINNÉ, 1758) マツノキイムシ

LINNÉ, 1758, Sys. Nat., ed. 10, p. 563 (*Dermestes*) (Europe).

major EGGERS, 1943, Ent. B1., 39, p. 50 (*Blastophagus*) (Kobe, Nord Korea).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Sakhalin, Siberia, Korea, Taiwan, China, Europe, North Africa.

T. puellus (REITTER, 1894) キタマツノキイムシ

REITTER, 1894, Best.-tab. Borkenkäfer p. 53 (*Myelophilus*) (Permskoe, Siberia).
Distr.: Japan (Hokkaido), Siberia.

Genus *Hylurgus* LATREILLE, 1807 ネノキクイムシ属

Hylurgus LATREILLE, 1807, Genera Crustaceorum et Insectorum secundum ordinem naturalem in familias disposita, 2, p. 274 (type-species: *Bostrichus ligniperda* FABRICIUS, 1792).

H. ligniperda (FABRICIUS, 1792) マツノネノキクイムシ

FABRICIUS, 1792, Ent. syst. 1, p. 367 (*Hylesinus*) (Europe).
Distr.: Japan (Honshu), Europe, North Africa.

Genus *Dendroctonus* Erichson マツオオキクイムシ属

Dendroctonus ERICHSON, 1836, Arch. Naturg., 2(1), p. 52 (type-species: *Bostrichus micans* KUGELMANN, 1794)

D. micans (KUGELMANN, 1794) エゾマツオオキクイムシ

KUGELMANN, 1794, Schneider Mag., 5, p. 523 (*Bostrichus*) (Europe)
Distr.: Japan (Hokkaido), Sakhalin, Siberia, China, Europe.

Tribe HYLESININI ヒレジニ族

Genus *Ficiphagus* MURAYAMA, 1958 イチジクキクイムシ属

Ficiphagus MURAYAMA, 1958, Bull. Fac. Agr. Yamaguti Univ. 10, p. 930 (type-species: *Phloeosinus goliatoides* MURAYAMA, 1955).

F. goliatoides MURAYAMA, 1955 ゴリアテキクイムシ

MURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., 6, p. 64 (*Phloeosinus*) (Kôyama, Kagoshima).
Distr.: Japan (Honshu, Kyushu), Taiwan.

Genus *Hylesinus* FABRICIUS, 1801 ヒレジヌス属

Hylesinus FABRICIUS, 1801, Syst. Eleuth. 2, p. 390 (type-species: *Hylesinus crenatus* FABRICIUS, 1787).

H. cingulatus BLANDFORD, 1894 シラオビキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 67 (Junsai).
Distr.: Japan (Hokkaido, Honshu, Kyushu), Siberia.

H. elatus NIIZIMA, 1913 タカスジキクイムシ

NIIZIMA, 1913, Trans. Sapporo nat. Hist. Soc., 5, p. 2 (Nishino, Hokkaido).
Distr.: Japan (Hokkaido, Honshu).

H. eos SPESSEVTEFF, 1919 シオジノキクイムシ

SPESSEVTEFF, 1919, Ent. month. Mag., 60, p. 248 (Vicinity of Vladivostok, Manchuria).
Distr.: Japan (Honshu), Siberia, China.

H. laticollis BLANDFORD, 1894, ヤチダモノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 65 (Sapporo).
Distr.: Japan (Hokkaido, Honshu), Siberia.

H. nobilis BLANDFORD, 1894 ヤチダモノオオキクイムシ

BLANDFORD, 1984, Trans. ent. Soc. London, **1894**, p. 64 (Sapporo).
 Distr.: Japan (Hokkaido, Honshu, Kyushu), Siberia.

- H. porcatus* CHAPUIS, 1873 ゴリアテキクイムシモドキ
 CHAPUIS, 1873, Synopsis des Scolytides, p. 239 (Nouvelle-Hollande).
 Distr.: Ryukyu Is., Taiwan, Philippines, Borneo, Sumatra, Java, Oceania, Australia.
- H. tiliae* (NIIJIMA, 1910) シナノキクイムシ
 NIIJIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 2 (*Kissophagus*) (Sapporo).
 Distr.: Japan (Hokkaido).
- H. tristis* BLANDFORD, 1894 ヤチダモノクロキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 66 (Ichiuchi, Junsai).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Ryukyu Is., Korea, Taiwan.

Genus *Neopteleobius* NOBUCHI, 1971 マダラキクイムシ属

Neopteleobius NOBUCHI, 1971, Bull. Gov. For. Exp. Sta., (236), p. 125 (type-species:
Hylesinus scutulatus BLANDFORD, 1894).

- N. scutulatus* (BLANDFORD, 1894) ハルニレノキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 64 (*Hylesinus*) (Kiga, Subashiri,
 Nagasaki, Omori, Oyama).
trepanatus WICHMANN, 1914, Ent. Bl., 10, p. 134 (Kioto, Japan).
 Distr.: Japan (Honshu, Kyushu), Korea.

Genus *Pruniphagus* MURAYAMA, 1954 サクラノキクイムシ属

Pruniphagus MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 9, p. 930 (type-species:
Pruniphagus gummensis MURAYAMA, 1954).

- P. gummensis* MURAYAMA, 1954 グンマキクイムシ
 MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 9, p. 931 (Mt. Manza, Gunma
 Pref.).
 Distr.: Japan (Honshu).

Genus *Alniphagus* SWAINE, 1918 ハンノカバイロキクイムシ属

Alniphagus SWAINE, 1918, Dom. Canada Dept. Agr. Ent. Br., Tech. Bull., **14**, p. 73 (type-
 species: *Hylesinus aspericollis* LECONTE, 1876).

- A. costatus* (BLANDFORD, 1894) ハンノカバイロキクイムシ (ヤチダモノコキクイムシ)
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 63 (*Hylesinus*) (Junsai).
alni (NIIJIMA, 1909), J. Coll. Agr., Tohoku imp. Univ., 3, p. 137 (*Hylastes*) (Toma-
 komai, Jozankei).
 Distr.: Japan (Hokkaido, Honshu), Kuril Is., Sakhalin, Siberia.

Tribe PHLOEOTRIBINI クシヒゲキクイムシ族

Genus *Phthorophloeus* REY, 1883 クシヒゲキクイムシ属

Phthorophloeus REY, 1883, Rév. Ent., 2, p. 127 (type-species: *Phthorophloeus spinulosus*
 REY, 1883).

- P. spinulosus* REY, 1883 アラゲキクイムシ
 REY, 1883, Rév. Ent., 2, p. 127
 Distr.: Japan (Hokkaido), Sakhalin, Siberia, Europe.

Tribe PHLOEOSININI ヒバノキクイムシ族

Genus *Phloeosinus* CHAPUIS, 1873 ヒバノキクイムシ属*Phloeosinus* CHAPUIS, 1873, Synopsis des Scolytides, p. 245 (type-species: *Olonthogaster nitidicollis* MOTSCHULSKY 1866).*P. gifuensis* MURAYAMA, 1954 ギフキクイムシMURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 5, p. 190 (Naka town, Gifu).
Distr.: Japan (Honshu, Shikoku).*P. izuensis* NOBUCHI, 1959 イズカワキクイムシNOBUCHI, 1959, Akitu, 8, p. 9 (Mt. Amagi).
Distr.: Japan (Honshu).*P. kiushuensis* MURAYAMA, 1955 キュウシュウキクイムシMURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., 6, p. 93 (Kôyama).
Distr.: Japan (Kyushu).*P. kumamotoensis* MURAYAMA, 1955 クマモトキクイムシMURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., 6, p. 91 (Waifu).
Distr.: Japan (Kyushu).*P. lewisi* CHAPUIS, 1875 ヒバノコキクイムシ (ヒバノヒメキクイムシ)CHAPUIS, 1875, Ann. Soc. ent. Belg., 18, p. 198 (without locality).
minutus BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 71 (Ichiuchi).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Taiwan.*P. osumiensis* MURAYAMA, 1955 オオスミキクイムシMURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., 6, p. 90 (Kishira).
Distr.: Japan (Kyushu).*P. perlatus* CHAPUIS, 1875 ヒバノキクイムシCHAPUIS, 1875, Ann. Soc. ent. Belg., 18, p. 198 (Hiogo).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Taiwan, China.*P. pulchellus* BLANDFORD, 1894 ユメノキクイムシBLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 69 (Wada Toge).
dubius BLANDFORD, 1894, *ibid.*, p. 70 (Kurigahara).
Distr.: Japan (Honshu, Shikoku, Kyushu).*P. rufus* BLANDFORD, 1894 ヒノキノキクイムシBLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 73 (Kashiwagi & Kobe).
Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Europe (France).*P. sannohensis* MURAYAMA, 1954 サンノヘキクイムシMURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 5, p. 161, 190 (Sannohe).
Distr.: Japan (Honshu, Shikoku, Kyushu).*P. seriatus* BLANDFORD, 1894 クスノキクイムシBLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 72 (Higo).
Distr.: Japan (Honshu, Shikoku, Kyushu).*P. shotoensis* MURAYAMA, 1955 ショトウキクイムシMURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., 6, p. 88 (Ônuda, Shôdojima).
Distr.: Japan (Honshu, Shikoku).

Tribe POLYGRAPHINI ヨツメキクイムシ族

Genus *Polygraphus* ERICHSON, 1836 ヨツメキクイムシ属

Polygraphus ERICHSON, 1836, Arch. Naturg., 2, p. 57 (type-species: *Hylesinus pubescens* FABRICIUS, 1792 = *Dermestes polygraphus* LINNÉ, 1758).

P. fulvipennis NIJJIMA, 1941 ヤツガタケキクイムシ

NIJJIMA, 1941, Ins. Matsum., 15, p. 131 (Yatsugatake).

Distr.: Japan (Honshu).

P. gracilis NIJJIMA, 1909 アカエゾキクイムシ

NIJJIMA, 1909, J. Coll. Agr., Tohoku imp. Univ., 3, p. 136 (Mihoni, Teshio).

Distr.: Japan (Hokkaido), Sakhalin, Siberia.

P. horyurensis MURAYAMA, 1937 ホリュウリキクイムシ (チョウセンキクイムシ)

MURAYAMA, 1937, Tenthredo, 1, p. 368 (Horyuri in Korea).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Korea, Taiwan, China.

P. jezoensis NIJJIMA, 1909 エゾキクイムシ

NIJJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 135 (Tayoroma, Onupunai, Tomakomai).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Kamtchatka, Siberia.

P. kisoensis NIJJIMA, 1941 キソキクイムシ

NIJJIMA, 1941, Ins. Matsum., 15, p. 131 (Kiso Kais Wald).

Distr.: Japan (Honshu).

P. magnus MURAYAMA, 1956 オオヨツメキクイムシ

MURAYAMA, 1956, Bull. Fac. Agr. Yamaguchi Univ., 7, pp. 282, 290 (Nishimata, Kochi).

Distr.: Japan (Shikoku).

P. meakanensis NIJJIMA, 1935 メアカンキクイムシ

NIJJIMA, 1935, Hokkaido Ringyo Kaihō, 33 (1), p. 2 (Mt. Meakan).

Distr.: Japan (Hokkaido, Honshu).

P. nigrielytris NIJJIMA, 1913 ナナカマドノキクイムシ

NIJJIMA, 1913, Trans. Sapporo nat. Hist. Soc., 5, p. 2 (Teshio).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Kamtchatka.

P. parvulus MURAYAMA, 1956 ヒメヨツメキクイムシ

MURAYAMA, 1956, Bull. Fac. Agr. Yamaguti Univ., 7, pp. 283, 292 (Fukuoka City, Hebitani (Kochi Pref.)).

Distr.: Japan (Honshu, Shikoku, Kyushu).

P. polygraphus (LINNÉ, 1758) ヨツメキクイムシ

LINNÉ, 1758, Syst. nat. ed. 10, 2, p. 562 (Europe).

Distr.: Japan (Hokkaido), Sakhalin, Siberia, China, Europe.

P. proximus BLANDFORD, 1894 トドマツノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 75 (Sapporo).

oblongus BLANDFORD, 1894, *ibid.*, p. 75 (Subashiri).

miser BLANDFORD, 1894, *ibid.*, p. 76 (Nikko).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Sakhalin, Siberia, Korea, China.

- P. shariensis* NIIZIMA, 1941 シャリキクイムシ
NIIZIMA, 1941, Ins. Matsum., 15, p. 127 (Kitami).
Distr.: Japan (Hokkaido, Honshu).
- P. squamulatus* NIIZIMA, 1941 チンナイキクイムシ
NIIZIMA, 1941, Ins. Matsum., 15, p. 129 (Chinnai and Hokkaido).
Distr.: Japan (Hokkaido), Sakhalin.
- P. ssiori* NIIZIMA, 1909 サクラノキクイムシ
NIIZIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 132 (Tomakomai, Sapporo).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Sakhalin.
- P. subopacus* THOMSON, 1871 トウヒノキクイムシ
THOMSON, 1871, Opusc. Ent., 4, p. 393 (Europe).
sachalinensis EGGERS, 1926, Ent. Bl., 22, p. 135 (Aihara, Sapporo).
Distr.: Japan (Hokkaido), Sakhalin, Kamtchatka, Siberia, Korea, China, Europe.
- P. uchimappensis* MURAYAMA, 1956 アオモリキクイムシ
MURAYAMA, 1956, Bulll. Fac. Agr. Yamaguti Univ., 7, p. 282.
granulatus NIIZIMA, 1941, Ins. Matsum., 15, p. 125 (nec EGgers, 1932) (Uchimappe-
staats wald bei Aomori).
japonicus NUNBERG, 1956, Ann. Zool. Polsc., 16, p. 208.
Distr.: Japan (Honshu).
- Genus *Nipponopolygraphus* NOBUCHI, 1981 ニッポンヨツメキクイムシ属
Nipponopolygraphus NOBUCHI, 1981, Kontyû, Tokyo, 49, p. 12 (type-species: *Nipponopoly-
graphus kaimochii* NOBUCHI, 1981).
- N. kaimochii* NOBUCHI, 1981 カイモチキクイムシ
NOBUCHI, 1981, Kontyû, Tokyo, 49, p. 13 (Shionomisaki).
Distr.: Japan (Honshu).

Subfamily IPINAE イピニ亞科

Tribe CRYPHALINI コキクイムシ族

Genus *Margadillius* HOPKINS, 1915 ミナミコキクイムシ

Margadillius HOPKINS, 1915, U.S. Dep. Agr., Rep. 99, p. 37 (type-species: *Margadillius
margadiliaonis* HOPKINS, 1515)

M. corpulentus (SAMPSON, 1919) オオハマボウノコキクイムシ
SAMPSON, 1919, Ann. Mag. nat. Hist., 9(4), p. 113 (*Cryphalus*) (India, Nilgiri Hills)
Distr.: Ryukyu Is. (Iejima, Okinawa, Iriomote), Philippines, Malaya, Java, India,
Africa.

Genus *Trypophloeus* FAIRMAIRE, 1868 トゲバネコキクイムシ属

Trypophloeus FAIRMAIRE, 1868, Faune Ent. France, 4, p. 105 (type-species: *Bostrichus
binodulus* RATZEBURG, 1937 = *Bostrichus asperatus* GYLLENHAL, 1813).
Cryphalus WOOD, 1954, Univ. Kansas Sci. Bull., 36, p. 987 (nec ERICHSON, 1836).

T. kurencovi NUNBERG, 1956 ポプラコキクイムシ
NUNBERG, 1956, Ann. Zool. Polsc., 16, p. 288.
populi KURENZOV, 1941, Bark-beetles Far East USSR, p. 164 (Upper course of the
River Suputinka) (nec HOPKINS, 1915).

Distr.: Japan (Hokkaido), Siberia.

Genus *Ernoporus* THOMSON, 1859 コブキクイムシ属

Ernoporus THOMSON, 1859, Skandinaviens Coleoptera, 1, p. 147 (type-species: *Bostrius tiliae* PANZER, 1793).

E. acanthopanaxi NIJJIMA, 1913 ウコギノコキクイムシ

NIJJIMA, 1913, Trans. Sapporo nat. Hist. Soc., 5, p. 4 (Sapporo).

Distr.: Japan (Hokkaido).

E. longus EGGLERS, 1926 ハンノナガコキクイムシ (ハンノナガキクイムシ)

EGGLERS, 1926, Ent. B1., 22, p. 136 (Tomakomai).

Distr.: Japan (Hokkaido), Siberia, Kamtchatka.

E. japonicus NOBUCHI, 1966 ツヤコキクイムシ

NOBUCHI, 1966, Bull. Gov. For. Exp. Sta., (185), p. 52 (Arashiyama, Kyoto City).

Distr.: Japan (Honshu).

E. shimanensis MURAYAMA, 1953 シマネコキクイムシ

MURAYAMA, 1953, Bull. Fac. Agr. Yamaguti Univ., 4, p. 37 (Suzuno-ohtani, Shimane Pref.).

Distr.: Japan (Honshu).

Genus *Ernopericus* BERGER, 1917 クロコキクイムシ属

Ernopericus BERGER, 1917, Rev. Russe Ent., 16, p. 242 (type-species: *Ernopericus spes-sivtzevi* BERGER, 1917).

Ernopocerus BALACHOWSKY, 1949, Faune de France, 50, p. 211 (type-species: *Ernopus caucasicus* LINDEMANN).

E. ater NOBUCHI, 1975 クロコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 42 (Sogoya, Saitama Pref.).

Distr.: Japan (Honshu).

E. alniphagus NOBUCHI, 1975 ヤマハンノキコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 43 (Konseitoge, Gunma Pref.).

Distr.: Japan (Honshu).

E. takahashii NOBUCHI, 1975 ナナカマドノコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 43 (Mt. Daisetsu, Hakkoda).

Distr.: Japan (Hokkaido, Honshu).

Genus *Ericryphalus* HOPKINS, 1915 トガリムネコキクイムシ属

Ericryphalus HOPKINS, 1915, U. S. Dep. Agr., Off. Secr. Rep., 99, p. 38 (type-species: *Ericryphalus henshawi* HOPKINS, 1915).

E. elongatus NOBUCHI, 1975 ホソナガコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 42 (Takao, Yashirogoya, Mt. Amagi, Mt. Hiko, Ukita).

Distr.: Japan (Honshu, Kyushu).

Genus *Scolytogenes* EICHHOFF, 1878 カクヘキコキクイムシ属

Scolytogenes EICHHOFF, 1878, Mém. Soc. roy. Sci. Liége (2) 8, pp. 475, 497 (type-species: *Scolytogenes darwini* EICHHOFF, 1878).

Cryphalomorphus SCHAFUSS, 1890, Tijd. Ent., 34, p. 12 (type-species: *Cryphalomorphus communis* SCHAFUSS, 1890).

Ernocryphalus MURAYAMA, 1958, Bull. Fac. Agr. Yamaguti Univ., 9, p. 934 (type-species: *Ernocryphalus birosimensis* MURAYAMA, 1958).

S. badius (NOBUCHI, 1975) クリイロコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 44 (*Cryphalomorphus*) (Inugakitoge, Tokaimura, Asakawa, Mt. Hiko).

Distr.: Japan (Honshu, Kyushu).

S. birosimensis (MURAYAMA, 1958) ピロウジマコキクイムシ

MURAYAMA, 1958, Bull. Fac. Agr. Yamaguti Univ., 9, p. 934 (*Ernocryphalus*) (Isl. Biro).

Distr.: Japan (Honshu, Shikoku, Kyushu).

S. camelliae (NOBUCHI, 1975) ツバキノコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 45 (*Cryphalomorphus*) (Tainagoya, Kawada).

Distr.: Japan (Honshu).

S. expers (BLANDFORD, 1894) タブノコキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 84 (*Hypothenemus*) (Kumamoto, Nagasaki).

Distr.: Japan (Shikoku, Kyushu).

S. fujisanus (NOBUCHI, 1975) フジコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 46 (*Cryphalomorphus*) (Mt. Fuji).

Distr.: Japan (Honshu).

S. candidus (NOBUCHI, 1975) イヅツヤコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 47 (*Cryphalomorphus*) (Gyokuto).

Distr.: Japan (Kyushu).

S. fulvipennis (NOBUCHI, 1975) キバネコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 47 (*Cryphalomorphus*) (Yona, Misato, Omotodake, Wulai).

Distr.: Ryukyu Is. (Okinawa, Ishigaki), Taiwan.

S. insularis (NOBUCHI, 1975) ツヤコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 48 (*Cryphalomorphus*) (Kawada).
Distr.: Japan (Mikura Is.).

S. punctatulus (NOBUCHI, 1976) フクオカコキクイムシ

NOBUCHI, 1976, Kontyû, Tokyo, 44, p. 72.

punctatus NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 49 (nec SCHEDL, 1951) (*Cryphalomorphus*) (Mt. Inugaki).

Distr.: Japan (Kyushu).

S. scolytomimoides (NOBUCHI, 1975) テイカカズラノコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 50 (*Cryphalomorphus*) (Asakawa, Mt. Tachibana, Kibune).

Distr.: Japan (Honshu, Kyushu).

Genus *Cryphalus* ERICHSON, 1836 コキクイムシ属

Cryphalus ERICHSON, 1836, Arch. Naturg., 2, p. 61 (type-species: *Bostricus asperatus* GYLLENHAL, 1813).

Taenioglyptes BEDEL, 1888, Faune des Coléoptères du Bassin de la Seine, 6, p. 398 (type-species: *Bostrichus abietis* RATZEBURG, 1837).

C. abbreviatus SCHEDL, 1943 ガジュマルノコキクイムシ

SCHEDL, 1943, Ent. B1., 39, p. 35 (Philippines).

Distr.: Japan (Honshu), Ryukyu Is. (Okinawa), Philippines.

C. abietis (RATZEBURG, 1837) モミノコキクイムシ

RATZEBURG, 1837, Die Forstsektien, 1, p. 163 (*Bostrichus*) (Europe).

Distr.: Japan (Honshu), Europe.

C. aquilonius (NOBUCHI, 1975) サワシバノコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 52 (*Taenioglyptes*) (Nigorikawa).

Distr.: Japan (Hokkaido).

C. babai (MURAYAMA, 1961) ババコキクイムシ

MURAYAMA, 1961, Akitu, 10, p. 29 (Noo).

Distr.: Japan (Honshu).

C. bicarinatus (NOBUCHI, 1975) フタスジコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 53 (*Taenioglyptes*) (Arakawa).

Distr.: Japan (Kyushu).

C. carpini BERGER, 1916 シデノコキクイムシ

BERGER, 1916, Rev. Russe d'Ent., 16, p. 324 (Ussuri).

Distr.: Japan (Honshu), Korea, Siberia.

C. cryptomeriae NIIJIMA, 1907 スギノコキクイムシ

NIIJIMA, 1907, Verh. zool. bot. Ges. Wien, 1907, p. 91 (Kumamoto).

Distr.: Japan (Honshu, Kyushu).

C. dissimilis (NOBUCHI, 1975) ハバビロコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 57 (*Taenioglyptes*) (Mt. Fuji).

Distr.: Japan (Honshu).

C. exiguus BLANDFORD, 1894 クワノコキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1984, p. 82 (Fukushima),

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Kuril Is., Korea, Taiwan, China.

C. ficatorius MURAYAMA, 1954 アコウノキクイムシ

MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 9, p. 933 (Muroto).

Distr.: Japan (Shikoku).

C. fulvus NIJJIMA, 1907 キイロコキクイムシ

NIJJIMA, 1907, Verh. zool. bot. Ges. Wien, 1907, p. 92 (Yatsuo-Berg).

pini EGGLERS, 1921, Ent. B1., 17, p. 39 (Kiautschou).

Distr.: Japan (Honshu, Shikoku, Kyushu), Bonin Is., Ryukyu Is., Korea, China.

C. hirsutus (NOBUCHI, 1975) フサゲコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 52 (*Taenioglyptes*) (Komanoyu).

Distr.: Japan (Honshu).

C. jehorensis MURAYAMA, 1939 ネツカコキクイムシ

MURAYAMA, 1939, Ann. zool. Japon., **18**, pp. 140, 143 (Shōtoku).
 Distr.: Japan (Hokkaido, Honshu, Shikoku), Korea, China.

- C. jezoensis* INOUYE et NOBUCHI, 1957 エゾマツノコキクイムシ
 INOUYE et NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (103), p. 49 (Sounkyo).
 Distr.: Japan (Hokkaido).
- C. juglans* NIIZIMA, 1913, クルミノコキクイムシ (クルミノキクイムシ)
 NIIZIMA, 1913, Trans. Sapporo nat. Hist. Soc., **5**, p. 3 (Prov. Kushiro).
 Distr.: Japan (Hokkaido, Honshu).
- C. kagoshimensis* (NOBUCHI, 1975) カゴシマコキクイムシ
 NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 56 (*Taenioglyptes*) (Satamisaki).
 Distr.: Japan (Kyushu).
- C. kyotoensis* NOBUCHI, 1966 キヨウトコキクイムシ
 NOBUCHI, 1966, Bull. Gov. For. Exp. Sta., (185), p. 53 (Mizorogaike).
 Distr.: Japan (Honshu).
- C. laricis* NIIZIMA, 1909 カラマツノコキクイムシ
 NIIZIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 152 (Nopporo).
 Distr.: Japan (Hokkaido, Honshu), Korea.
- C. longisetosus* (NOBUCHI, 1975) ケナガコキクイムシ
 NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 57 (*Taenioglyptes*) (Hirogawara, Karisakatoge, Mt. Fuji).
 Distr.: Japan (Honshu).
- C. malus* NIIZIMA, 1909 リンゴノコキクイムシ (リンゴノキクイムシ)
 NIIZIMA, 1909, J. Agr. Coll. Tohoku imp. Univ., **3**, p. 144 (Hakodate).
 Distr.: Japan (Hokkaido, Honshu), Korea, China.
- C. meridionalis* (NOBUCHI, 1975) オキナワコキクイムシ
 NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 55 (*Taenioglyptes*) (Yona, Omotodake).
 Distr.: Ryukyu Is. (Ishigaki).
- C. montanus* NOBUCHI, 1964 ミヤマコキクイムシ
 NOBUCHI, 1964, Bull. Gov. For. Exp. Sta., (171), p. 129 (Shirotoge, Sanpeitoge, Fujimitoge, Mt. Kinpu, Karisakatoge).
 Distr.: Japan (Honshu).
- C. nipponensis* INOUYE et NOBUCHI, 1957 ニホンコキクイムシ
 INOUYE et NOBUCHI, 1957, Bull. Gov. For. Exp. Sta., (103), p. 51 (Sounkyo).
 Distr.: Japan (Hokkaido).
- C. parvulus* NIIZIMA, 1910 トウキヨウコキクイムシ
 NIIZIMA, 1910, Trans. Sapporo nat. Hist. Soc., **3**, p. 8 (Tokio).
 Distr.: Japan (Honshu).
- C. piceae* (RATZEBURG, 1837) トドマツコキクイムシ (トウヒノコキクイムシ)
 RATZEBURG, 1837, Die Forstinsekten **1**: 163 (*Bostrichus*) (Europe).
hattorii KÔNO, 1939, Ins. Matsum., **12**, p. 67 (Nakahira).
 Distr.: Japan (Hokkaido, Honshu, Shikoku), Sakhalin, Kuril Is., Korea, China.
- C. piceus* EGGLERS, 1926 ジョウザンコキクイムシ
 EGGLERS, 1926, Ent. Bl., **22**, p. 136 (Hokkaido, Jozankei).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Kuril Is., Siberia, China.

C. pulchellus (NOBUCHI, 1975) トチノコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 54 (*Taenioglyptes*) (Higashikawa).

Distr.: Japan (Honshu).

C. rhusii NIJIMA, 1909 ツタウルシノコキクイムシ

NIJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 145 (Chitose).

Distr.: Japan (Hokkaido, Honshu).

C. sawadai NOBUCHI et K. TAKAHASHI, 1965 サワダコキクイムシ

NOBUCHI et K. TAKAHASHI, 1965, Akitu, 13, p. 3 (Mt. Hayachine, Mt. Fuji).

Distr.: Japan (Honshu).

C. scopiger BERGER, 1916 ブナノコキクイムシ

BERGER, 1916, Rev. Russ. d'Ent., 16, p. 228 (Ussuri).

Distr.: Japan (Honshu, Kyushu), Siberia, Korea.

C. sordidus (NOBUCHI, 1975) ハヤチネコキクイムシ

NOBUCHI, 1975, Bull. Gov. For. Exp. Sta., (277), p. 51 (*Taenioglyptes*) (Mt.

Hayachine).

Distr.: Japan (Honshu).

C. yamaguchi INOUE et NOBUCHI, 1957 ヤマグチコキクイムシ

INOUE et NOBUCHI, 1957, Bull. Gov. For. Exp. Sta., (103), p. 55 (Sounkyo).

Distr.: Japan (Hokkaido).

Genus *Hypothenemus* WESTWOOD, 1834 チビコキクイムシ属

Hypothenemus WESTWOOD, 1834, Trans. ent. Soc. London, 1, p. 34 (type-species: *Hypothenemus eruditus* WESTWOOD, 1834).

Stephanoderes EICHHOFF, 1872, Berl. ent. Z., 15, p. 132 (type-species: *Stephanoderes chapuisi* EICHHOFF, 1872).

H. amakusanus (MURAYAMA, 1934) アマクサコキクイムシ

MURAYAMA, 1934, Annot. zool. Japon., 14, p. 187 (Kadoyama).

Distr.: Japan (Honshu, Shikoku, Kyushu).

H. birmanus (EICHHOFF, 1878) ナガサキコキクイムシ

EICHHOFF, 1878, Mém. Soc. roy. Sci. Liége (2) 8, p. 486 (*Triarmocerus*) (Birma).

peritus BLANDFORD, 1894, Trans. ent. Soc. Londōn, 1894, p. 84 (Nagasaki).

Distr.: Japan (Honshu, Kyushu), Bonin Is., China, Philippines, Borneo, Burma, Malaya, Sumatra, Java, Oceania, Florida, Cuba.

H. chamaecypariae NIJIMA, 1910 ヒノキノコキクイムシ

NIJIMA, 1910, Tarns. Sapporo nat. Hist. Soc., 3, p. 10 (*Cryphalus (Hypothenemus)*) (Tokio).

Distr.: Japan (Honshu).

H. cosmoderoides MURAYAMA, 1961 エチゴコキクイムシ

MURAYAMA, 1961, Akitu, 10, p. 30 (Noo).

Distr.: Japan (Honshu).

H. eruditus WESTWOOD, 1836 チビコキクイムシ

WESTWOOD, 1836, Trans. ent. Soc. London, 1, p. 34 (England).

ehlersii EICHHOFF, 1878, Mém. Soc. Roy. Sci. Liége, (2) 8, p. 493 (*Stephanoderes*)

(Spain) (イチジクノコキクイムシ).
basjoo NIIZIMA, 1910, Trans. Sapporo Nat. Hist. Soc., 3, p. 9 (*Cryphalus*) (Tokio)
 (バショウノコキクイムシ).

Distr.: Japan (Honshu, Shikoku, Kyushu), Bonin Is., Asia to Africa, S. Europe, N.
 & S. America.

H. furukawai MURAYAMA, 1934 フルカワコキクイムシ

MURAYAMA, 1934, J. Chosen nat. Hist. Soc., 17, p. 3 (*Cryphalus*) (Chochiin).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Korea.

H. japonicus (NIIZIMA, 1910) ヤマトコキクイムシ

NIIZIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 10 (*Cryphalus (Hypothenemus)*)
 (Honshu).
 Distr.: Japan (Honshu).

H. kraunhiae (MURAYAMA, 1950) フジノコキクイムシ

MURAYAMA, 1950, Trans. Shikoku ent. Soc., 1, p. 50 (*Cryphalus (Stephanoderes)*)
 (Fuchū).
 Distr.: Japan (Honshu).

H. modestus (MURAYAMA, 1940) ノニレコキクイムシ

MURAYAMA, 1940, Annot. zool. Japon., 19, pp. 233, 236 (*Cryphalus (Stephanoderes)*)
 (Seiryū dans Jōhōl).
 Distr.: Japan (Honshu, Shikoku), Korea, NE. China.

H. oblongus (NIIZIMA, 1910) アカマツノコキクイムシ

NIIZIMA, 1910, Trans. Sapporo nat. Hist. Soc. 3, p. 9 (*Cryphalus (Hypothenemus)*)
 (Tokio).
 Distr.: Japan (Honshu).

H. sapporoensis (NIIZIMA, 1910) サッポロコキクイムシ

NIIZIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 3 (*Cryphalus (Hypothenemus)*)
 (Sapporo).
 Distr.: Japan (Hokkaido, Honshu).

H. tristis (EICHHOFF, 1875) クロヒメコキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., 18, p. 200 (*Stephanoderes*) (Japan).
 Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Macrocyphalus* NOBUCHI, 1981 オオガタコキクイムシ属

Macrocyphalus NOBUCHI, 1981, Kontyū, Tokyo, 49, p. 14 (type-species: *Macrocyphalus oblongus* NOBUCHI, 1981).

M. oblongus NOBUCHI, 1981 チャイロコキクイムシ

NOBUCHI, 1981, Kontyū, Tokyo, 49, p. 14 (Takao, Wakasugi, Oshima in Wakayama Pref.).

Distr.: Japan (Honshu, Kyushu).

Genus *Cosmoderes* EICHHOFF, 1878 ホソコキクイムシ属

Cosmoderes EICHHOFF, 1878, Stettin. ent. Ztg., 39, p. 387 (type-species: *Cosmoderes monilicollis* EICHHOFF, 1878).

C. consobrinus BLANDFORD, 1894 ホソコキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 86 (without locality).

Distr.: Japan (Shikoku).

Genus *Pseudocosmoderes* NOBUCHI, 1981 ホソコキクイムシモドキ属

Pseudocosmoderes NOBUCHI, 1981, Kontyû, Tokyo, **49**, p. 16 (type-species: *Pseudocosmoderes attenuatus* NOBUCHI, 1981).

P. attenuatus NOBUCHI, 1981 ツルノコキクイムシ

NOBUCHI, 1981, Kontyû, Tokyo, **49**, p. 17 (Shimizutoge, Yabitsutoge, Mt. Hikosan, Ohto, Asakawa).

Distr.: Japan (Honshu, Kyushu).

Tribe XYLOTERINI カレザイノキクイムシ族

Genus *Trypodendron* STEPHENS, 1830 カレザイノキクイムシ属

Trypodendron STEPHENS, 1830, Ill. British Ent., **3**, p. 353 (type-species: *Dermestes domesticus* LINNÉ, 1758).

Xyloterus ERICHSON, 1836, Arch. Naturg., **1**, p. 60 (type-species: *Bostrichus lineatus* OLIVIER, 1795).

T. ashuense (MURAYAMA, 1950) アシュウキクイムシ

MURAYAMA, 1950, Trans. Shikoku ent. Soc., **1**, p. 51 (*Xyloterus*) (Ashu).

Distr.: Japan (Hokkaido, Honshu).

T. gaimense (MURAYAMA, 1937) ガイマザイノキクイムシ

MURAYAMA, 1937, Tenthredo, **1**, p. 368 (*Xyloterus*) (Horyuri).

Distr.: Japan (Honshu), Korea.

T. lineatum (OLIVIER, 1795) シラベザイノキクイムシ

OLIVIER, 1795, Ent. Hist. nat. Ins., Coleopt., **4**, p. 18 (*Bostrichus lineatus*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Kamtchatka, Siberia, Korea, China, Europe, N. Africa, N. America.

T. obtusum EGGERS, 1939 スギザイノキクイムシ

EGGERS, 1939, Arb. morph. tax. Ent., **6**, p. 121 (Karafuto, Nopporo).

Distr.: Japan (Hokkaido, Honshu, Kyushu), Sakhalin.

T. proximum (NIIJIMA, 1909) クロツヤキクイムシ

NIIJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 165 (*Xyloterus*) (Jozankei).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Sakhalin, Siberia, Korea.

T. signatum (FABRICIUS, 1782) カシワノキクイムシ

FABRICIUS, 1782, Syst. Eleuth. **2**, p. 383 (*Apate*) (Europe).

quercus EICHHOFF var. *niponicum* BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, 124 (Miyamoshita, Oyayama).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Kuril Is., Sakhalin, Kamtchatka, Siberia, Europe.

Genus *Indocryphalus* EGGERS, 1939 ヨツメザイノキクイムシ属

Indocryphalus EGgers, 1939, Ark. Zool., **31** (A), 4, p. 6 (type-species: *Indocryphalus malaisei* EGgers, 1939).

Dendrotrypum SCHEDL, 1951, Mitt. forstl. Bunds-Ver. Mariabrum, **47**, p. 76 (type-species: *Trypodendron sordidum* BLANDFORD, 1893).

I. aceris (NIIJIMA, 1910) イタヤノキクイムシ

NIIJIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 4 (*Xyloterus*) (Sapporo).
Distr.: Japan (Hokkaido, Honshu).

I. dainichiensis (MURAYAMA, 1954) ダイニチキクイムシ
MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 5, pp. 167, 191 (*Xyloterus*) (Gifu,
Mt. Dainichi)
Distr.: Japan (Honshu).

I. majus (EGGERS, 1926) オオザイノキクイムシ
EGGERS, 1926, Ent. B1., 22, p. 148 (*Trypodendron*) (Tomakomai).
Distr.: Japan (Hokkaido, Honshu, Shikoku).

I. pubipennis (BLANDFORD, 1894) カナクギノキキクイムシ
BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 125 (*Trypodendron*) (Sapporo,
Kiga, Miyanoshita, Ichiuchi).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Kuril Is., Sakhalin, Korea,
Taiwan.

I. pulchellus (MURAYAMA, 1957) エゴザイノキクイムシ
MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., 7, p. 582 (*Xyloterus*) (Todoro,
Kochi).
Distr.: Japan (Shikoku).

I. sordidus (BLANDFORD, 1894) キシマキクイムシ
BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 577 (*Trypodendron*) (Japan).
Distr.: Japan (Honshu, Shikoku, Kyushu).

Tribe CRYPTURGINI ホソキクイムシ族

Genus *Crypturgus* ERICHSON, 1836 ホソキクイムシ属

Crypturgus ERICHSON, 1836, Arch. Naturg., 1, p. 60 (type-species: *Bostrichus pusillus*
GYLLENHAL, 1813),

C. pusillus (GYLLENHAL, 1813) トウヒノホソキクイムシ
GYLLENHAL, 1813, Insecta svec. 3, p. 371 (*Bostrichus*) (Europe).
Distr.: Japan (Hokkaido, Honshu), Kuril Is., Sakhalin, Siberia, Korea, Taiwan,
China, Himalaya, Europe, Africa.

C. tuberosus NIIJIMA, 1909 カバイロホソキクイムシ
NIIJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 139 (Onupunai, Tomakomai,
Berg Mackarinupuri).
Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia.

Genus *Lymantor* LOEVENDAL, 1889 リマントール属

Lymantor LOEVENDAL, 1889, Ent. Med., 2, p. 68 (type-species: *Lymantor sepicola*
LOEVENDAL, 1889).

L. kabei MURAYAMA, 1955 カベアトマルキクイムシ
MURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., 6, p. 87 (Shiraito, Kumamoto).
Distr.: Japan (Kyushu).

Genus *Taphrorychus* EICHHOFF, 1878 ナガシンクイモドキ属

Taphrorychus EICHHOFF, 1878, Mém. Soc. roy. Sci, Liége (2) 8, p. 204 (type-species:
Bostrichus bicolor HERBST, 1793).

T. apatoides (EICHHOFF, 1875) ニセナガシンクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., 18, p. 201 (*Dryocoetes?*) (without locality).
Distr.: Japan.

T. coffeeae (EGGERS, 1923) コーヒーキクイムシ

EGGERS, 1923, Zool. Meded. R. Mus. nat. Hist. Leiden, 7, p. 161 (*Dryocoetes*)
(Java).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan, Borneo, Malaya, Java, India,
New Guinea.

T. dinoderoides (BLANDFORD, 1894) ナガシンクイモドキ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 97 (*Dryocoetes*) (Ichiuchi).
Distr.: Japan (Kyushu).

T. modestus (BLANDFORD, 1894) クロケブカキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 94 (*Dryocoetes*) (Nikko).
Distr.: Japan (Honshu, Shikoku).

T. striatus (NOBUCHI, 1966) ズイノキクイムシ

NOBUCHI, 1966, Bull. Gov. For. Exp. Sta., (185), p. 52 (Mt. Homan, Fukuoka).
Distr.: Japan (Kyushu).

Genus *Pseudopoecilips* MURAYAMA, 1957 ニセアトマルキクイムシ属

Pseudopoecilips MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., 8, p. 614 (type-species: *Pseudopoecilips mikuniyamensis* MURAYAMA, 1957).

P. mikuniyamensis MURAYAMA, 1957 ミクニヤマキクイムシ

MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., 8, p. 616 (Mt. Mikuni, Niigata,
Kamikochi).
Distr.: Japan (Honshu).

P. pilosus (BLANDFORD, 1894) プナノコアトマルキクイムシ (ツガノキクイムシ)

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 92 (*Dryocoetes*) (Nikko).
pilosus MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., 8, pp. 612, 620 (nec
BLANDFORD, 1884) (Mt. Nagatani, Gifu).
Distr.: Japan (Honshu, Kyushu).

P. taradakensis MURAYAMA, 1957 タラダケキクイムシ

MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., 8, pp. 616, 618 (Mt. Tara,
Shimizu town).
Distr.: Japan (Shikoku, Kyushu).

Genus *Dryocoetes* EICHHOFF, 1864 アトマルキクイムシ属

Dryocoetes EICHHOFF, 1864, Berl. ent. Z., 8, p. 38 (type-species: *Bostrius autographus*
RATZEBURG, 1837).

D. affinis BLANDFORD, 1894 トオホクキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 93 (Oyayama).
Distr.: Japan (Honshu, Kyushu).

D. autographus (RATZEBURG, 1837) トウヒノネノキクイムシ

RATZEBURG, 1837, Die Forstinsekten, p. 160 (Europe).
Distr.: Japan (Hokkaido, Honshu, Kyushu), Sakhalin, Siberia, Korea, Taiwan,
China, Europe, N. America.

- D. baicalicus* REITTER, 1899 グイマツアトマルキクイムシ
 REITTER, 1899, Dtsch. ent. Z., **1899**, p. 286 (Irkutsk).
 Distr.: Japan (Honshu), Sakhalin, Siberia, Mongolia.
- D. brevipilosus* MURAYAMA, 1957 ツガアトマルキクイムシ
 MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., **8**, pp. 601, 604 (Mt. Manza, Gunma).
 Distr.: Japan (Honshu).
- D. cristatus* INOUYE et NOBUCHI, 1959 ホッカイドウキクイムシ
 INOUYE et NOBUCHI, 1959, Ins. Matsum., **22**, p. 122 (Nopporo).
 Distr.: Japan (Hokkaido).
- D. hectographus* REITTER, 1913 アカアトマルキクイムシ
 REITTER, 1913, Best.-tab. Borkenkäfer, p. 76 (Transbaikalien).
 Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, China, Europe.
- D. karamatsu* SAWAMOTO, 1940 カラマツアトマルキクイムシ
 SAWAMOTO, 1940, Ins. Matsum., **14**, p. 102 (Mitonogawa, Nagano).
 Distr.: Japan (Honshu).
- D. niijimai* NOBUCHI, 1959 ニイジマアトマルキクイムシ
 NOBUCHI, 1959, Akitu, **8**, p. 11 (Misono, Yamabe).
 Distr.: Japan (Honshu, Hokkaido).
- D. picipennis* EGGERS, 1926 ベニイタヤノキクイムシ
 EGgers, 1926, Ent. B1., **22**, p. 138 (Jozankei).
 Distr.: Japan (Hokkaido, Kyushu).
- D. pini* NIIJIMA, 1909 ハイマツアトマルキクイムシ
 NIJJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 152 (Makkarinupuri).
 Distr.: Japan (Hokkaido, Honshu, Shikoku), Sakhalin, Korea.
- D. rugicollis* EGGERS, 1926 アトマルキクイムシ
 EGgers, 1926, Ent. B1., **22**, p. 137 (Sachalin (Osawa), Kurilen).
 Distr.: Japan (Hokkaido, Honshu, Shikoku), Kuril Is., Sakhalin, Siberia, China.
- D. striatus* EGGERS, 1933 トドマツアトマルキクイムシ
 EGgers, 1933, Ent. B1., **29**, p. 8 (Wladiwostok).
abietinus KÔNO et TAMANUKI, 1939, Ins. Matsum., **13**, p. 90 (Sachalin (Naibuchi, Konuma, Kashiho, Horo), Hokkaido (Toman)).
 Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, NE. China.
- D. uniseriatus* EGGERS, 1926 アカマツアトマルキクイムシ
 EGgers, 1926, Ent. B1., **22**, p. 145 (Sapporo).
 Distr.: Japan (Hokkaido, Honshu).
- Genus *Cyrtogenius* STROHMEYER, 1910 ホソアトマルキクイムシ属
- Cyrtogenius* STROHMEYER, 1910, Ent. B1., **6**, p. 127 (type-species: *Cyrtogenius bicolor* STROHMEYER, 1910).
- Carposinus* HOPKINS, 1915, U. S. Dept. Agr., Rep., **99**, p. 47 (type-species: *Carposinus pini* HOPKINS, 1915).
- Orosiotes* NIJJIMA, 1917, A collection of essays for Mr. Yasushi MIWA, written in commemoration of his sixtieth birthday, p. 1 (type-species: *Orosiotes kumamotoensis* NIJJIMA 1917).

Pelicerus EGGERS, 1923, Zool. Meded. R. Mus. nat. Hist. Leiden, 7, p. 216 (type-species: *Lepicerus nitidus* HAGEDORN, 1910).

C. brevior (EGGERS, 1927) フィリピンキクイムシ

EGGERS, 1927, Philip. J. Sci., 33, p. 86 (*Pelicerus*) (Philippinen (Masbate, Mindanao)).
philippinensis EGGERS, 1927, Philip. J. Sci., 33, p. 87 (*Pelicerus*) (Philippines (Catanduanes)).

Distr.: Japan (Honshu, Kyushu), Ryukyu Is. (Okinawa), Korea, Philippines, Sumba.

C. kumamotoensis (NIIJIMA, 1917) クマモトアトマルキクイムシ

NIIJIMA, 1917, A collection of essays for Mr. Yasushi MIWA, written in commemoration of his sixtieth birthday, p. 2 (*Orosiotes*) (Kumamoto).

Distr.: Japan (Kyushu).

C. luteus (BLANDFORD, 1894) コガネキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 94 (without locality).

Distr.: Japan (Honshu), Korea, China.

Genus *Coccotrypes* EICHHOFF, 1878 タネノキクイムシ属

Coccotrypes EICHHOFF, 1878, Mém. Soc. roy. Sci. Liége (2) 8, pp. 57, p. 308 (type-species: *Bostrichus dactyliperda* FABRICIUS, 1801).

C. dactyliperda (FABRICIUS, 1801) タネノキクイムシ

FABRICIUS, 1801, Syst. Eleuth. 2, p. 387 (*Bostrichus*).

Distr.: Japan (Bonin Is.), Cosmopolitan in tropical and subtropical areas.

C. perditor BLANDFORD, 1894 ナガサキタネノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 99 (Nagasaki).

Distr.: Japan (Kyushu).

Genus *Poecilips* SCHAUFUSS, 1897 ヒメアトマルキクイムシ属

Poecilips SCHAUFUSS, 1897, Berl. ent. Z., 42, p. 110 (type-species: *Poecilips sannio* SCHAUFUSS, 1897).

P. advena (BLANDFORD, 1894) ヒメアカキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 100 (*Coccotrypes*) (Nagasaki).
persicae (HOPKINS, 1915), U. S. Dept. Agr., Rep., 99, p. 45 (*Thamnurgides*) (Honolulu).

Distr.: Japan (Kyushu, Bonin Is.), Ryukyu Is. (Okinawa), Philippines, Borneo, Malaya, Java, Caroline Is., Hawaii, Samoa, Queensland, Tropical Africa, N. America, Cuba.

P. babai (MURAYAMA, 1961) ババキクイムシ

MURAYAMA, 1961, Akitu, 10, p. 29 (*Thamnurgus*) (Mt. Kinpuku).

Distr.: Japan (Honshu).

P. cardamomi (SCHAUFUSS, 1905) クリノミキクイムシ

SCHAUFUSS, 1905, Insekten-Börse, 22, p. 87 (*Coccotrypes*) (Ceylon).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan, China, Vietnam, Malaya, Nepal, Borneo, Sumatra, Ceylon, S. India.

P. fagi NOBUCHI, 1959 プナヒメアトマルキクイムシ

NOBUCHI, 1959, Akitu, 8, p. 12 (River Daido).

Distr.: Japan (Honshu).

P. graniceps (EICHHOFF, 1877) ドングリキクイムシ

EICHHOFF, 1877, Dtsch. ent. Z., 21, p. 120 (*Dryocoetes*) (Japan).

Distr.: Japan (Honshu, Kyushu).

P. naidaijinensis (MURAYAMA, 1957) ナイダイジンキクイムシ

MURAYAMA, 1957, Bull. Fac. Agr. Yamaguti Univ., 8, pp. 603, 605 (*Dryocoetes*) (Mt. Naidaijin, Niimi).

Distr.: Japan (Honshu, Kyushu).

P. norimasanus (MURAYAMA, 1954) ノリマサキクイムシ

MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 5, pp. 173, 192 (*Dryocoetes*) (Norimasa).

Distr.: Japan (Honshu).

P. nubilus (BLANDFORD, 1894) ケブカキクイムシ (ヤマトキクイムシ)

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 95 (*Dryocoetes*) (Kiga, Suyama).

japonicus EGGERS, 1926, Ent. B1., 22, p. 145 (Takao-Berg bei Hachiōji).

Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Taiwan.

P. oblongus EGGERS, 1927 サザンカコアトマルキクイムシ

EGGERS, 1927, Philip. J. Sci., 30, p. 83 (Philippines).

Distr.: Japan (Kyushu), Philippines.

Tribe XYLEBORINI ザイノキクイムシ族

Genus *Xylosandrus* REITTER, 1913 ハバビロザイノキクイムシ属

Xylosandrus REITTER, 1913, Wien. ent. Ztg., 32, p. 83 (type-species: *Xyleborus morigerus* BLANDFORD, 1894).

X. borealis NOBUCHI, 1981 ヒメハネミジカキクイムシ

NOBUCHI, 1981, Bull. For. & For. Prod. Res. Inst., (314), p. 34, ♀ (Mt. Takao, Ikeda).

Distr.: Japan (Honshu, Kyushu), Korea.

X. brevis (EICHHOFF, 1877) ハネミジカキクイムシ

EICHHOFF, 1877, Dtsch. ent. Z., 21, p. 211, ♀ (*Xyleborus*) (Hagi).

cucullatus BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 121, ♂ (*Xyleborus*) Kurigahara, Konose).

Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Taiwan, Thailand.

X. compactus (EICHHOFF, 1875) シイノコキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., 18, p. 202, ♀ (Japan).

Distr.: Japan (Honshu, Shikoku, Kyushu, Bonin Is.), Ryukyu Is., Taiwan, China, Philippines, Tonkin, Indochina, India, Ceylon, Malaya, Borneo, Sumatra, Java, Celebes, Fiji, Samoa, Hawaii, Seychelles, Mauritius, Madagascar, Africa, United States.

X. crassiusculus (MOTSCHULSKY, 1866) サクキクイムシ (サカクレノキクイムシ)

MOTSCHULSKY, 1866, Bull. Soc. imp. Nat. Moscou, 39, p. 403.

semiopacus EICHHOFF, 1878, Mém. Soc. roy. Sci. Liège 2 (8), 334, ♀ (Hagi).

ebriosus NIIZIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 154, ♀ (Sapporo).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Bonin Is.), Korea, China. Tropicopolitan.

X. germanus (BLANDFORD, 1894) ハンノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 106, ♀ (*Xyleborus*) (Oyama, Nikko, Subashiri, Kiga, Miyano Shita).

orbatus BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 123, ♂ (*Xyleborus*) (Kurigahara).

Distr.: Japan (Honshu, Shikoku, Kyushu), Kuril Is., Ryukyu Is., Korea, China, Vietnam, U.S.A., Central Europe.

Genus *Xyleborus* EICHHOFF, 1864 ザイノキクイムシ属

Xyleborus EICHHOFF, 1864, Berl. ent. Z., 8, p. 37 (type-species: *Bostrichus monographus* FABRICIUS, 1792).

X. adumbratus BLANDFORD, 1894 ツヤナシキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 115, ♀ (Nagasaki, Hitoyoshi, Oyama, Subashiri).

septentrionalis NIJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 162, ♀ (Tayoroma, Yoshino).

Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Andaman Is., Germany.

X. amputatus BLANDFORD, 1894 ツヅミキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 575, ♀ (Higo).

Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan.

X. andrewesi BLANDFORD, 1896 ホソバネキクイムシ

BLANDFORD, 1896, Trans. ent. Soc. London, 1896, p. 227, ♀ (India, Belgaum).

Distr.: Japan (Shikoku, Kyushu), Ryukyu Is. (Okinawa, Ishigaki), Philippines, Borneo, Thailand, Burma, Malaya, Andaman Is., Java, Seychellus, Bangla Desh, India, Ceylon.

X. apicalis BLANDFORD, 1894 ニレザイノキクイムシ (リングマルキクイムシ)

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 105, ♀ (without locality).

Distr.: Japan (Hokkaido, Honshu, Kyushu), Korea.

X. aquilus BLANDFORD, 1894 アカマツザイノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 109, ♀ (Oyayama, Hitoyoshi, Kashiwagi).

Distr.: Japan (Honshu, Shikoku, Kyushu), Korea, Taiwan, China.

X. ashuensis MURAYAMA, 1954 アシュウキクイムシ

MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 5, p. 193, ♀ (Ashu).

Distr.: Japan (Honshu, Shikoku, Kyushu).

X. atratus EICHHOFF, 1875 クワノキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., 18, p. 201, ♀ (Hagi).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Taiwan, China, Indochina.

X. attenuatus BLANDFORD, 1894 サクラノホソキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 114, ♀ (Nikko).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Taiwan.

X. bicolor BLANDFORD, 1894 フタイロキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 113, ♀ (Nagasaki, Inasa).

Distr.: Japan (Shikoku, Kyushu), Philippines, Borneo, Burma, Malaya, Java, Samoa,

India, Andamans.

X. calamoides MURAYAMA, 1934 ナガオキクイムシ

MURAYAMA, 1934, Annot. zool. Japon., **14**, p. 191, ♀, ♂ (Nagao).
Distr.: Japan (Kyushu).

X. canus NIJIMA, 1909 カヌスキクイムシ

NIJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 161, ♀ (Sapporo).
Distr.: Japan (Hokkaido, Honshu).

X. collis NIJIMA, 1910 コリスキクイムシ

NIJIMA, 1910, Trans. Sapporo nat. Hist. Soc., **3**, p. 12, ♀ (Kumanotaira).
Distr.: Japan (Honshu).

X. concisus BLANDFORD, 1894 アカガシノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 107, ♀ (without locality).
Distr.: Japan (Kyushu), Philippines, Borneo, Malaya, Sumatra, Java, Burma,
Bengal.

X. cornivorus MURAYAMA, 1953 ミズキノキクイムシ

MURAYAMA, 1953, Trans. Shikoku ent. Soc., **3**, p. 157, ♂, ♀ (Irazuyama).
Distr.: Japan (Honshu, Shikoku).

X. defensus BLANDFORD, 1894 シイノホソキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 118, ♀ (Sapporo).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

X. exesus BLANDFORD, 1894 シイノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 119, ♀ (Miyanoshita).
Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan, China.

X. festivus EICHHOFF, 1875 オキノキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 202, ♀ (Japan).
Distr.: Japan (Honshu, Shikoku), Ryukyu Is.

X. formicatus EICHHOFF, 1868 ナンヨウキクイムシ

EICHHOFF, 1868, Berl. ent. Z., **12**, p. 151, ♀ (Ceylon).
Distr.: Japan (Bonin Is.), Taiwan, Philippines, Indochina, Malaya, Sumatra, Java,
New Guinea, Hawaii, Fiji, Sandwich Is., Ceylon, India.

X. fulvus MURAYAMA, 1936 ケヤキノキクイムシ

MURAYAMA, 1936, Tenthredo, **1**, p. 132, ♀ (Hōkidaisen).
Distr.: Japan (Honshu, Shikoku, Kyushu).

X. galeatus BLANDFORD, 1894 ヘルメットキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 133, ♂ (Nagasaki).
Distr.: Japan (Kyushu).

X. ganshoensis MURAYAMA, 1952 ガンショキクイムシ

MURAYAMA, 1952, Bull. Fac. Agr. Yamaguti Univ., **3**, p. 16 (Ganshō).
Distr.: Japan (Honshu, Kyushu).

X. glabratrus EICHHOFF, 1877 ハギキクイムシ (ヤマモモノキクイムシ)

EICHHOFF, 1877, Dtsch. ent. Z., **21**, p. 127, ♂, ♀ (Hagi).
Distr.: Japan (Honshu, Shikoku, Kyushu), Bonin Is., Taiwan, Assam, Bengal.

X. interjectus BLANDFORD, 1984 アイノキクイムシ

- BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 576, ♀ (Japan, China, Chusan Is.).
 Distr.: Japan (Kyushu), Taiwan, China, Vietnam, Indochina, Burma, Borneo, Sumatra, Java, Mentawai, Ceylon, Nepal, India.
- X. ishidai* NIJJIMA, 1909 イシダキクイムシ
 NIJJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 156, ♀ (Jozankei, Sapporo).
 Distr.: Japan (Hokkaido, Honshu).
- X. izuensis* MURAYAMA, 1952 イズキクイムシ
 MURAYAMA, 1952, Bull. Fac. Agr. Yamaguti Univ., **3**, p. 16, ♀ (Ganshô).
 Distr.: Japan (Honshu).
- X. kadoyamensis* MURAYAMA, 1934 カドヤマキクイムシ
 MURAYAMA, 1934, Annot. zool. Japon., **14**, p. 290, ♀ (Kadoyama).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Taiwan.
- X. kaimochii* NOBUCHI, 1981 カイモチザイノキクイムシ
 NOBUCHI, 1981, Kontyû, Tokyo, **49**, p. 143, ♀ (Takimoto).
 Distr.: Japan (Honshu).
- X. kirishimanus* MURAYAMA, 1955 キリシマキクイムシ
 MURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., **6**, p. 85, ♂ (Mt. Kirishima).
 Distr.: Japan (Honshu, Kyushu).
- X. kojimai* MURAYAMA, 1936 コジマキクイムシ
 MURAYAMA, 1936, Tenthredo, **1**, pp. 134, 143, ♀ (Uchinoura).
 Distr.: Japan (Shikoku, Kyushu).
- X. kraunhiae* NIJJIMA, 1910 フジノキクイムシ
 NIJJIMA, 1910, Trans. Sapporo nat. Hist. Soc., **3**, p. 14, ♀ (Kumanotaira).
 Distr.: Japan (Honshu).
- X. kumamotoensis* MURAYAMA, 1934 クマモトキクイムシ
 MURAYAMA, 1934, Annot. zool. Japon., **14**, p. 288, ♀, ♂ (Jisso).
 Distr.: Japan (Honshu, Shikoku, Kyushu).
- X. kumanoensis* NOBUCHI, n. nom. フタトゲザイノキクイムシ
 NOBUCHI, 1981, Kontyû, Tokyo, **49**, p. 147, ♀, ♂ (Kumano) (n. nam. for *Xyleborus bispinus* NOBUCHI, nec SCHEDL, 1978).
 Distr.: Japan (Honshu).
- X. laetus* NIJJIMA, 1909 ヨシカワキクイムシ
 NIJJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 159, ♂ (Sapporo).
 Distr.: Japan (Hokkaido, Honshu, Kyushu).
- X. lewisi* BLANDFORD, 1894 ルイスザイノキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 104, ♀ (Nikko, Hakone, Miyanoshita).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Taiwan, Borneo, Philippines, Indochina, Malaya, Burma, Sumatra, Java, India.
- X. longipilus* EGGLERS, 1926 ザイノケナガキクイムシ
 EGGLERS, 1926, Ent. Bl., **22**, p. 146 (Kumanotaira).
 Distr.: Japan (Honshu).
- X. machili* NIJJIMA, 1910 アオガシノキクイムシ

NIIJIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 14 (Tokio).
 Distr.: Japan (Honshu).

X. magnus NIIJIMA, 1910 ザイノオオキクイムシ

NIIJIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 11 (without locality).
 Distr.: Japan (Shikoku).

X. minutus BLANDFORD, 1894 ザイノコキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 116, ♀ (Inasa).
 Distr.: Japan (Honshu, Kyushu), Korea, Borneo, Malaya, Java.

X. misatoensis NOBUCHI, 1981 ミサトザイノキクイムシ

NOBUUHI, 1981, Kontyû, Tokyo, 49, p. 146, ♀ (Misato).
 Distr.: Japan (Kyushu).

X. miyazakiensis MURAYAMA, 1936 ミヤザキキクイムシ

MURAYAMA, 1936, Tenthredo, 1, p. 135, ♀ (Sakurakawa).
 Distr.: Japan (Kyushu).

X. montanus NIIJIMA, 1910 オオザイノキクイムシ

NIIJIMA, 1910, Trans. Sapporo nat. Hist. Soc., 3, p. 13, ♀ (Kumanotaira).
 Distr.: Japan (Honshu).

X. muticus BLANDFORD, 1894 カシワギキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 112, ♀ (Kashiwagi).
 Distr.: Japan (Honshu, Kyushu).

X. mutilatus BLANDFORD, 1894 クスノオオキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 103, ♀ (Japan).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Taiwan, Thailand,
 Malaya, Borneo, Sumatra, Java, Batoe.

X. nagaoensis MURAYAMA, 1934 ナガオキクイムシ

MURAYAMA, 1934, Annot. zool. Japon., 14, p. 294, ♂ (Tarumidzu).
 Distr.: Japan (Shikoku, Kyushu).

X. nameranus MURAYAMA, 1954 ナメラキクイムシ

MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., 5, p. 194, ♀ (Mt. Namera).
 Distr.: Japan (Honshu).

X. obliquecauda (MOTSCHEULSKY, 1863) タカネザイノキクイムシ

MOTSCHEULSKY, 1863, Bull. Soc. ent. Mosc., 36, p. 315, ♀ (*Phloeotrogus*) (Ceylon).
 Distr.: Japan (Honshu, Kyushu), Ceylon.

X. octiesdentatus MURAYAMA, 1930 ヒサカキノキクイムシ

MURAYAMA, 1930, J. Chosen nat. Hist. Soc., 15, p. 26 (without description).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Korea.

X. ohtoensis NOBUCHI, 1981 オオトザイノキクイムシ

NOBUCHI, 1981, Kontyû, Tokyo, 49, p. 149, ♀ (Mt. Ohto).
 Distr.: Japan (Honshu).

X. okinosenensis MURAYAMA, 1961 オキノセンキクイムシ

MURAYAMA, 1961, Akitu, 10, p. 31, ♀ (Mt. Okinosen, Kami-ishikawa).
 Distr.: Japan (Honshu).

X. onoharaensis MURAYAMA, 1934 オノハラキクイムシ

MURAYAMA, 1934, Annot. zool. Japon., 3, p. 293, ♀, ♂ (Tarumidzu, Onohara).

- Distr.: Japan (Shikoku, Kyushu).
- X. osumiensis** MURAYAMA, 1934 オオスミキクイムシ
 MURAYAMA, 1934, Annot. zool. Japon., **14**, p. 292, ♀ (Nagao).
 Distr.: Japan (Kyushu).
- X. pelliculosus** EICHHOFF, 1878 ヴラジロカシノキクイムシ
 EICHHOFF, 1878, Mém. Soc. roy. Sci. Liége (2) **8**, p. 339, ♀ (Hagi).
 Distr.: Japan (Honshu, Kyushu).
- X. perforans** (WOLLASTON, 1857) フィリピンザイノキクイムシ
 WOLLASTON, 1857, Cat. Coleopt. Madeira, p. 96, ♀ (*Tomicus*) (Madeira).
 Distr.: Bonin Is., Ryukyu Is. Tropicopolitan.
- X. permarginatus** SCHEDL, 1933 ミズサワキクイムシ
 SCHEDL, 1933, Philip. J. Sci., **52**, p. 200, ♀ (Mt. Maquiling).
 Distr.: Ryukyu Is., (Amami-oshima), Philippines.
- X. pfeili** (RATZEBURG, 1837) ファイルキクイムシ
 RATZEBURG, 1837, Die Forstinsenketen, **1**, p. 168, ♂, ♀ (*Bostrichus*) (Europe).
 Distr.: Japan (Hokkaido, Honshu, Kyushu), Siberia, Korea, Europe.
- X. praevius** BLANDFORD, 1894 センダツキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 110, ♀ (Japan).
 Distr.: Japan (Honshu, Kyushu), Korea.
- X. quercicola** EGGLERS, 1926 カシワザイノキクイムシ
 EGGLERS, 1926, Ent. B1., **22**, p. 146, ♀ (Tokio).
 Distr.: Japan (Honshu).
- X. rubricollis** EICHHOFF, 1875 アカクビキクイムシ
 EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 202, ♀ (*Bostrichus*) (Japan).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Bonin Is., Korea, Taiwan, Malaya, N. America.
- X. saxeseni** (RATZEBURG, 1937) サクセスキクイムシ
 RATZEBURG, 1937, Die Forstinsenketen, **1**, p. 167, ♀ (*Bostrichus*) (Europe).
sobrinus EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 202, ♀.
alni NIIZIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., **3**, p. 160, ♀ (Sapporo, Tomakomai).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Bonin Is., Kuril Is., Sakhalin, Kamchatka, Siberia, Korea, China, Taiwan, Vietnam, Thailand, Europe, Australia, America.
- X. schaufussi** BLANDFORD, 1894 シャウフスキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 117, ♀, ♂ (Junsai, Nikko).
 Distr.: Japan (Hokkaido, Honshu, Shikoku), Taiwan.
- X. seiryorensis** MURAYAMA, 1930 セイリョウリキクイムシ
 MURAYAMA, 1930, J. Chosen nat. Hist. Soc., **11**, p. 25, ♀ (Seiryori).
 Distr.: Japan (Honshu, Shikoku, Kyushu), Korea.
- X. seriatus** BLANDFORD, 1894 ハンノスジキクイムシ
 BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 111, ♀ (Nikko, Miyanoshita).
todo KÔNO, 1938, Ins. Matsum., **12**, p. 71, ♀ (Hokkaido).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Taiwan.

X. shionomisakiensis MURAYAMA, 1951 シオノミサキキクイムシ

MURAYAMA, 1951, Bull. Fac. Agr. Yamaguti Univ., **2**, p. 3, ♀, ♂ (Shiono-misaki).
Distr.: Japan (Honshu, Shikoku).

X. similis FERRARI, 1867

FERRARI, 1867, Die forst-und baumzuchtschädlichen Borkenkäfer, **1867**, p. 23, ♀.
Distr.: Bonin Is., Taiwan, Philippines, Borneo, Indochina, Malaya, Sumatra,
Mentawai, Java, Celebes, Micronesia, New Guinea, Australia, Solomon Is.,
Samoa, Tahiti, Seychellus, India, Ceylon, Madagascar, Africa.

X. takinoyensis MURAYAMA, 1953 タキノヤキキクイムシ

MURAYAMA, 1953, Report of the pine bark-beetle control investigation society, p.
109, ♂ (Takinoya).
Distr.: Japan (Honshu, Shikoku, Kyushu).

X. tsukubanus MURAYAMA, 1954 ツクバネヤマキクイムシ

MURAYAMA, 1954, Bull. Fac. Agr. Yamaguti Univ., **5**, pp. 184, 195, ♀ (Mt. Tsukuba).
Distr.: Japan (Honshu).

X. validus EICHHOFF, 1875 トドマツオオキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 202, ♀ (Japon).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Bonin Is., Ryukyu Is.,
Korea, China, Taiwan, Vietnam, Burma, Malaya, India, Ceylon.

X. volvulus (FABRICIUS, 1775) ユズリハノキクイムシ

FABRICIUS, 1775, Syst. Ent. p. 454, ♀ (*Bostrius*) (Cuba).
badius EICHHOFF, 1868, Berl. ent. Z., **12**, p. 280, ♀ (Madagascar, St. mauritius).
vicarius EICHHOFF, 1877, Ann. Soc. ent. Belg., **18**, p. 202, ♂, ♀ (Japon).
Distr.: Japan (Honshu, Shikoku, Kyushu), Ryukyu Is. Tropicopolitan.

Genus *Arixyleborus* HOPKINS, 1915 スジザイノキクイムシ属

Arixyleborus HOPKINS, 1915, U. S. Dept. Agr., Off. Sec., Rep. **99**, p. 59 (type-species: *Arixyleborus rugosipes* HOPKINS, 1915).

A. yakushimanus MURAYAMA, 1955 ヤクシマスジザイノキクイムシ

MURAYAMA, 1955, Bull. Fac. Agr. Yamaguti Univ., **6**, p. 83, ♀ (Miyanoura).
Distr.: Japan (Yakushima).

Genus *Cnestus* SAMPSON, 1911 ウスイロキクイムシ属

Cnestus SAMPSON, 1911, Ann. Mag. nat. Hist., (8) **7**, p. 383 (type-species: *Cnestus magnus* SAMPSON, 1911).

Tosaxyleborus MURAYAMA, 1950, Trans. Shikoku ent. Soc., **1**, p. 49 (type-species: *Tosaxyleborus pallidipennis* MURAYAMA, 1950).

C. murayamai SCHEDL, 1962 ウスキイロキクイムシ

SCHEDL, 1962, Ent. B1., **58**, p. 207.
pallidipennis MURAYAMA, 1950, Trans. Shikoku ent. Soc., **1**, p. 49, ♀ (*Tosaxyleborus*)
(nec EGGLERS, 1940) (Isa).
Distr.: Japan (Honshu, Shikoku, Kyushu), Ryukyu Is., Taiwan.

Tribe PITYOPHTHORINI ヒメキクイムシ族

Genus *Eidophelus* EICHHOFF, 1875 ヒメマルキクイムシ属

Eidophelus EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 200 (type-species: *Eidophelus imitans* EICHHOFF, 1875).

E. imitans EICHHOFF, 1875 クヌギノキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 200 (Japan).

Distr.: Japan (Kyushu), Korea.

E. minutus BLANDFORD, 1894 チュウゼンジキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 88. (Chuzenji).

Distr.: Japan (Honshu).

Genus *Pityophthorus* EICHHOFF, 1864 ヒメキクイムシ属

Pityophthorus EICHHOFF, 1864, Berl. ent. Z., **8**, p. 39 (type-species: *Bostrichus lichtensteini* RATZEBURG, 1837).

Myeloborus BLACKMAN, 1928, New York St. Coll. For., Tech. Bull., **25**, p. 16 (type-species: *Pityophthorus ramiperda* SWAINE, 1917).

Cladoborus SAWAMOTO, 1942, Ins. Matsum., **16**, p. 162 (type-species: *Cladoborus arakii* SAWAMOTO, 1942).

P. arakii SAWAMOTO, 1942 アラキサエダキクイムシ

SAWAMOTO, 1942, Ins. Matsum., **16**, p. 167 (Kutchan).

Distr.: Japan (Hokkaido).

P. jucundus BLANDFORD, 1894 トウヒノヒメキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, **1894**, p. 87 (Nagasaki).

Distr.: Japan (Honshu, Shikoku, Kyushu), Sakhalin, Siberia, Korea, NE. China.

Tribe IPINI イビニ族

Genus *Pityogenes* BEDEL, 1888 ホシガタキクイムシ属

Pityogenes BEDEL, 1888, Faune des Coléoptères du basin de la Seine, **6**, pp. 397, 418 (type-species: *Dermestes chalcographus* LINNÉ, 1761).

P. bistridentatus (EICHHOFF, 1879) フタツノキクイムシ

EICHHOFF, 1879, Mém. Soc. roy. Sci. Liège (2) **8**, p. 282 (var. of *Tomicus bidentatus* HERBST, 1783) (Europe).

Distr.: Japan (Hokkaido, Honshu), NE. China, Europe.

P. chalcographus (LINNÉ, 1761) ホシガタキクイムシ

LINNÉ, 1761, Fauna Suecica, **2**, p. 143 (*Dermestes*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Kuril Is., Sakhalin, Siberia, Korea, China, Europe.

P. foveolatus EGGERS, 1926 オンタケキクイムシ

EGGERS, 1926, Ent. Bl., **22**, p. 137 (Ontake, Etorof).

Distr.: Japan (Hokkaido, Honshu), Kuril Is., Sakhalin, Kamtchatka, Siberia.

P. japonicus NOBUCHI, 1974 ニホンホシガタキクイムシ

NOBUCHI, 1974, Bull. Gov. For. Exp. Sta., (266), p. 39 (Mt. Hakoda).

Distr.: Japan (Honshu).

P. seirindensis MURAYAMA, 1929 セイリンドウキクイムシ

MURAYAMA, 1929, J. Chosen nat. Hist. Soc., **9**, pp. 26, 30 (Seirindo).

aizawai KÔNO, 1938, Ins. Matsum., **12**, p. 69 (Horo, Rikubetsu).

nitidus EGGERS, 1941, Stettin. ent. Ztg., **102**, p. 121 (Ussuri).

Distr.: Japan (Hokkaido), Kuril Is., Sakhalin, Siberia, Korea.

Genus *Pityokteines* FUCHS, 1911 キヨクシキクイムシ属

Pityokteines FUCHS, 1911, Morphologische Studien über Borkenkäfer, p. 37 (type-species: *Bostichus curvidens* GERMAN, 1824).

P. curvidens (GERMAN, 1824) キヨクシキクイムシ

GERMAN, 1824, Insectorum species novae aut minus congnitae, descriptionibus illustratae, p. 462 (*Bostichus*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Europe.

Genus *Orthotomicus* FARRARI, 1867 ツノキクイムシ属

Orthotomicus FARRARI, 1867, Die forst- und baumzuchtschädlichen Borkenkäfer, p. 44 (type-species: *Bostichus laricis* FABRICIUS, 1792).

O. angulatus (EICHHOFF, 1875) マツノツノキクイムシ

EICHHOFF, 1875, Ann. Soc. ent. Belg., **18**, p. 200 (*Tomicus (Cyrtotomicus)*) (Japon).

Distr.: Japan (Honshu, Shikoku, Kyushu), Ryukyu Is., Taiwan, Korea, China, Cambodia.

O. golovjankoi PJATNITZKY, 1930 ゴロウヤンコキクイムシ

PJATNITZKY, 1930, Ent. Bl., **26**, p. 179 (Ostsibirien).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, NE. China.

O. kuniyoshii (NOBUCHI, 1959) クニヨシキクイムシ

NOBUCHI, 1959, Bull. Gov. For. Exp. Sta., (185), p. 23 (*Ips*) (Kitanakagashuku).

Distr.: Ryukyu Is. (Okinawa).

O. laricis (FABRICIUS, 1792) カラマツキクイムシ

FABRICIUS, 1792, Ent. syst., **1**, p. 365 (*Bostichus*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Sakhalin, Siberia, Kamtchatka, Korea, China, Europe.

O. proximus (EICHHOFF, 1867) マツカワノキクイムシ

EICHHOFF, 1867, Berl. ent. Z., **11**, p. 403 (*Tomicus*) (Europe).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Siberia, Korea, China, Europe.

O. suturalis (GYLLENHAL, 1827) ホンスンキクイムシ

GYLLENHAL, 1827, Insecta svec. **4**, p. 622 (*Bostichus*) (Europe).

Distr.: Japan (Hokkaido, Honshu), Siberia, Korea, China, Europe.

O. tosaensis (MURAYAMA, 1950) トサキクイムシ

MURAYAMA, 1950, Trans. Shikoku ent. Soc., **1**, p. 52 (*Ips*) (Irinohama, Unzen).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus *Ips* DEGEER, 1775 ハキクイムシ属

Ips DEGEER, 1775, Mémoires pour servir à l'histoire des insectes **5**, p. 190 (type-species:

Dermestes typographus LINNÉ, 1758).

I. acuminatus (GYLLENHAL, 1827) マツノムツバキクイムシGYLLENHAL, 1827, Insecta svec. 4, p. 620 (*Bostrichus*) (Europe).

Distr.: Japan (Hokkaido, Honshu, Shikoku), Sakhalin, Siberia, Korea, China, Thailand, Europe.

I. cembrae (HEER, 1836) カラマツヤツバキクイムシ (マツノオオキクイムシ, カラマツオオキクイムシ)HEER, 1836, Observations entomologica, p. 28 (*Bostrichus*) (Europe).*subelongatus* MOTSCHULSKY, 1860, Schrenks Reise, 2, p. 155 (*Bostrichus*).

Distr.: Japan (Hokkaido, Honshu), Kuril Is., Sakhalin, Siberia, Korea, NE. China, Taiwan, Europe.

I. duplicatus (SAHLBERG, 1836) オオハキクイムシSAHLBERG, 1836, Dissertatio entomological insecta Fennica enumerans, 1, p. 144 (*Bostrichus*) (Europe).

Distr.: Japan (Honshu), Sakhalin, Siberia, NE. China, Europe.

I. multidentatus MURAYAMA, 1953 マツノトゲキクイムシ

MURAYAMA, 1953, Bull. Fac. Agr. Yamaguti Univ., 4, p. 35 (Mt. Jakuchi).

Distr.: Japan (Honshu, Shikoku, Kyushu).

I. typographus LINNÉ, 1758 ヤツバキクイムシLINNÉ, 1758, Syst. Nat., ed. 10, p. 335 (*Dermestes*) (Europe).Distr.: Japan, Kuril Is., Sakhalin, Kamchatka, Siberia, Korea, NE. China, Europe.
subsp. *japonicus* NIJIMA, 1909, J. Coll. Agr. Tohoku imp. Univ., 3, p. 147 ((Tomas-
komai, Iburī, Jozankei, Nopporo, Uriu)).

Distr.: Japan (Hokkaido, Honshu), Sakhalin.

Genus *Acanthotomicus* BLANDFORD, 1894 ナガハキクイムシ属*Acanthotomicus* BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 89 (type-species:
Acanthotomicus spinosus BLANDFORD, 1894).*A. spinosus* BLANDFORD, 1894 シラカシノキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 90 (Oyama, Nikko).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Subfamily SCOLYTOPLATYPINAE キザハシキクイムシ亜科

Genus *Scolytoplatypus* SCHAUFUSS, 1861 キザハシキクイムシ属*Scolytoplatypus* SCHAUFUSS, 1861, Tijd. Ent., 34, p. 31 (type-species: *Scolytoplatypus permirus* SCHAUFUSS, 1861).*S. daimio* BLANDFORD, 1893 ダイミョウキクイムシ

BLANDFORD, 1893, Trans. ent. Soc. London, 1893, p. 433 (Nikko).

siomio BLANDFORD, 1893, Trans. ent. Soc. London, 1893, p. 434 (Nikko, Sendai).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Kuril Is.

S. mikado BLANDFORD, 1893 ミカドキクイムシ

BLANDFORD, 1893, Trans. ent. Soc. London, 1893, p. 437 (Nikko, Oyama, Sapporo).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Ryukyu Is., Sakhalin, Korea, Taiwan, Malaya, India.

S. shogun BLANDFORD, 1894 ショウグンキクイムシ

BLANDFORD, 1894, Trans. ent. Soc. London, 1894, p. 126 (Higo).

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 31

December 29, 1989

The Japanese Society of Coleopterology
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family CHRYSOMELIDAE ハムシ科
(CHRYSOMELINAE ハムシ亜科)

By

Shinsaku KIMOTO

Subfamily CHRYSOMELINAE ハムシ亜科

Genus *Chrysolina* MOTSCHULSKY, 1860 ヨモギハムシ属

Chrysolina MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 210 (type species: *Chrysomela staphylea* LINNAEUS, 1758).—SILFVERBERG, 1980, Bull. zool. Nomencl., 31, p. 57 (placed in Official List of Generic Names in Zoology).

Atechna CHEVROLAT, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 427.—SILFVERBERG, 1980, Bull. zool. Nomencl., 27, p. 57 (type designated as *Chrysomela vulpina* FABRICIUS, 1781; requested not to be given priority over *Chrysolina* MOTSCHULSKY).

Polysticta HOPE, 1840, Coleopt. Man., Mand., 3, p. 164 (nec *Polysticta* EYTON, 1836; Aves).

Rhyssoloma WOLLASTON, 1854, Ins. Mader., p. 458.

Apterosoma MOTSCHULSKY, 1860, Étud. ent., 9, p. 23.

Anopachys MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 202.

Bittotaenia MOTSCHULSKY, 1860, ibid., p. 206.

Centoptera MOTSCHULSKY, 1860, ibid., p. 207.

Chalcoidea MOTSCHULSKY, 1860, ibid., p. 209.

Chrysomorph MOTSCHULSKY, 1860, ibid., p. 204.

Colaphodes MOTSCHULSKY, 1860, ibid., p. 212.

Colaphoptera MOTSCHULSKY, 1860, ibid., p. 215.

Colaphosoma MOTSCHULSKY, 1860, ibid., p. 216.

Craspeda MOTSCHULSKY, 1860, ibid., p. 191.

Dlochrysa MOTSCHULSKY, 1860, ibid., p. 203.

Distr.: Japan (Hokkaido, Honshu, Kyushu), Taiwan.

S. tycon BLANDFORD, 1893 タイコンキクイムシ

BLANDFORD, 1893, Trans. ent. Soc. London, 1893, p. 432 (Nikko, Kiga).

ussuriensis BERGER et CHOLODOKOVSKY, 1916, Rev. russ. Ent., 16, p. 1 (Vladivostok).

Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu), Kuril Is., Sakhalin, Siberia,
Korea, Taiwan.