

CHECK-LIST
OF
COLEOPTERA OF JAPAN

日本産甲虫目録

No. 31

December 29, 1989

The Japanese Society of Coleopterology
c/o Department of Zoology, National Science Museum (Nat. Hist.)
3-23-1 Hyakunin-chô, Shinjuku, Tokyo 160, Japan

Family CHRYSOMELIDAE ハムシ科
(CHRYSOMELINAE ハムシ亜科)

By

Shinsaku KIMOTO

Subfamily CHRYSOMELINAE ハムシ亜科

Genus *Chrysolina* MOTSCHULSKY, 1860 ヨモギハムシ属

Chrysolina MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 210 (type species: *Chrysomela staphylea* LINNAEUS, 1758).—SILFVERBERG, 1980, Bull. zool. Nomencl., 31, p. 57 (placed in Official List of Generic Names in Zoology).

Atechna CHEVROLAT, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 427.—SILFVERBERG, 1980, Bull. zool. Nomencl., 27, p. 57 (type designated as *Chrysomela vulpina* FABRICIUS, 1781; requested not to be given priority over *Chrysolina* MOTSCHULSKY).

Polysticta HOPE, 1840, Coleopt. Man., Mand., 3, p. 164 (nec *Polysticta* EYTON, 1836; Aves).

Rhyssoloma WOLLASTON, 1854, Ins. Mader., p. 458.

Apterosomalia MOTSCHULSKY, 1860, Étud. ent., 9, p. 23.

Anopachys MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 202.

Bittotaenia MOTSCHULSKY, 1860, ibid., p. 206.

Centoptera MOTSCHULSKY, 1860, ibid., p. 207.

Chalcoidea MOTSCHULSKY, 1860, ibid., p. 209.

Chrysomorph MOTSCHULSKY, 1860, ibid., p. 204.

Colaphodes MOTSCHULSKY, 1860, ibid., p. 212.

Colaphoptera MOTSCHULSKY, 1860, ibid., p. 215.

Colaphosoma MOTSCHULSKY, 1860, ibid., p. 216.

Craspeda MOTSCHULSKY, 1860, ibid., p. 191.

Dlochrysa MOTSCHULSKY, 1860, ibid., p. 203.

- Heliolestola* MOTSCHULSKY, 1860, *ibid.*, p. 190.
Hoplasoma MOTSCHULSKY, 1860, *ibid.*, p. 211.
Lithoptera MOTSCHULSKY, 1860, *ibid.*, p. 210.
Ovomorpha MOTSCHULSKY, 1860, *ibid.*, p. 213.
Ovosoma MOTSCHULSKY, 1860, *ibid.*, p. 214.
Ovostoma MOTSCHULSKY, 1860, *ibid.*, p. 215.
Pelurosticha MOTSCHULSKY, 1860, *ibid.*, p. 191.
Stichoptera MOTSCHULSKY, 1860, *ibid.*, p. 209.
Stichosoma MOTSCHULSKY, 1860, *ibid.*, p. 208.
Taeniosticha MOTSCHULSKY, 1860, *ibid.*, p. 207.
Timarchoptera MOTSCHULSKY, 1860, *ibid.*, p. 118.
Threnosoma MOTSCHULSKY, 1860, *ibid.*, p. 213.
Zeugotaenia MOTSCHULSKY, 1860, *ibid.*, p. 206.
Crositops MARSEUL, 1883, *Abeille*, Paris, **21**, p. 105.
Hypericia BEDEL, 1892, *Faune Coleopt. Bassin Seine*, **5**, p. 258 (as a division of *Chrysomela*).
Sphaeromela BEDEL, 1892, *ibid.*, p. 260, nota (as a division of *Chrysomela*).
Pezocrosita JACOBSON, 1901, *Öfvers. Finska Vetensk Soc. Förh.*, **43**, p. 120 (as a subgenus of *Crosita*).
Timarchomela ACHARD, 1922, *Fragm. ent.*, **1922**, p. 17.
Minckia STRAND, 1935, *Folia zool. hydrobiol.*, **7**, p. 292 (n. n. for *Hoplasoma* MOTSCHULSKY, nec AGASSIZ, 1846).
Lithopteroides STRAND, 1935, *ibid.*, p. 295 (n. n. for *Lithoptera* MOTSCHULSKY, nec MÜLLER, 1858).
Naluhia BECHYNÉ, 1948, *Rev. suisse zool.*, **55**, p. 540 (as a subgenus of *Chrysolina*).
Camerounia JOLIVET, 1949, *Bull. Inst. Sci. nat. Belg.*, **15** (40), p. 7 (as a subgenus of *Iscadida*).
Allochrysolina BECHYNÉ, 1950, *Ent. Arb. Mus. Frey*, **1**, p. 133 (as a subgenus of *Chrysolina*).
Allohypericia BECHYNÉ, 1950, *ibid.*, p. 159 (as a subgenus of *Chrysolina*).
Caudatochrysa BECHYNÉ, 1950, *ibid.*, p. 149 (as a subgenus of *Chrysolina*).
Chrysocrosita BECHYNÉ, 1950, *ibid.*, p. 90 (as a subgenus of *Chrysolina*).
Chrysolinopsis BECHYNÉ, 1950, *ibid.*, p. 82 (as a subgenus of *Chrysolina*).
Erythrochrysa BECHYNÉ, 1950, *ibid.*, p. 91 (as a subgenus of *Chrysolina*).
Euchrysolina BECHYNÉ, 1950, *ibid.*, p. 83 (as a subgenus of *Chrysolina*).
Ghesquiereita BECHYNÉ, 1950, *ibid.*, p. 171 (as a subgenus of *Chrysolina*).
Maenadochrysa BECHYNÉ, 1950, *ibid.*, p. 116 (as a subgenus of *Chrysolina*).
Melasomoptera BECHYNÉ, 1950, *ibid.*, p. 141 (as a subgenus of *Chrysolina*).
Menthastrilla BECHYNÉ, 1950, *ibid.*, p. 74 (as a subgenus of *Chrysolina*).
Parkaniola BECHYNÉ, 1950, *ibid.*, p. 130 (as a subgenus of *Chrysolina*).
Pierryvettia BECHYNÉ, 1950, *ibid.*, p. 68 (as a subgenus of *Chrysolina*).
Taeniochrysea BECHYNÉ, 1950, *ibid.*, p. 87 (as a subgenus of *Chrysolina*).
Timarcholina BECHYNÉ, 1950, *ibid.*, p. 66 (as a subgenus of *Chrysolina*).
Timarchomima BECHYNÉ, 1950, *ibid.*, p. 65 (as a subgenus of *Chrysolina*).
Palaesticta BECHYNÉ, 1952, *ibid.*, **3**, p. 383 (as a subgenus of *Chrysolina*).
Polystictella BECHYNÉ, 1952, *ibid.*, **3**, p. 384 (as a subgenus of *Chrysolina*).
Cobosorina DAVID, 1953, *Rev. fr. Ent.*, **20**, p. 186 (as a subgenus of *Chrysochloa*).

- Diachalcoidea* BECHYNÉ, 1955, Bull. Res. Counc. Israel, **4**, p. 349 (as a subgenus of *Chrysolina*).
Arctolina KONTKANEN, 1957, Ann. ent. fenn., Helsinki, **25**, p. 31 (as a subgenus of *Chrysolina*).
Oreina: KIMOTO, 1964, J. Fac. Agr. Kyushu Univ., **13**, p. 264.
Bechynea MEDVEDEV, 1966, Forest Ent.-Fauna Kuril, p. 40 (as a subgenus of *Chrysolina*).
Jacobsonia MEDVEDEV, 1970, Trudy Biol.-Pochvenn. Ins., Vladivostok, **2**, p. 162 (as a subgenus of *Chrysolina*).
Pseudololithoptera MEDVEDEV, 1970, ibid., p. 159 (as a subgenus of *Chrysolina*).
Bechynia BOURDONNE, 1977, Nouv. Rev. Ent., **7**, p. 330 (as a subgenus of *Chrysolina*).
Vittatochrysa LOPATIN, 1977, Chrysom. Middle Asia and Kazacstan, p. 146 (as a subgenus of *Chrysolina*).
Paradiochaloidea DACCORDI, 1978, Rev. Zool. afr., **92**, p. 752 (as a subgenus of *Chrysolina*).
Pseudotaeniochrysea DACCORDI, 1980, ibid., **94**, p. 302 (as a subgenus of *Chrysolina*).
Paramenthastriella DACCORDI, 1980, ibid., p. 304 (as a subgenus of *Chrysolina*).
Pseudotimarchomima DACCORDI, 1980, ibid., p. 307 (as a subgenus of *Chrysolina*).
C. aino TAKIZAWA, 1970 アイヌヨモギハムシ
TAKIZAWA, 1970, Kontyû, Tokyo, **38**, p. 117 (Hokkaido).
Distr.: Japan (Hokkaido).
C. angusticollis (MOTSCHULSKY, 1860) オオヨモギハムシ
MOTSCHULSKY, 1860, Étud. ent., **9**, p. 23 (*Apterosoma*) (Japan).
japana BALY, 1874, Trans. ent. Soc. London, **1874**, p. 171 (*Chrysomela*) (Hakodate; Manchuria).
Distr.: Japan (Hokkaido), E. Siberia, NE. China.
C. aurichalcea (MANNERHEIM, 1825) ヨモギハムシ
MANNERHEIM, 1825, in HUMMEL, Essais Ent., **4**, p. 39 (*Chrysomela*) (Altai).
asclepiadis VILLA, 1833, Coleopt. eur. dupl., p. 36 (*Chrysomela*) (no locality cited).
villae CRISTOFORI, 1833, Conspr. Meth., p. 82 (*Chrysomela*) (It. bor.).
gibbipennis FALDERMANN, 1835, Mém. Acad. St.-Pétersb., **2**, p. 441 (*Chrysomela*) (Irkutzk).
elevata SUFFRIAN, 1851, Linn. Ent., **5**, p. 189 (*Chrysomela*) ("?London").
stali BALY, 1860, J. Ent., **1**, p. 95 (*Chrysomela*) (N. China).
quadrangulata MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 226, pl. 11, fig. 10 (*Chrysomorpha*) (Amur).
violaceicollis MOTSCHULSKY, 1861, Étud. ent., **10**, p. 21 (*Anopachys*) (Japan).
wallacei BALY, 1862, Annls. Mag. nat. Hist., (3), **10**, p. 21 (*Chrysomela*) (Manchuria).
cyanopurpurea BALLION, 1878, Bull. Soc. Natur. Moscow, **53**, p. 379 (*Chrysomela*) (Kuldsha).
aurichalcea eurina FRIVALDSZKY, 1883, Természetr. Füz., **7**, p. 17 (*Chrysomela*) (Mehadia).
amethystina KOLBE, 1886, Arch. Naturg., **52** (1), p. 228 (*Chrysomela*) (Korea).
cupararia KOLBE, 1886, ibid., p. 228 (*Chrysomela*) (Korea).
aurichalcea var. *japana* MARSEUL, 1886, Abeille, Paris, **24**, p. 41 (*Chrysomela*) (nec BALY, 1874; Japan).

- pekinensis* FAIRMAIRE, 1887, Rev. Ent., 6, p. 331 (*Chrysomela*) (China).
aurichalcea var. *lineella* WEISE, 1887, Arch. Naturg., 53 (1), p. 182 (*Chrysomela*) (Amur).
aurichalcea var. *recticollis* WEISE, 1887, ibid., pp. 182, 185 (*Chrysomela*) (Japan).
aurichalcea var. *nigricans* JACOBSON, 1900, Horae Soc. ent. ross., 35, p. 100 (*Chrysomela*) (Siberia).
aurichalcea var. *collaris* WEISE, 1916, Coleopt. Cat. Junk, 68, p. 56 (*Chrysomela*) (n. n. for *aurichalcea* var. *recticollis* WEISE).
ascepiadis schatzmayri MÜLLER, 1916, Ent. Bal., 12, p. 96, figs. (*Chrysomela*) (Istria).
ascepiadis thurntaxisi SCHATZMAYR, 1927, Boll. Soc. Adriat. Sci. Trieste, 29, p. 151 (*Chrysomela*) (Istria).
ascepiadis bohemica MÜLLER, 1948, Atti Mus. Civ. Stor. nat. Trieste, 17, p. 95 (*Chrysomela*) (C. Europe).
aurichalcea kwanghsiensis BECHNYÉ, 1950, Ent. Arb. Mus. Frey, 1, p. 147 (China).
aurichalcea omisiensis BECHYNÉ, 1950, ibid., p. 147 (China).
aurichalcea fokiensis BECHYNÉ, 1950, ibid., p. 147 (China).
aurichalcea yunnanica BECHYNÉ, 1950, ibid., p. 148 (China, Laos).
aurichalcea vagesplendens BECHYNÉ, 1950, ibid., p. 148 (Annam).
aurichalcea problematica KASZAB, 1960, Rovart. Kozl., 15 (3), p. 55 (*Chrysomela*) (Hungary).
Distr.: Japan (Hokkaido, Honshu, Hachijo, Sado, Awa-shima, Shikoku, Kyushu, Goto, Tsushima, Yakushima), Ryukyu Is. (Tokara, Amami-Oshima, Okierabu, Okinawa), Taiwan, Vietnam, Laos, C. Europe, Siberia, Mongolia, China, Korea.

C. brunneipennis (MATSUMURA, 1911) オドリコソウハムシ

MATSUMURA, 1911, J. Agr. Tohoku imp. Univ., Sapporo, 4, p. 149 (*Chrysomela*) (S. Sachalin).

Distr.: Japan (Hokkaido), S. Kuriles (Kunashiri), Sachalin.

subsp. *lamii* TAKIZAWA, 1970

TAKIZAWA, 1970, Kontyû, Tokyo, 38, p. 118 (as *Chrysolina lamii*) (Hokkaido, Kurile Is.).

Distr.: Japan (Hokkaido), S. Kuriles (Kunashiri).

C. exanthematica (WIEDEMANN, 1821) ハッカハムシ

WIEDEMANN, 1821, Germar's Mag. Ent., 4, p. 178 (*Chrysomela*) (Siberia).

guttata GEBLER, 1817, Mém. Acad. Mosc., 5, p. 316 (*Chrysomela*) (nec FABRICUS, 1792) (Siberia).

musiva GEBLER, 1830, Ledeb. Reise, 2, p. 215 (*Lithoptera*) (Altai).

speculifera REDTENBACHER, 1848, in HUGEL, Reise Kaschmir, 4, p. 558 (*Chrysomela*) (Caschmir).

subaenea MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 229, pl. 11, fig. 13 (*Lithoptera*) (nec SUFFRIAN, 1851) (Japan).

consimilis BALY, 1874, Trans. ent. Soc. London, 1874, p. 172 (*Chrysomela*) (nec CLARK, 1864) (n. n. for *subaenea* MOTSCHULSKY) (Nagasaki; E. Siberia, N. China).

laevipunctata LEWIS, 1879, Cat. Coleopt. Japanese Archipelago, p. 27 (*Chrysomela*) (n. n. for *subaenea* MOTSCHULSKY, and *consimilis* BALY) (Japan).

guttata var. *marseuli* WEISE, 1898, Arch. Naturg., 64 (1), p. 200 (*Chrysomela*) (no

- locality cited).
- Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima), Siberia, Mongolia, Korea, China, Taiwan, N. India.
- C. nikkoensis* (JACOBY, 1885) ニッコウルリハムシ
JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 207 (*Chrysomela*) (Nikko, Yunoshiku, Urasa).
Distr.: Japan (Honshu, Sado).
- C. nikolskyi* (JACOBSON, 1898) カワカミハムシ
JACOBSON, 1898, Ann. Mus. St.-Pétersb., **2**, p. 200 (*Chrysomela*) (E. Siberia).
kawakamii MATSUMURA, 1911, J. Coll. Agr. Tohoku imp. Univ., Sapporo, **4**, p. 141 (*Timarcha*) (Rishiri Is., S. Saghalien).
Distr.: Japan (Hokkaido), E. Siberia, Sachalin.
- C. pirka* TAKIZAWA, 1970 サメハダヨモギハムシ
TAKIZAWA, 1970, Kontyû, Tokyo, **38**, p. 118 (Hokkaido, Kurile Is.: Kunashiri).
Distr.: Japan (Hokkaido), S. Kuriles (Kunashiri).
- C. porosirensis* TAKIZAWA, 1970 ミヤマヨモギハムシ
TAKIZAWA, 1970, Kontyû, Tokyo, **38**, p. 120 (Hokkaido).
Distr.: Japan (Hokkaido).
- C. virgata* (MOTSCHULSKY, 1860) オオルリハムシ
MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 224, pl. 11, fig. 8 (*Dlochrysa*) (Amur).
obscurofasciata JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 208, pl. 11, fig. 11 (*Chrysomela*) (Niigata).
Distr.: Japan (Honshu, Sado), E. Siberia, China, Korea.
- C. watanabei* TAKIZAWA, 1970 ワタナベハムシ
TAKIZAWA, 1970, Kontyû, Tokyo, **38**, p. 121 (Hokkaido, Sachalin).
Distr.: Japan (Hokkaido), Sachalin.
- C. yezoensis* (MATSUMURA, 1911) クロルリハムシ
MATSUMURA, 1911, J. Arg. Tohoku imp. Univ., Sapporo, **4**, p. 149 (*Chrysomela*) (S. Sachalin, Japan).
geminata: JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 207 (*Chrysomela*) (Hakodate).
pseudogeminata BECHYNÉ, 1950, Ent. Arb. Mus. Frey, **1**, p. 156 (Kioto, Innai in N. Hondo).
shikokensis NAKANE, 1963, Fragm. coleopt., ed. NAKANE, (5), p. 19 (Mt. Tsurugi).
aeruginosa: KIMOTO, 1964, J. Fac. Agr. Kyushu Univ., **13**, p. 268 (Japan).
Distr.: Japan (Hokkaido, Honshu, Shikoku), Sachalin.

Genus *Agrosteomela* GISTEL, 1857 オオハムシ属

Agrosteomela GISTEL, 1857, Achthundert und zwanzig neue oder unbeschriebene wirbellose Thiere, **2**, p. 605 (type species: *Chrysomela indica* HOPE, 1831).

Paralina BALY, 1859, Trans. ent. Soc. London, (2), **5**, p. 155.

A. indica (HOPE, 1831) オオハムシ

HOPE, 1831, in GRAY, Misc. zool., p. 39 (*Chrysomela*) (Nepal).

caschmirensis REDTENBACHER, 1843, in HUGEL, Reise Kashmîr, **4**, p. 558 (*Chrysomela*) (Cashmir).

elata STÅL, 1857, Öfvers. Vetensk.-Akad. Förh., 14, p. 60 (*Lina*) (Thibet).
impressiuscula FAIRMAIRE, 1878, Annls. Soc. ent. France, (5), 8, p. 135 (*Paralina*) (C. China).
chinensis WEISE, 1922, Tijdschr. Ent., 65, p. 54 (*Paralina*) (Fukien).
Distr.: Japan (Honshu), N. India, Nepal, Bhutan, Vietnam, S. China, Taiwan.

Genus *Phaedon* LATREILLE, 1829 ダイコンハムシ属

Phaedon LATREILLE, 1829, in CUVIER, Règne Anim., ed. 2, 5, p. 151 (type species: *Chrysomela armoraciae* LINNAEUS, 1758).
Alitene GISTEL, 1857, Achthundert und zwanzig neue oder unbeschriebene wirbellose Thiere, 2, p. 530.
Orthosticha MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 196.
Emmetrus MOTSCHULSKY, 1860, ibid., p. 221.
Stenoplatys MOTSCHULSKY, 1860, ibid., p. 220.
Neophaedon JACOBSON, 1900, Horae Soc. ent. ross., 38, p. 90.
Hemiphaedon JACOBSON, 1901, Öfvers. Finska Vetensk Soc. Förh., 43, p. 131.
Paraphaedon SHARP, 1910, Ent. month. Mag., 46, p. 4.
Austrosternoplatus JACOBSON, 1921, Bull. Sta. Reg. Prot. Plantes Petrograd, 2, p. 24.
Allophaedon KONTKANEN, 1933, Annl. Soc. Zool-Bot. Fenn., 14, p. 70.

P. brassicae BALY, 1874 ダイコンハムシ

BALY, 1874, Trans. ent. Soc. London, 1874, p. 174 (Nagasaki).
incertum BALY, 1874, ibid., p. 175 (Nagasaki).
nigritus CHÛJÔ, 1951, Trans. Shikoku ent. Soc., 2, p. 38, fig. 2 (Mt. Saraga-mine in Ehime Pref.).
armoraciae: WEISE, 1898, Arch. Naturg., 64 (1), p. 203 (Japan).
Distr.: Japan (Honshu, Sado, Shikoku, Kyushu, Tsushima, Yakushima, Kuchi-erabu-jima), Vietnam, China, Taiwan, Korea.

Genus *Gastrophysa* CHEVROLAT, 1837 コガタルリハムシ属

Gastrophysa CHEVROLAT, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 429 (type species: *Chrysomela polygoni* LINNAEUS, 1758).
Gastroidea HOPE, 1840, Coleopt. Man., 3, p. 164.
Gastroidea GEMMINGER et HAROLD, 1874, Cat. Coleopt., 11, p. 3403.
Exiguipenna JOLIVET, 1951, Bull. Inst. Roy. Sci. nat. Belg., 27 (21), p. 3.

G. atrocyanea MOTSCHULSKY, 1860 コガタルリハムシ

MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 222, pl. 11, fig. 3 (E. Siberia).
tonkinea ACHARD, 1926, Fragm. ent., 1926, p. 134 (*Gastroidea*) (Tonkin).
Distr.: Japan (Honshu, Sado, Shikoku, Kyushu, Tsushima), E. Siberia, Korea, China, Taiwan, Vietnam.

Genus *Gastrolina* BALY, 1859 クルミハムシ属

Gastrolina BALY, 1859, Annls. Mag. nat. Hist., (3), 4, p. 61 (type species: *Gastrolina depressa* BALY, 1859).
Linastica MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 200.

G. depressa BALY, 1859 クルミハムシ

BALY, 1859, Annls. Mag. nat. Hist., (3), 4, p. 61 (N. China).
thoracica BALY, 1864, Trans. ent. Soc. London, (3), 2, p. 228 (Amur).

depressa pallipes CHEN, 1974, Acta ent. sin., **17**, p. 196 (W. Yunnan).

Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu), Siberia, China, Korea.

G. peltoidea (GEBLER, 1832) ミヤマヒラタハムシ

GEBLER, 1832, Nouv. Mém. Moscou, **2**, p. 74 (*Chrysomela*) (E. Siberia).

japonica JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 210, pl. 11, fig. 12 (Japan).

Distr.: Japan (Hokkaido, Honshu, Shikoku), E. Siberia, N. China.

Genus **Gastrolinoides** CHÛJÔ et KIMOTO, 1960 ズグロキハムシ属

Gastrolinoides CHÛJÔ et KIMOTO, 1960, Niponius, Takamatsu, **1** (4), p. 7 (type species:

Melasoma japonica HAROLD, 1877).

G. japonica (HAROLD, 1877) ズグロキハムシ

HAROLD, 1877, Dtsch. ent. Z., **21**, p. 362 (*Melasoma*) (Hagi).

Distr.: Japan (Honshu, Shikoku, Kyushu).

Genus **Plagiодера** CHEVROLAT, 1837 ヤナギルリハムシ属

Plagiодера CHEVROLAT, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 404 (type species: *Chrysomela versicolora* LAICHARTING, 1781).

Plagiosterna MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 196.

Linamorpha MOTSCHYLSKY, 1860, ibid., p. 197.

Plagiomorpha MOTSCHULSKY, 1860, ibid., p. 200.

Pseudolina SCHAEFFER, 1919, J. New York ent. Soc., **27**, p. 331.

Melasomida SCHAEFFER, 1920, Bull. Brooklyn ent. Soc., **15**, p. 117.

P. versicolora (LAICHARTING, 1781) ヤナギルリハムシ

LAICHARTING, 1781, Verz. Tirol. Ins., **1**, p. 148 (*Chrysomela*) (Tyrol).

armoraicae: FABRICIUS (nec LINNAEUS), 1775, Syst. Ent., p. 103 (*Chrysomela*) (no locality cited).

similis HERBST, 1783, in FUESSLY, Arch. Insecteng., **4**, p. 58 (*Chrysomela*) (Berlin).

salicis PANZER, 1795, Dtschl. Ins.-f., Nürnberg, p. 172 (*Galleruca*) (no locality cited).

distinct BALY, 1874, Trans. ent. Soc. London, **1874**, p. 174 (Nagasaki).

versicolora laevicollis LOPATIN, 1962, Annls. hist.-nat. Mus. natn. hung., **54**, p. 325 (Nordl. China).

Distr.: Japan (Hokkaido, Honshu, Awa-shima, Sado, Shikoku, Kyushu, Tsushima), Ryukyu Is. (Amami-Oshima), Siberia, Korea, China, Europe, N. Africa, Taiwan, India.

Genus **Chrysomela** LINNAEUS, 1758 ドロノキハムシ属

Chrysomela L., 1758, Syst. Nat. ed. 10, p. 368 (type species: *Chrysomela populi* LINNAEUS, 1758).

Melasoma STEPHENS, 1831, Illustr. Brit. Ent., Mabdib., **4**, p. 349.

Gymnota GISTEL, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 402.

Lina MEGERLE, 1837, in DEJEAN, ibid., p. 402.

Microdera STEPHENS, 1839, Man. Brit. Coleopt., p. 307.

Macrolina MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 198.

Strickcerus LUCAS, 1920, Catalog. Alphabeticus., p. 413.

Pachylina MEDVEDEV et CHERNOV, 1969, Zool. Zh., **48**, p. 536.

C. cuprea FABRICIUS, 1775 ムラサキハムシ

FABRICIUS, 1775, Syst. Ent., p. 96 (Germania).
ruficaudis DEGEER, 1775, Mém. Hist. Ins., 5, p. 305 (no locality cited).
humeralis ESCHSCHOLTZ, 1818, Mém. Acad. St.-Pétersb., 6, p. 465 (Livoniae).
cuprea var. *sarmatica* WEISE, 1887, Arch. Naturg., 53, p. 187, nota (*Melasoma*) (Minsk).
Distr.: Japan (Hokkaido), Siberia, Mongolia, Europe.

C. lapponica LINNAEUS, 1758 ウズマキハムシ

LINNAEUS, 1758, Syst. Nat., ed. 10, p. 371 (Europe).
curvilinea DEGEER, 1775, Mém. Hist. Ins., 5, p. 302 (no locality cited).
bulgharensis FABRICIUS, 1798, Suppl. Ent. Syst., p. 88 (Germania).
interrupta FABRICIUS, 1801, Syst. Eleuth., 1, p. 438 (Carolinae).
quadripunctata SUFFRIAN, 1858, Stett. ent. Ztg., 19, p. 389 (N. America).
lapponica var. *rubrocoeruleum* GRANDL, 1881, Ent. Nachr., 7, p. 305 (*Melasoma*) (Egerlandes).
lapponica var. *bimaculatum* GRANDL, 1881, ibid., p. 305 (*Melasoma*) (Egerlandes).
lapponica var. *bipustulatum* GRANDL, 1881, ibid., p. 305 (*Melasoma*) (Egerlandes).
lapponica var. *sexmaculatum* GRANDL, 1881, ibid., p. 306 (*Melasoma*) (Egerlandes).
lapponica var. *quadrimeaculatum* GRANDL, 1881, ibid., p. 306 (*Melasoma*) (Egerlandes).
lapponica var. *coerulea* GRANDL, 1881, ibid. p., 306 (*Melasoma*) (Egerlandes).
lapponica var. *marginatum* GRANDL, 1881, ibid., p. 306 (*Melasoma*) (Egerlandes).
lapponica var. *altaica* WEISE, 1884, Ins. Deutschl., 6 (3), p. 557 (*Melasoma*) (Siberia).
lapponica var. *unicolor* MARSEUL, 1888, Abeille, Paris, 26, p. 13 (*Lina*) (no locality cited).
lapponica var. *litua* MARSEUL, 1888, ibid., p. 13 (*Lina*) (no locality cited).
lapponica var. *quadripustulatum* JACOBSON, 1900, Horae Soc. ent. ross., 35, p. 90 (*Melasoma*) (Siberia occidentalis).
lapponica var. *crucliatum* JACOBSON, 1901, Öfvers. Finska. Vetensk Soc. Förh., 43, p. 133 (*Melasoma*) (W. Siberia).
lapponicum var. *multipunctatum* JACOBSON, 1901, ibid., p. 137 (*Melasoma*) (W. Siberia).
lapponicum var. *violaceipenne* JACOBSON, 1901, ibid., p. 133 (*Melasoma*) (W. Siberia).
interrupta var. *quadriguttata* SCHAEFFER, 1928, Canad. Ent., 60, p. 43 (*Lina*) (N. America).
interrupta var. *aeneicollis* SCHAEFFER, 1928, ibid., p. 43 (*Lina*) (N. America).
Distr.: Jpn (Hokkaido), N. China, Siberia, Europe, N. America.

C. populi LINNAEUS, 1758 ドロノキハムシ

LINNAEUS, 1758, Syst. Nat., ed. 10, p. 370 (no locality cited).
populi var. *janaceki* REITTER, 1892, Wien, ent. Ztg., 11, p. 240 (*Melasoma*) (Braunberg in Mahren).
Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu), Siberia, Mongolia, India, China, Korea, Europe, W. Asia.

C. tremulae FABRICIUS, 1787 ポプラハムシ

FABRICIUS, 1787, Man. Ins., 1, p. 69 (Europe).
longicollis SUFFRIAN, 1851, Linn. Ent., 5, p. 199 (Europe).
Distr.: Japan (Hokkaido), China, Korea, Sachalin, Siberia, Mongolia, Europe,

N. America.

C. vigintipunctata (SCOPOLI, 1763) ヤナギハムシ

- SCOPOLI, 1763, Ent. Carn., p. 78 (*Coccinella*) (no locality cited).
octodecimpunctata LINNAEUS, 1767, Syst. Nat., ed. 12, p. 582 (*Coccinella*) (Europe).
vigintimaculata HARRER, 1784, Beschr. Ins. Schaff., p. 157 (no locality cited).
vigintipunctata var. *pustulata* WEISE, 1884, Ins. Deutschl., 6 (3), p. 559 (*Melasoma*) (no locality cited).
vigintipunctata var. *incontaminata* HEYDEN, 1887, Horae Soc. ent. ross., 21, p. 263 (*Melasoma*) (Korea).
vigintipunctata var. *costella* MARSEUL, 1888, Abeille, Paris, 26, p. 10 (*Lina*) (Japan).
vigintipunctata var. *diluta* MARSEUL, 1888, ibid., p. 10 (*Lina*) (no locality cited).
vigintipunctata var. *miniata* AUEL, 1909, Z. Wiss.-Inseken-biol., 5, p. 318 (*Melasoma*) (Pfakuerinsel bei Potsdam).
vigintipunctata ab. *quadripunctata* SCHULTZ, 1911, Int. ent. Z., Guben, 5, p. 153 (*Melasoma*) (Finkenkrug).
vigintipunctata ab. *deegeneri* SCHULTZ, 1911, ibid., 5, p. 153 (*Melasoma*) (Finkenkrug).
vigintipunctata ab. *extrema* SCHULTZ, 1911, ibid., 5, p. 153 (*Melasoma*) (Finkenkrug).
vigintipunctata ab. *melanina* SCHULTZ, 1911, ibid., 5, p. 153 (*Melasoma*) (Finkenkrug).
Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yakushima, Tsushima, Kuchinoerabu), Siberia, Mongolia, China, Taiwan, Korea, Europe.

Genus *Linaeidea* MOTSCHULSKY, 1860 ルリハムシ属

Linaeidea MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 199 (type species: *Chrysomela aenea* LINNAEUS, 1798).

Macromela CHÛJÔ, 1958, Quart. J. Taiwan Mus., 9, p. 31.

L. aenea (LINNAEUS, 1758) ルリハムシ

- LINNAEUS, 1758, Syst. Nat., ed. 10, p. 369 (*Chrysomela*) (Europe).
betulae SCOPOLI, 1763, Ent. Carn., p. 71 (*Coccinella*) (no locality cited).
vitellinae SCOPOLI, 1763, ibid., p. 72 (*Coccinella*) (no locality cited).
coeruleoviolacea DEGEER, 1775, Mém. Hist. Ins., 5, p. 316 (*Chrysomela*) (no locality cited).
haemorrhoidalis MÜLLER, 1776, Zool. danic. Prodr., p. 81 (*Chrysomela*) (no locality cited).
fuscicornis FABRICIUS, 1781, Spec. Ins., 1, p. 126 (*Chrysomela*) (Russia).
amethystica GMELIN, 1790, in LINNAEUS, Syst. Nat., ed. 13, 1, 4, p. 1688 (*Chrysomela*) (no locality cited).
violacea OLIVIER, 1790, Encycl. Méth., 5, p. 718 (*Chrysomela*) (Suede).
aenea var. *bicolor* SCHILSKY, 1908, Dtsch. ent. Z., 1908, p. 603 (*Melasoma*) (Europe).
Distr.: Japan, Siberia, Mongolia, NE. China, Sachalin, Kuriles, Europe.
subsp. *aenea* (LINNAEUS, 1758)
LINNAEUS, 1758, Syst. Nat., ed. 10, p. 369 (*Chrysomela*) (Europe).
Distr.: Japan (Hokkaido, Honshu, Sado), Siberia, NE. China, Sachalin, Kuriles, Europe.
subsp. *insularis* (CHÛJÔ, 1940)
CHÛJÔ, 1940, Kontyû, Tokyo, 14, p. 78 (*Chrysomela*) (as subsp. of *adamsii* BALY) (Kochi).

- C.** Distr.: Japan (Shikoku, Kyushu, Yakushima).
 subsp. *tsutsuii* NAKANE, 1955
 NAKANE, 1955, Sci. Rept. Saikyo Univ., **2** (1), p. A38 (as subsp. of *aenea* LINNAEUS)
 (Wakayama).
 Distr.: Japan (S. Honshu).
- Genus *Paropsides* MOTSCHULSKY, 1860 ヤマトヨダンハムシ属
- C.** *Paropsides* MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., **2**, p. 192 (type species:
Paropsis duodecimpustulata GEBLER, 1825).
Aparopsis WEISE, 1908, Nova Guinea, **5**, p. 315.
- P. duodecimpustulata** (GEBLER, 1825) ヤマトヨダンハムシ
 GEBLER, 1825, in HUMMEL, Essai Ent., **4**, p. 54 (*Paropsis*) (Siberia).
duodecimpustulata var. *hieroglyphica* GEBLER, 1825, ibid., p. 55 (*Paropsis*) (Siberia).
maculicollis JACOBY, 1889, Entomologist, **23**, p. 118, pl. 1, fig. 10 (China).
duodecimpustulata melli REINECK, 1922, Dtsch. ent. Z., **1922**, p. 370 (Canton).
duodecimpustulata yuasai OHNO, 1958, Kontyû, Tokyo, **16**, p. 34, figs. (Yamanashi,
 Nagano, Gumma).
 Distr.: Japan (Honshu), E. Siberia, China, Korea, Vietnam, Burma, India.
- Genus *Gonioctena* CHEVROLAT, 1837 トホシハムシ属
- Gonioctena* CHEVROLAT, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 403 (type species:
Chrysomela viminalis LINNAEUS, 1758).
Phytodecta KIRBY, 1837, Fauna Bor. Amer., **4**, p. 213 (type species: *Chrysomela rufipes*
 DEGEER, 1775).
- Subgenus *Gonioctena* s. str.
- G. (G.) honshuensis** NAKANE, 1963 ヤナギホシハムシ
 NAKANE, 1963, Fragm. coleopt., ed. NAKANE, (5), p. 19 (Shimashima).
sibirica: KIMOTO, 1963, Fragm. Coleopt., ed. NAKANE, (3), p. 4 (Hokkaido).
chujoii MEDVEDEV, 1966, Forest Ent.-Fauua Kuril, Kamch. & Magda., p. 41, fig. 2
 (Kuriles).
 Distr.: Japan (Hokkaido, Honshu), Kuriles, Siberia.
- G. (G.) japonica** CHÛJÔ et KIMOTO, 1960 トホシハムシ
 CHÛJÔ et KIMOTO, 1960, Niponius, Takamatsu, **1** (4), p. 5, figs. (Hokkaido, Honshu,
 Shikoku).
rufipes: BALY, 1874, Trans. ent. Soc. London, **1874**, p. 173 (*Gonioctena*) (Hakodadi).
 Distr.: Japan (Hokkaido, Honshu, Shikoku, Kyushu).
- G. (G.) kamiyai** KIMOTO, 1963 カンバムジハムシ
 KIMOTO, 1963, Fragm. coleopt., ed. NAKANE, (3), p. 14 (Masutomi, Shimashima).
 Distr.: Japan (Honshu).
- G. (G.) morimotoi** KIMOTO, 1963 サクラムジハムシ
 KIMOTO, 1963, Fragm. coleopt., ed. NAKANE, (4), p. 15 (Karuizawa, Masutomi).
 Distr.: Japan (Honshu).
- G. (G.) shibatai** TAKIZAWA, 1982 シバタホシハムシ
 TAKIZAWA, 1982, Ent. Rev. Japan, **37**, p. 57 (Nukumidaira, Yamagata Pref.).
 Distr.: Japan (Honshu).

- G. (G.) sibirica** (WEISE, 1893) ヤナギムジハムシ
 WEISE, 1893, Ins. Deutschl., 2, p. 1128 (*Phytodecta*) (Siberia).
sorbina WEISE, 1916, Coleopt. Cat. Junk, 68, p. 176 (*Phytodecta*) (n. n. for *sorbi*
 MOTSCHULSKY, nec GRIMMER, 1841).
sorbi MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 223, pl. 11, fig. 6
 (Dauria).
cheni BECHYNÉ, 1947, Acta Mus. natn. Prag., 3B (3), pp. 102, 134, pl. 1, fig. 8, pl. 5,
 figs. 6–7 (Ussuri, Amur).
 Distr.: Japan (Hokkaido), E. Siberia, Mongolia.
- G. (G.) springlovae** (BECHYNÉ, 1947) クロモンハムシ
 BECHYNÉ, 1947, Acta Mus. natn. Prag., 3B (3), pp. 115, 116, figs. (*Phytodecta*)
 ("Kioto").
springlovae ab. *grandialis* BECHYNÉ, 1947, ibid., p. 116, fig. (*Phytodecta*) ("Kioto").
gracilicornis? JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 210 (*Phytodecta*)
 (Japan).
 Distr.: Japan (Hokkaido).
- G. (G.) takahashii** KIMOTO, 1963 ヒメトホンハムシ
 KIMOTO, 1963, Fragm. coleopt., ed. NAKANE, (4), p. 15 (Tochigi, Tokushima,
 Nagano).
 Distr.: Japan (Honshu, Shikoku).
- Subgenus *Brachyphytodecta* BECHYNÉ, 1948
- Brachyphytodecta* BECHYNÉ, 1948, Acta Mus. natn. Prag., 3B (3), p. 101 (type species:
Spartophila fulva MOTSCHULSKY, 1860).
- G. (B.) rubripennis** BALY, 1862 フジハムシ
 BALY, 1862, Annls. Mag. nat. Hist., (3), 10, p. 28 (Japan).
rubripennis var. *tenebrosa* WEISE, 1910, Verh. naturf. Ver. Brünn, 48, p. 37 (*Phytodecta*) (Japan).
rubripennis var. *plagipennis* ACHARD, 1924, Čas. Čs. Spol. ent., 21, p. 33 (*Phytodecta*) (Yokohama).
 Distr.: Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima).
- Subgenus *Sinomela* CHEN, 1935
- Sinomela* CHEN, 1935, Chinese J. Zool., 1, p. 126 (type species: *Phytodecta aeneipennis*
 BALY, 1862).
- G. (S.) hikosanus** KIMOTO, 1974 ヒコサンクロボシハムシ
 KIMOTO, 1974, Kontyû, Tokyo, 42, p. 144, fig. (Hikosan).
 Distr.: Japan (Kyushu).
- G. (S.) nagaii** NAKANE, 1963 リュウキュウクロボシハムシ
 NAKANE, 1963, Sci. Rept. Kyoto pref. Univ., (Nat. Sci. & Lib. Sci.), 3 (5), p. 12
 (Okinoerabu Is.).
 Distr.: Ryukyu Is. (Okinoerabu).
- G. (S.) nigroplagiata** BALY, 1862 ヤツボシハムシ
 BALY, 1862, Annls. Mag. Nat. Hist., (3), 10, p. 28 (Japan).
robusta JACOBY, 1885, Proc. zool. Soc. London, 1885, p. 209 (*Phytodecta*) (Japan).
 Distr.: Japan (Honshu), E. China.

Genus *Phratora* CHEVROLAT, 1837 ホソルリハムシ属

Phratora CHEVROLAT, 1837, in DEJEAN, Cat. Coléopt., ed. 3, p. 405 (type species: *Chrysomela vitellinae* LINNAEUS, 1758).

Phyllodecta KIRBY, 1837, Fauna Bor. Amer., 4, p. 216.

Subgenus *Phratora* CHEVROLAT, s. str.

P. (P.) grandis (CHÛJÔ, 1956) オオホソルリハムシ

CHÛJÔ, 1956, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., 2 (31), p. 6, fig. 3 (*Phyllodecta*) (Gumma, Yamagata, Rishiri-tô).

Distr.: Japan (Hokkaido, Honshu).

P. (P.) laticollis SUFFRIAN, 1851 ヤマナラシハムシ

SUFFRIAN, 1851, Linn. Ent., 5, p. 262 (Europe).

multipunctatus JACOBY, 1890, Entomologist, 23, p. 117 (*Phyllodecta*) (China).

Distr.: Japan (Hokkaido, Honshu), China, Mongolia, Siberia, Asia Minor, Armenia, Europe.

Subgenus *Chaetocera* WEISE, 1884

Chaetocera WEISE, 1884, Ins. Deutschl., 6 (3), p. 514 (type species: *Chrysomela vulgatissima* LINNAEUS, 1758).

Chaetoceroides STRAND, 1935, Fol. zool. hydrobiol., 7, p. 285.

P. (C.) obtusicollis MOTSCHULSKY, 1860 ホソルリハムシ

MOTSCHULSKY, 1860, Schrenck's Reisen Amurl., 2, p. 230 (Amur).

inhonesta WEISE, 1884, Ins. Deutschl., 9 (3), p. 514, nota (*Phyllodecta*) (Kjachata, Sarepta).

longulus: CHÛJÔ, 1959, Mem. Fac. Lib. Arts & Educ. Kagawa Univ., 2 (81), p. 9 (*Phyllodecta*) (Hokkaido).

Distr.: Japan (Hokkaido), S. Kuriles (Etorofu), Korea, Sachalin, NE. China, Mongolia, E. Siberia.

Genus *Phola* WEISE, 1890 ハマゴウハムシ属

Phola WEISE, 1890, Horae Soc. ent. ross., 24, p. 482 (type species: *Chalcolampra octodecimguttata* FABRICIUS, 1775).

Phyllophila STÅL, 1857, Öfvers. Vetensk.-Akad. Förh., 14, p. 59.

P. octodecimguttata (FABRICIUS, 1775) ハマゴウハムシ

FABRICIUS, 1775, Syst. Ent., 100 (*Chrysomela*) ("Nova Hollandia").

cybele STÅL, 1860, Öfvers. Vetensk-Akad. Förh., 17, p. 464 (*Chalcolampra*) (Hong-kong).

viticis FAIRMAIRE, 1888, Annls. Soc. ent. Belg., 1888, p. 39 (*Chalcolampra*) (China).

keyserlingi WEISE, 1890, Horae Soc. ent. ross., 24, p. 482 (*Chalcolampra*) (China).

Distr.: Japan (Kyushu), Ryukyu Is. (Tokara, Amami-Oshima, Kikaiga-shima, Oki-erabu, Okinawa, Ishigaki, Iriomote, Yonaguni), China, Taiwan, Vietnam, Burma, India, Ceylon, Malaya, Philippines, New Guinea.

Genus *Potaninia* WEISE, 1889 アカゾハムシ属

Potaninia WEISE, 1889, Horae Soc. ent. ross., 23, p. 603 (type species: *Potaninia polita* WEISE, 1889).

Suizone CHEN, 1931, Bull. Soc. ent. France, **1931**, p. 130.

P. cyrtonoides (JACOBY, 1885) アカゾハムシ

JACOBY, 1885, Proc. zool. Soc. London, **1885**, p. 206 (*Chrysomela*) (Konose).

Distr.: Japan (Honshu, Shikoku, Kyushu).