命名した. 後者は、中国から知られている H. sculpticollis FAIRMAIRE である.

References

KASZAB, Z., 1941. Die indomalayischen Misolampinen (Coleopt., Tenebr.). Annls. Mus. nat. hung., 34: 1-44, 1 pl.

Elytra, Tokyo, 18 (2): 230, November 15, 1990

A New Record of the Taiwanese Opisthiine *Paropisthius* masuzoi (Coleoptere, Carabidae)¹⁾

Shun-Ichi UÉNO

Department of Zoology, National Science Museum (Nat. Hist.), Shinjuku, Tokyo, 169 Japan

Paropisthius masuzoi Kasahara (1989, Elytra, Tokyo, 17, p. 114, figs. 1–3) was described from three localities in the northern part and one locality in the southern part of the Taiwanese high mountains. Through the autumn expedition made in 1990, a new locality of this interesting carabid beetle was found by Mr. Kun Fu Shih in the central part of the island. The collecting data are as given below.

1 ♀, Kuan-kao on the Yü-shan Mountains, 2,550 m in altitude, in Hsin-i Hsiang of Nan-t'ou Hsien, central Taiwan, 26-X-1990, K. F. Shih leg. (NSMT).

Like the original material, the specimen recorded above was found from beneath a stone lying in a very wet spot at the side of a narrow stream which cascaded down a steep slope in a coniferous forest.

¹⁾ This study is supported by the Grant-in-aid No. 01041099 for Field Research of the Monbusho International Scientific Research Program, Japan.