

つこと、雄交尾器の背板が長いことによって区別できる。ヤエヤマタテスジジョウカイは、外見がハナタニジョウカイにきわめてよく似ているが、体がやや大きく全体的に濃色であること、雄交尾器の背板の先端がいくぶん尖ることで区別できる。

References

- ISHIDA, K., 1986. A revision of the two genera, *Athemus* and *Athemellus*, of Japan (Coleoptera: Cantharidae). *Trans. Shikoku ent. Soc.*, **17**: 193–213.
- SATÔ, M., 1985. Cantharidae. In KUROSAWA, Y., et al. (eds.), *The Coleoptera of Japan in Color*, **3**: 107–118, pls. 17–19. Hoikusha, Osaka. (In Japanese.)
- WITTMER, W., 1970. Coleoptera of the Loo-Choo Archipelago (II). 17. Families Cantharidae and Malachiidae. Cantharidae und Malachiidae von den Inseln Okinawa und Ryukyu, gesammelt durch Herrn Prof. Dr. Michio CHÛJÔ. *Mem. Fac. Educ. Kagawa Univ.*, (II), (192): 22–27.
- 1972. 55. Beitrag zur Kenntnis der palaearktischen Cantharidae und Malachiidae (Col.). *Ent. Arb. Mus. Frey*, **23**: 122–141.
- 1983. Die Familie Cantharidae (Col.) auf Taiwan (2. Teil). *Ent. Rev. Japan*, **38**: 147–172, pls. 10–14.

Elytra, Tokyo, **24** (1): 124, May 15, 1996

A New Record of *Stenhomalus fenestratus* (Coleoptera, Cerambycidae) in Indochina

Tatsuya NIISATO

Bioindicator Co., Ltd., Takada 3–16–4, Toshima-ku, Tokyo, 171 Japan

Stenhomalus fenestratus WHITE is a species widespread in the continental side of Asia including Taiwan, though previously recorded from only northern Thailand in the Indochinese Region. Recently, I had an opportunity to examine a male specimen of the species collected from northern Vietnam, as recorded below:

Specimen examined. 1 ♂, near Sapa, Lao Cai Province of northern Vietnam, 20–V–1995, N. KATSURA leg.

This *Stenhomalus* is known as a species showing slight geographical variation (NIISATO & MAKIHARA, 1991). The present specimen is rather similar to the specimens from northern Thailand in the darker coloration and broad body form. The measurements of body parts are as follows (in mm): BL 9.7, AL 14.5, HW 1.8, PL 1.75, PA 1.3, PB 1.2, PW 1.63, EL 5.8, EW 2.5.

I am indebted to Mr. Takao ARAI for his kind support of material.