

New Localities of *Pseudotarphius lewisi*  
WOLLASTON (Coleoptera, Colydiidae)

Keiji OKADA

Bioindicator Co., Ltd., Takada 3-16-4, Toshima-ku, Tokyo, 171 Japan

The colydiid beetle, *Pseudotarphius lewisi* WOLLASTON, 1873, was originally described from Nagasaki and Yuyama in Kyushu, and Hagi in Honshu, Japan. After that, SASAJI (1985) recorded the species from Yaku-shima Island and Amami-Ōshima Island without detailed collecting data. Recently, I had an opportunity to examine some specimens of this colydiid species collected from Shikoku and the Tokara Islands in the collection of the Entomological Laboratory, Ehime University, and the private collection of Dr. Masahiro SAKAI of the same university. These are the first records of the species from the two areas.

*Specimens examined.* [Shikoku] (Kagawa Pref.) 1 ex., Mt. Zōzu-san, 16-VII-1978, M. SAKAI leg.; 2 exs., Mt. Zōzu-san, 16-VII-1978, A. ODA leg. [Tokara Isls.] 1 ex., Nakanoshima Is., 28-VII-1969, M. SAKAI leg.; 2 exs., same locality and collector, 2-V-1971; 1 ex., same locality and collector, 4-V-1971.

*Distribution.* Japan (Honshu, Shikoku, Kyushu, Yaku-shima Is., Tokara Isls., Amami-Ōshima Is.).

I wish to express my sincere gratitude to Dr. Masahiro SAKAI for his kind supply of specimens.

References

- DAJOZ, R., 1977. Coléoptères Colydiidae et Anommatidae Paléarctiques. *Fauna de l'Europe et du Bassin Méditerranéen*, 8: 126-127.
- SASAJI, H., 1985. Colydiidae. In KUROSAWA, Y., S. HISAMATSU & H. SASAJI (eds.), *Coleopt. Japan Col., Osaka*, 3: 291-294, pl. 48. (In Japanese.)
- WOLLASTON, T. V., 1873. On a new genus of Colydiidae, from Japan. *Trans. ent. Soc. London*, 1873: 1-4.