

New Record of *Flavohelodes burmensis* (KLAUSNITZER, 1974)  
(Coleoptera, Scirtidae) from Thailand

Hiroyuki YOSHITOMI

Zoological Laboratory, Faculty of Agriculture, Meijo University,  
Tempaku-ku, Nagoya, 468 Japan

Up to the present, only two males of *Flavohelodes burmensis* (KLAUSNITZER, 1974) have been known from Myanmar (type area) and Nepal. Recently, I had an opportunity to examine four male specimens of this species collected from Thailand and Nepal. In this short paper, I am going to give a new record from Thailand and an additional record from Nepal.

Before going further, I wish to sincerely thank Professor M. SATÔ, Nagoya Women's University, for giving me the opportunity to examine the specimens, and Prof. Dr. Y. ARITA and Dr. S. HASHIMOTO, Meijo University, for their advice and encouragement.

*Flavohelodes burmensis* (KLAUSNITZER, 1974)

*Helodes burmensis* KLAUSNITZER, 1974, Ent. Nachr., Dresden, **18**: 19–20.

*Flavohelodes burmensis*: KLAUSNITZER, 1980, Reichenbachia, **18**: 85; 1980, Ent. basil., **5**: 196, 199.

Measurements (n=4). Pronotal width 1.3–1.4 ( $x=1.35$ ) mm; pronotal length 0.8–1.0 ( $x=0.88$ ) mm; elytral length 2.8–3.2 ( $x=2.98$ ) mm; elytral width 1.7–1.9 ( $x=1.78$ ) mm; total length (pronotal length plus elytral length) 3.6–4.2 ( $x=3.85$ ) mm.

*Specimens examined.* [Thailand] 2 ♂♂, Maeo Khun Klang, 1,350 m, Doi Inthanon, 20–X–1983, M. SAKAI leg. (male genitalia on slides Nos. HY 293–294). [Nepal] 2 ♂♂, Bogara, 1,700–2,200 m, 83°23'E, 28°35'N, 19–20–IX–1971, Malaise trap, A. NAKANISHI leg. (male genitalia on slide No. HY 295).

*Distribution.* Myanmar, Nepal, Thailand (new record).



Fig. 1. *Flavohelodes burmensis* (KLAUSNITZER), male, from Nepal.