


On the Male of *Trichophya japonica*
(Coleoptera, Staphylinidae)

Kiminori MIYASHITA

Tôkô-chô 110-23, Ebetu-shi, Hokkaido, 067 Japan

Trichophya japonica Y. WATANABE et Y. SHIBATA has previously been known from only female specimens (WATANABE & SHIBATA, 1962; NAOMI, 1995). Recently, I reported a collection of male specimens of this species from Hokkaido (MIYASHITA, 1996). In the present paper, the aedeagus and the 8th and 9th male sternites of *T. japonica* are illustrated for the first time (Figs. 1-3).



Figs. 1-3. *Trichophya japonica* Y. WATANABE et Y. SHIBATA, ♂; 1, sternite 8; 2, sternite 9; 3, aedeagus. Scale: 0.1 mm.

Remarks. All the specimens examined were collected by myself by beating a kind of fungi, *Pleurotus cornucopiae* var. *citrinopileatus* (Japanese name: tamogitake).

References

- MIYASHITA, K., 1996. Coleoptera fauna of the upper reaches of Riv. Otofuke. *Antenna*, 3: 21-44. (In Japanese.)
NAOMI, S., 1995. Description of a new species of the genus *Trichophya* (Coleoptera, Staphylinidae) from Japan. *Spec. Bull. Jpn. Soc. Coleopterol., Tokyo*, (4): 347-350.
WATANABE, Y., & Y. SHIBATA, 1962. Description of a new species of the genus *Trichophya* MANNERHEIM from Japan (Col. Staphylinidae). *J. agric. Sci. Tokyo*, 7: 95-96.