

ratanai sp. nov. を記載した。この新種は、スマトラ産の種とは、頭盾の前縁が弓形にえぐれること、触角が短いこと、小盾板に点刻があること、そして上翅に3本の黒帯がないこと、などの差異によって容易に区別できる。

References

- ENDO, S., 1992. A new species of *Pseudohyperantha* from Malaysia (Coleoptera, Buprestidae). *Jewel Beetles, Tokyo*, (1): 6-7.
 SAUNDERS, E., 1869. Descriptions of nine new species of Buprestidae. *Trans. ent. Soc. London*, **1869**: 1-8.
 THÉRY, A., 1935. Note sur les Buprestidae du muséum de Leiden. *Zool. Meded., Leiden*, **18**: 241-256.
 TÔYAMA, M., 1989. A note on the buprestid genus *Pseudohyperantha* (Coleoptera, Buprestidae). *Kanagawa-Chûhō, Yokohama*, (90): 193-196.

Elytra, Tokyo, **25** (1): 108, May 15, 1997

New Localities of *Cis sasakawai* NOBUCHI (Coleoptera, Ciidae)

Makoto KAWANABE

Bioindicator Co., Ltd., Takada 3-16-4, Toshima-ku, Tokyo, 171 Japan

The ciid beetle, *Cis sasakawai* NOBUCHI, 1960 [Japanese name: Sasakawa-tsutsukinokomushi], was originally described from Hokkaido, Japan. This is one of the rare species of Japanese ciids, and MIYATAKE (1985) recorded it from Shikoku without detailed collecting data. These are all the records of this species up to the present. In the course of my revisional study of the Japanese Ciidae, I gathered up the collecting data of *C. sasakawai* on the basis of the collection of Ehime University and other private collections. These data contain the first records from Honshu and Kyushu.

Specimens examined. [Hokkaido] 1 ♂, 1 ♀, Takinosawa, near Sapporo, 8-IV-1956, A. NOBUCHI leg. (cotypes); 1 ♀, Pyuka, near Nayoro, 5-X-1955, A. NOBUCHI leg. (cotype). [Honshu] (Hiroshima Pref.) 15 exs., Ōasa, Ōasa-chō, 25-VI-1994, M. KAWANABE leg. [Shikoku] (Ehime Pref.) 1 ex., Naose near Kuma-chō, 16-V-1953, M. MIYATAKE leg.; 2 exs., Yoshinogawa, Oda-chō, 15-VII-1993, M. KAWANABE leg. (Kōchi Pref.) 59 exs., Cape Ashizuri-misaki, 25-VI-1989, M. KAWANABE leg. [Kyushu] (Ōita Pref.) 12 exs., Kōbaru, 20-VII-1989, M. KAWANABE leg. (Nagasaki Pref.) 3 exs., Tsushima Isls., Kamiagata, 16-XII-1974, M. MOGI leg.

Distribution. Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima Isls.).

Host fungi. *Coriolus versicolor* (L.: FR.) QUÉL. (Kawaratake in Japanese) and *Pycnoporus coccineus* (FR.) BOND. et SING. (Hirotake in Japanese).

References

- MIYATAKE, M., 1985. Ciidae. In KUROSAWA, Y., S. HISAMATSU & H. SASAJI (eds.), *The Coleoptera of Japan in Color*, **3**: 278-285 [incl. pl. 46]. Hoikusha, Osaka. (In Japanese.)
 NOBUCHI, A., 1960. Ciidae from Hokkaido (Coleoptera). *Ent. Rev. Japan, Osaka*, **11**: 37-41 [incl. pl. 5].