

Haut Yunnan", "Zool. Mus. Berlin"; 2 ♀♀, "Kiukiang China", "Zool. Mus. Berlin".

Two specimens from Sapa are donated to the Museum für Naturkunde der Humboldt-Universität zu Berlin and the Museum of Nature and Human Activities, Hyôgo, respectively.

要 約

伊藤 武：ベトナム産アカチャコガネの一種，*Brahmina ciliaticollis* MOSERの再記載。——本種は、北ベトナムから得られた3個体の雌標本に基づいて、MOSERにより記載されて以来、現在まで形態学的な補足説明も追加記録もなされていなかった。今回、本種の雄個体を含めて多数の標本を検視することができたので、改めてここに再記載を行った。

Literature Cited

MOSER, J., 1914. Neue Melolonthiden-Arten aus der Sammlung des Herrn Dr. H. J. VETH (Haag). *Notes Leyden Mus.*, **36**: 148–156.

Elytra, Tokyo, **25** (1): 116, May 15, 1997

Occurrence of *Zoodes formosanus* (Coleoptera, Cerambycidae) in Northern Vietnam

Tatsuya NIISATO

Bioindicator Co., Ltd., Takada 3-16-4, Toshima-ku, Tokyo, 161 Japan

Zoodes formosanus is a little-known hesperophanine cerambycid beetle and has so far been known as an endemic to Taiwan. Recently, I had an opportunity to examine a single female specimen of the same species collected near the northern border of Vietnam. It is an interesting example from the zoogeographical viewpoint, exhibiting a relationship of the cerambycid faunas between northern Vietnam and Taiwan. The facies of the Vietnamese specimen are similar to those of the type population, except for some weak differences regarded as infraspecific variation.

Zoodes formosanus NIISATO, 1982

Zoodes formosanus NIISATO, 1982, Trans. Shikoku ent. Soc., **16**, pp. 41–43, figs. 1–2.

Specimen examined. 1 ♀, near Sapa, Lao Cai Prov., N. Vietnam, V-1996.

Distribution. Taiwan, N. Vietnam (new record).