

が多かった。本研究において、既知種の再検討とともに、第1の群では9新種、第2の群では4新種、第3の群では2新種を記載した。なお、これまでニューギニアおよびその周辺の島じまのみから知られていた *Strongylium tuberipenne* FAIRMAIRE, 1881 が、オーストラリア（北部クイーンズランド）にまで分布していることがわかった。

### References (Additional)

- FAIRMAIRE, L., 1881. Diagnoses de Coléoptères de la Micronésie et de la Polynésie. *Natural., Paris*, **3** (45): 359.
- 1883. Essai sur les Coléoptères de l'Archipel de la Nouvelle-Bretagne. *Annl. Soc. ent. Belg.*, **27**: 1–58.
- 1893. Coléoptères nouveaux des Indes orientales, de la famille des Scarabaeidae, Rhipidoceidae, Tenebrionidae et Oedemeridae. *Not. Leyden Mus.*, **15**: 17–64.
- 1900. Description d'une espèce nouvelle du genre *Strongylium* [Col.]. *Bull. Soc. ent. Fr.*, **1900**: 45–46.
- GEBIEN, H., 1913. Die Tenebrioniden der Philippinen. *Philipp. J. Sci.*, **8** (D): 401–433.
- 1920. Coleoptera, Tenebrionidae. *Nova Guinea*, XIII, Zool., **3**: 213–500.
- 1921. Philippine Tenebrionidae, II. *Philipp. J. Sci.*, **19**: 439–519, 2 pls.
- KASZAB, Z., 1977. Die Tenebrioniden des Papuanischen Gebietes. 1. Strongyliini (Coleoptera: Tenebrionidae). *Pacif. Ins. Mon.*, **33**: 1–219.
- KIRBY, W., 1818. A century of insects, including several new genera described from his cabinet. *Trans. Linn. Soc. London*, **12**: 375–478, 3 pls.
- KULZER, H., 1966. Australische und papuanische Strongyliini. *Ent. Arb. Mus. Frey*, **17**: 338–396.
- PIC, M., 1917. Descriptions abrégées diverses. *Mél. exot.-ent.*, **23**: 2–20.

---

*Elytra, Tokyo*, **25** (2): 309–310, November 15, 1997

### New Synonymy in the Genus *Strongylium* (Coleoptera, Tenebrionidae, Strongyliini) Described by FAIRMAIRE and PIC

**Kimio MASUMOTO**

Institute of Human Living Sciences, Otsuma Women's University,  
12 Sanbancho, Chiyoda-ku, Tokyo, 102 Japan

In the course of my study on the Asian Strongyliini, I had the opportunity of examining type specimens of the species described by FAIRMAIRE and PIC preserved in the Muséum National d'Histoire Naturelle, Paris. The following new synonyms were confirmed in the course of this study.

***Strongylium bidentaticolle* PIC, 1927**

*Strongylium bidentaticolle* PIC, 1927, Bull. Soc. ent. Fr., **1927**: 112.

*Strongylium hirtipes* PIC, 1940, Échange, Moulins, (479): 3. [Syn. nov.]

***Strongylium brevesulcatum* PIC, 1931**

*Strongylium brevesulcatum* PIC, 1931, Mél. exot.-ent., (57): 36.

*Strongylium dalatense* PIC, 1931, Mél. exot.-ent., (57): 36. [Syn. nov.]

*Notes.* PIC gave the two names cited above on the same page. Though *Strongylium dalatense* has precedence over *S. brevesulcatum*, the former is based on a female. As the first reviser, I prefer to select the latter as the senior synonym.

***Strongylium fisicolle* FAIRMAIRE, 1903**

*Strongylium fisicolle* FAIRMAIRE, 1903, Anns. Soc. ent. Belg., **47**: 18.

*Strongylium sulcaticeps* PIC, 1925, Mél. exot.-ent., (43): 11. [Syn. nov.]

***Strongylium infoveatum* PIC, 1917**

*Strongylium infoveatum* PIC, 1917, Mél. exot.-ent., (23): 15.

*Strongylium diversefoveipenne* PIC, 1936, Mél. exot.-ent., (68): 20. [Syn. nov.]

***Strongylium semiviolaceum* PIC, 1917**

*Strongylium semiviolaceum* PIC, 1917, Mél. exot.-ent., (23): 16.

*Strongylium curvatipes* PIC, 1922, Échange, Moulins, (408): 23. [Syn. nov.]

*Strongylium hoanum* PIC, 1938, Échange, Moulins, (474): 15. [Syn. nov.]

***Strongylium serricorne* FAIRMAIRE, 1888**

*Strongylium serricorne* FAIRMAIRE, 1888, Anns. Soc. ent. Fr., (6), **8**: 364.

*Strongylium impubens* PIC, 1922, Échange, Moulins, (408): 24. [Syn. nov.]