

Further Records of *Cis subrobustus* (Coleoptera, Ciidae)

Makoto KAWANABE

Bioindicator Co., Ltd., Takada 3-16-4, Toshima-ku, Tokyo, 171-0033 Japan

The ciid beetle, *Cis subrobustus* MIYATAKE, 1955 [Japanese name: Kimune-tsutsukinokomushi], was originally described from Shikoku, Japan (MIYATAKE, 1954). Though this is one of the easily distinguishable species in the genus *Cis*, little is known about its distribution. Up to the present, Shikoku and Kyushu are known as its distributional range (MIYATAKE, 1985), but the record with detailed collecting data is not found. In this report, I will sum up the collecting data of the species mainly on the basis of the collection of Ehime University and other private collections. These data contain the first records from Honshu. Second records from the same localities are omitted.

Specimens examined. [Honshu] <Kanagawa Pref.> 1 ex., Ôiso, 9-I-1972, Y. HIRANO leg.; 1 ex., Takeyama, 1-V-1988, Y. HIRANO leg. <Kyoto Pref.> 2 exs., Mt. Nanzenjisan, 1-VIII-1989, M. KAWANABE leg.; 2 exs., Mt. Hieizan, 5-VIII-1989, M. KAWANABE leg. <Nara Pref.> 1 ex., Mt. Kasugayama, 5-V-1955, T. SHIBATA leg. <Tottori Pref.> 1 ex., Mt. Daisen, 26~27-VII-1989, M. KAWANABE leg. <Okayama Pref.> 3 exs., Mt. Gagyûsan, 28-VII-1989, M. KAWANABE leg. <Hiroshima Pref.> 1 ex., Gokurakuji, 9-IV-1955, M. OKADA leg.; 9 exs., Ôasa, Ôasa-chô, 25~26-VI-1994, M. KAWANABE leg. [Shikoku] <Ehime Pref.> 9 exs., Naose near Kuma-chô, 16-V-1953, M. MIYATAKE leg. (including the holotype and 7 paratypes); 2 exs., Tobe-chô, 24-V-1955, K. SASAKI leg.; 1 ex., Ônogahara, 20-VIII-1953, T. MOHRI leg. (paratype); 9 exs., Kurotaki, Shigenobu-chô, 9-IV-1989, M. KAWANABE leg.; 2 exs., Ôtani-ike, Iyo-shi, 25-IV-1989, M. KAWANABE leg.; <Kôchi Pref.> 1 ex., Kuroson, 17-VII-1953, S. HISAMATSU leg. (paratype); 1 ex., Matsubagawa, 8-VII-1961, M. MIYATAKE leg. [Kyushu] <Fukuoka Pref.> 2 exs., Mt. Fukuchiyama, 3-V-1970, Y. TAKAKURA leg.; 2 exs., Mt. Ôsakayama, 1-V-1977, Y. TAKAKURA leg. <Ôita Pref.> 1 ex., Mt. Sobosan, 18~19-VII-1989, M. KAWANABE leg.

Distribution. Japan (Honshu, Shikoku, Kyushu).

Host fungi. *Coriolus versicolor* (L.: FR.) QUÉL. [Kawaratake], *Coriolus hirsutus* (WULF.: FR.) QUÉL. [Aragekawaratake], *Pycnoporus coccineus* (FR.) BOND. et SING. [Hiitake] and *Microporus vernicipes* (BERK.) O. KUNTZE [Tsuyauchiwatake].

References

- MIYATAKE, M., 1954. Studies on the Japanese Ciidae, I (Coleoptera). *Sci. Rept. Matsuyama agric. Coll., Matsuyama*, (14): 40-67, pls. 1-11.
——— 1955. Notes on some Japanese Ciidae (Coleoptera). *Trans. Shikoku ent. Soc., Matsuyama*, 4: 94-95.
——— 1985. Ciidae. In KUROSAWA, Y., S. HISAMATSU & H. SASAJI (eds.), *The Coleoptera of Japan in Color*, 3: 278-285 [incl. pl. 46]. Hoikusha, Osaka. (In Japanese.)