Occurrence of *Mecocerina angoonae* (Coleoptera, Anthribidae) in Northern Vietnam

Toshio SENOH

Department of Biology, Chuo University High School, Koganei, Tokyo, 184–8575 Japan

Mecocerina angoonae was described by SENOH (1997, p. 250) on the basis of a pair of the specimens collected by me from Doi Suthep, about 1,000 m alt., Chiang Mai Province, northern part of Thailand. There has been no other records than the original.

Late in the spring of 1998, I had an opportunity to visit the northern part of Vietnam and succeeded in obtaining a series of this species. The collecting site, Mt. Tam Dao, is distant for 700 km from the type locality of the species. The collecting data of the specimens taken are as follows:

6 & & QQ, Mt. Tam Dao (about 900–1,000 m alt.), Vinh Phu Prov., N. Vietnam, $4\sim5$ –V−1998, T. Senoh leg.




Fig. 1. Distribution of Mecocerina angoonae SENOH. — T, Tam Dao; D, Doi Suthep.

Distribution. N. Vietnam (new record) and N. Thailand.

Reference

Senoh, T., 1997. Two new species of the genus *Mecocerina* (Coleoptera, Anthribidae) from Vietnam and Thailand. *Elytra, Tokyo*, **25**: 247–253.