

Occurrence of *Obrium laosicum* (Coleoptera, Cerambycidae) in South China

Tatsuya NIISATO

Bioindicator Co., Ltd., Takada 3–16–4, Toshima-ku, Tokyo, 171–0033 Japan

and

Pu Fu-Ji

Institute of Zoology, Academia Sinica, Beijing 100080, China

In this short report, additional records are given for *Obrium laosicum* originally described from Phou Kham Khoay of Laos, Indochina. Two Chinese specimens examined agree well with the holotype female of *O. laosicum* deposited in the B. P. Bishop Museum, Honolulu, which was recently examined by the senior author. The species is unique among the congeners in having narrowly elongate pronotum with dense recumbent pubescence and simple sternites of female abdomen like that of *Ibidionidum* species. Apparently, *O. laosicum* and its relatives have to be discriminated in an isolated group in the tribe Obriini (NIISATO & OHMOTO, 1993).

***Obrium laosicum* GRESSITT et RONDON**

Obrium laosicum GRESSITT et RONDON, 1970, *Pacif. Ins. Mon.*, **24**, p. 107, fig. 20 d; type locality: Phou Kham Khoay, Vientiane Prov., Laos.

Specimens examined. 1 ♀, Jian-Feng Tian-Chi, Hainan Is., S. China, 12–V–1980, collected at light, PU F.-J. leg.; 1 ♀, He-Kong, 80 m alt., Yunnan, SW. China, 5–V–1956, HUANG K.-L. leg. (in coll. Academia Sinica, Beijing.)

Distribution. Laos; South China: Hainan and Yunnan (new records).

References

- GRESSITT, J. L., & J. A. RONDON, 1970. Cerambycids of Laos (Disteniidae, Prioninae, Philinae, Asemiae, Lepturinae, Cerambycinae). *Pacific. Ins. Mon.*, **24**: 1–314.
NIISATO, T., & T. OHMOTO, 1993. A new obriine species (Coleoptera, Cerambycidae, Cerambycinae) discovered from Iriomote-jima Island of the Ryukyu Islands. *Elytra, Tokyo*, **22**: 349–352.