

closely related to *C. shikokuensis* in expansion of the male fore and mid tarsi than to *C. shibatai*. This relationship agrees with that between *Trechiana onocoro* S. UÉNO from Awaji Is. and *T. tenuis* S. UÉNO / *T. satoui* S. UÉNO from Shikoku.

At the moment I hesitate to determine the taxonomic position of the Awaji population, since the number of available materials is too small to make discussion on its exact status including individual variation.

Lastly, I cordially thank Mr. Y. OKUDA and Mr. M. MORI for their kind offer of interesting materials.

Elytra, Tokyo, **27** (2): 598, November 13, 1999

New Records of Two Harpaline Beetles (Coleoptera, Cababidae)

Noboru Ito

1-7-18 Higashiuneno, Kawanishi City, Hyôgo Pref., 666-0117 Japan

Recently, I had an opportunity to examine many harpaline beetles from Asia and found two interesting species newly recorded from Sichuan and Thailand.

I wish to express my deep gratitude to Dr. Jan FARKAČ of the Česka Zemědělská Universita v Praze for his kind loan of a part of the materials used in this study.

1. *Coleolissus (Tenuistilus) cyanescens* N. ITO et WRASE, 1998
Coleolissus cyanescens N. ITO et WRASE, 1998, Ent. Rev. Japan, **53**: 43-46. (Yunnan in China).

Specimens examined. 2♂♂, Moxi, 29°13'N, 102°10'E, alt. 1,600 m, Sichuan, China, VII-1998 (N. ITO coll.).

This species was described only from Yunnan. I have now examined its specimens from Sichuan. Though the species is apterous, its distribution may be wide. Apterous trichotichine species are usually restricted in distribution, while *C. cyanescens* makes a sharp contrast in this respect to other species.

Distribution. Yunnan, Sichuan (new record).

2. *Pseudognathaphanus festivus* (ANDREWES, 1921)
Gnathaphanus festivus ANDREWES, 1921, Trans. ent. Soc. London, **1921**: 107-111. (Vietnam).
Pseudognathaphanus festivus: SCHAUBERGER, 1932, Kol. Rdsch., **18**: 57. ——— CSIKI, 1932, Coleopt. Cat., (121): 1041. ——— NOONAN, 1973, Quaest. ent., **9**: 344.

Specimens examined. 1♂, 1♀, Kanchanaburi, Thailand, 3-15-IV-1991, J. FARKAČ leg. (preserved in J. FARKAČ coll.); 1♂, 1♀, ditto, 31-III-1-IV-1994, J. HORÁK leg. (preserved in N. ITO coll.).

Distribution. Vietnam, Thailand (new record).