

quences. *J. mol. Evol.*, **42**: 124–129.

SU, Z.-H., O. TOMINAGA, M. OKAMOTO & S. OSAWA, 1998. Origin and diversification of hindwingless *Damaster* ground beetles within the Japanese Islands as deduced from mitochondrial ND5 gene sequences (Coleoptera, Carabidae). *Mol. Biol. Evol.*, **15**: 1026–1039.

Elytra, Tokyo, **28** (1): 20, May 15, 2000

A Third Record of *Curculio amabilis* MORIMOTO (Coleoptera, Curculionidae, Curculioninae)

Hiraku YOSHITAKE

Laboratory of Entomology, Tokyo University of Agriculture, Sakuragaoka 1–1–1,
Setagaya, Tokyo, 156–8502 Japan

The curculionid weevil *Curculio amabilis* was described by MORIMOTO (1962) on the basis of one male collected from Hikosan, Fukuoka Prefecture, Kyushu. This species seems to be very rare, and only one additional record has so far been known from Gifu Prefecture, Chûbu District, Honshu (NOHIRA, 1990). In the middle of May, 1999, the author obtained one male of this species by net-sweeping of *Quercus crispula* BLUME [Mizu-nara in Japanese] (Fagaceae) in Kanagawa Prefecture, Kantô District, Honshu. The collecting data will be shown below as a third record of the species.

Curculio amabilis MORIMOTO, 1962

Curculio amabilis MORIMOTO, 1962, *Mushi*, Fukuoka, **36**: 36–37.

Specimen examined. 1♂, Mt. Daigatake, Hakone, Kanagawa Pref., 17–V–1999, H. YOSHITAKE leg.

Distribution. Japan (Honshu, Kyushu).

References

- MORIMOTO, K., 1962. Revision of the subfamily Curculioninae from Japan II (Coleoptera). *Mushi, Fukuoka*, **36**: 21–45.
- NOHIRA, T., 1990. *Curculio amabilis* collected from Gifu Prefecture. *Gekkan-Mushi, Tokyo*, (237): 37–38. (In Japanese.)