

Occurrence of *Corticeus amamiensis* (Coleoptera, Tenebrionidae) in Honshu, Japan

Atsushi KATÔ¹⁾ and Isao MATOBA²⁾

¹⁾ 8-4-12-503, Kano, Higashi-osaka, Osaka, 578-0901 Japan

²⁾ Wakayama Prefectural Museum of Natural History,
Funao, Kainan, Wakayama, 642-0001 Japan

Corticeus amamiensis has hitherto been known from the Ryukyu Islands (Amami-ohshima Is., Okinawa Is., Miyako Is., Ishigaki Is., Iriomote Is. and Yonaguni Is.). We recently discovered this species from Wakayama Prefecture, Honshu, mainland of Japan. This is the northernmost record of the distribution of the species.

Corticeus amamiensis (KASZAB, 1964)

Hypophloeus (*Paraphloeus*) *amamiensis* KASZAB, 1964, p. 48 (Type locality: Amami, Kominato). — M. T. CHÛJÔ, 1982, p. 99.

Corticeus amamiensis: BREMER, 1999, p. 51.

Specimens examined. [Honshu] 2♂♂, 2♀♀, Esuzaki, Susami, Wakayama Pref., 1-VI-1996, I. MATOBA leg.

Distribution. Japan (Honshu and the Ryukyu Islands).

We are indebted to Dr. Hans J. BREMER, Melle, and Mr. Kiyoshi ANDO, Ehime University, Matsuyama, for their kindness in offering relevant literature.

References

- BREMER, H. J., 1999. Revision der orientalischen *Corticeus*-Arten (Col., Tenebrionidae, Hypophloeini). II. Teil. *Acta coleopterol., München*, **15** (1): 3-64.
- CHÛJÔ, M. T., 1982. Tenebrionidae of the Nansei Islands VII (Coleoptera). *Spec. Iss. Mem. Retir. Emer. Prof. M. Chûjô*, 87-101. (In Japanese.)
- KASZAB, Z., 1964. Über die Tenebrioniden einiger japanischer Inseln. I. (Coleoptera). *Ent. Rev. Japan*, **16**: 39-49.