A New Arrangement of the *Stenhomalus incongruus* Complex (Coleoptera, Cerambycidae)

Tatsuya Niisato

Bioindicator Co., Ltd., Yarai-chô 126, Shinjuku-ku, Tokyo, 162-0805 Japan

Stenhomalus incongruus and its relatives have so far been known from three sibling species, viz., S. incongruus Gressitt from East China, S. parallelus Niisato from the Korean Peninsula and the Tsushima Islands, and S. muneaka Hayashi from Hokkaido and Honshu. After the revisional study of these relatives by Niisato (1988), some additional records of S. muneaka and S. parallelus were made known, and slight geographical variation in the male genital organ was recognized among their local populations. It seems to me now that these allopatric "species" had better be regarded as geographical variations of the same species. In the following lines, I am going to propose the new arrangement of the subspecies under the oldest name S. incongruus Gressitt, 1939.

Stenhomalus incongruus incongruus GRESSITT

Stenhomalus incongruus Gressitt, 1939, Lingnan Sci. J., **18**, p. 211, pl. 8, fig. 3; type locality: Nanking. —— Niisato, 1988, Kontyû, Tokyo, **56**, pp. 790–791, figs. 2–3.

Distribution. East China.

Stenhomalus incongruus parallelus Niisato, stat. nov.

Stenhomalus parallelus Niisato, 1988, Kontyû, Tokyo, **56**, pp. 791–193, figs. 1, 4–5; type locality: Mine, Tsushima Isls. off northern Kyushu, SW. Japan; 1998, Elytra, Tokyo, **26**, p. 460.

Stenhomalus muneaka: Lee, 1987, Longicorn Beetles of Korean Peninsula, p. 79, pl. 10, fig. 94; locality record: Mt. Pal-Cong, Gyeong-Bug, Korea.

Distribution. South Korea and Tsushima Isls.

Stenhomalus incongruus muneaka HAYASHI, stat. nov.

Stenhomalus muneaka Hayashi, 1981, Ent. Rev. Japan, **36**, p. 29; type locality: Kuzuba-tôge Pass in Niigata Pref.; 1984, Coleopt. Japan Col., Osaka, **4**, p. 52, pl. 10, fig. 22. — Kusama & Takakuwa, 1984, Longicorn-Beetles of Japan in Color, p. 271, pl. 31, fig. 212. — Niisato, 1998, Kontyû, Tokyo, **56**, pp. 795–796, figs. 6–7; 1998, Elytra, Tokyo, **26**, p. 172; locality record, Taishakukyô, Hiroshima, W. Honshu. — Nagao, 2001, Gekkan-Mushi, Tokyo (in press); locality record: Mt. Maruyama, Sapporo-shi, Hokkaido.

Distribution. Japan (Hokkaido and Honshu).