

Occurrence of *Morimotozo shigematsui* (Coleoptera, Curculionidae) in Hiroshima Prefecture, Japan¹⁾

Hiraku YOSHITAKE²⁾ and Iwao OKAMOTO³⁾

²⁾Entomological Laboratory, Division of Bioresource and Bioenvironmental Science,
Graduate School, Kyushu University, 812-8581 Japan

³⁾Hiroshima 3-1-49, Kure, Hiroshima, 737-0101 Japan

Morimotozo shigematsui was originally described by KOJIMA and MORIMOTO (1995) as *Gryporrhynchus shigematsui* on the basis of five female specimens collected from the southern parts of the Kii Peninsula, Japan. After that, this species has never been found until now. Recently, OKAMOTO had an opportunity to collect an additional female specimen of *M. shigematsui* in Hiroshima Prefecture, Japan. To record the new locality is essential in the future for understanding the distributional range of this species. The collecting data are as shown in the following lines.

Morimotozo shigematsui (KOJIMA et MORIMOTO, 1995)

Gryporrhynchus shigematsui KOJIMA et MORIMOTO, 1995, 117-134.

Specimen examined. 1 ♀, Kure, Hiroshima, 31-VIII-1999, I. OKAMOTO leg.

References

- KOJIMA, H., & K. MORIMOTO, 1995. Systematics of the weevil genus *Gryporrhynchus* ROELOFS (Coleoptera, Curculionidae). *J. Fac. Agric. Kyushu Univ.*, **40**: 117-134.
- ALONSO-ZARAZAGA, M. A., & C. H. C. LYAL, 1999. A World Catalogue of Families and Genera of Curculionoidea (Insecta: Coleoptera). (Excepting Scolytidae and Platypodidae). 315 pp. Entomopraxis, S. C. P., Barcelona.

1) Contribution from the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka (Ser. 5, No. 55).