

## An Adult Host Record of *Oberea nigriventris* (Coleoptera, Cerambycidae)

**Yoshiyasu KUSAKABE**

Sumiregaoka 21–12, Tsuzuki-ku, Yokohama, Kanagawa, 224–0013 Japan

It has been known that the adult and larva of *Oberea nigriventris* BATES are associated with *Cynanchum caudatum* (MIQ.) MAXIM., *Marsdenia tomentosa* MORREN et DECAISNE of the Asclepiadaceae, and *Pyrus pyrifolia* NAKAI var. *culta* NAKAI of the Platanaceae. Recently, I collected many adults of this species on *Tylophora aristolochioides* MIG. (Asclepiadaceae). The adult *Oberea* fed leaves and stems of the plant, and made spawning behavior. *Tylophora aristolochioides* is the fourth host plant of the species. The collecting data are as follows:

6♂♂, 12♀♀, Mt. Takao-san, Hachiōji City, Tokyo, Japan, 10~21-VII-2000 (observations on total 5 days), Y. KUSAKABE leg.

I thank Mr. Kaoru SAKAI for his useful information of the collecting site, and Mr. Nobuyuki KOBAYASHI for determination of the plant.

### References

- KOJIMA, K., & S. NAKAMURA, 1986. Food Plants of Cerambycid Beetles (Cerambycidae, Coleoptera) in Japan. 336 pp. Hiba Society of Natural History, Hiroshima. (In Japanese with English title.)  
KUSAKABE, Y., 1992. Genus *Oberea* (part). In: OHBAYASHI, N., M. SATŌ & K. KOJIMA (eds.), *An Illustrated Guide to Identification of Longicorn Beetles of Japan*, 197–198, 646–650. Tokai Univ. Press, Tokyo. (In Japanese with English title.)  
SHIRAI, K., 1998. Distributional record of *Oberea nigriventris* BATES from Aichi Prefecture. *Gekkan-Mushi, Tokyo*, (334): 11. (In Japanese.)