

References

- BOPPE, P.-L., 1911. Un genre nouveau de Prioniens du Yunnan (Col.). *Bull. Soc. ent. Fr.*, **1911**: 29–33.
- GAHAN, C. J., 1906. Coleoptera. Cerambycidae, Vol. 1. *In the: Fauna of British India, including Ceylon and Burma*. XVIII+329 pp. Taylor & Francis, London.
- GRESSITT, J. L., 1940. The longicorn beetles of Hainan Island (Col.: Cerambycidae). *Philipp. J. Sci.*, **72**: 1–239, 8 pls.
- HÜDEPOHL, K.-E., 1997. Eine neue Art Gattung *Megopsis* SERVILLE, 1832 aus Ost-Malaysia, Sabah (Coleoptera Cerambycidae Prioninae). *Entomofauna, Ansfelden*, **18**: 45–47.
- LAMEERE, A., 1916. Trois Prioninae nouveaux (Col. Cerambycidae). *Bull. Soc. ent. Fr.*, **1916**: 233–235.
- 1919. Coleoptera Longicornia, Fam. Cerambycidae, Subfam. Prioninae. *In*: WYTSMAN, P. (ed.), *Genera Insectorum*, (172): 1+1–189, 8 pls.

Elytra, Tokyo, **29** (2): 400, November 15, 2001

New Localities of *Merionoeda tosawai* (Coleoptera, Cerambycidae)

Tatsuya NIISATO

Bioindicator Co., Ltd., Yarai-chô 126, Shinjuku-ku, Tokyo, 162–0805 Japan

Merionoeda tosawai (KOBAYASHI, 1932, Mushi, Fukuoka, **5**, p. 1, fig. 1) has so far been known to occur on two main islands of the Ogasawaras, i.e., Chichi-jima and Haha-jima Islands. I recently examined the specimens of this stenopterine collected from Imouto-jima Island off southern Haha-jima and Muko-jima Island of the Muko-jima island group as recorded below. All the specimens examined were collected by H. KARUBE of the Kanagawa Prefectural Museum of Natural History and are preserved in his museum.

1 ♂, Muko-jima Is., 27–VI–2001; 3 ♂♂, Imouto-jima Is., 26–VI–2001.

In closing this short report, I wish to thank Mr. Haruki KARUBE for his permission to examine the interesting material.