

- 319–370, pls. 16–17.
PIC, M., 1916. Diagnoses génériques et spécifiques. *Mél. exot.-ent.*, **18**: 2–20.
——— 1917. Descriptions abrégées diverses. *Ibid.*, **23**: 2–23
——— 1925. Nouveaux malacodermes asiatiques. *Bull. Mus. Hist. nat., Paris*, **31**: 72–73.
WITTMER, W., 1997. Neue Rhagophthalmidae und Phengodidae. *Mitt. schweiz. ent. Ges.*, **70**: 257–263.
——— & N. Ohba, 1994. Neue Rhagophthalmidae (Coleoptera) aus China und Benachbarten Ländern. *Jpn. J. Ent.*, **62**: 341–350.
-

Elytra, Tokyo, **29** (2): 434, November 15, 2001

Occurrence of *Lucidina biplagiata* (Coleoptera, Lampyridae) on Teuri-tô Island, off Hokkaido, Japan

Itsuro KAWASHIMA

Nagasawa 1–50–9, Yokosuka-shi, Kanagawa, 239–0842 Japan

A middle-sized lampyrine beetle, *Lucidina biplagiata* (MOTSCHULSKY, 1866), is widely distributed in the region from the Kuril Islands in the northeast to Taiwan in the southwest including Japan. I had an opportunity to examine new materials of this species through the courtesy of Mr. T. SHIMADA. In this brief report, I will record it as a new locality of this species.

Materials examined. 3♂♂, 2♀♀, Benten, Teuri-tô Is., off NW. Hokkaido, 7~8–VII–2001, T. SHIMADA leg.

I am indebted to Mr. Takashi SHIMADA of the Laboratory of Insect Resources, Tokyo University of Agriculture, Atsugi, Kanagawa, for his supplying with the materials.