

An Additional Record of *Paramimistena brevis*
(Coleoptera, Cerambycidae)

Tatsuya NIISATO¹⁾ and Hiroshi MAKIHARA²⁾

¹⁾ Bioindicator Co., Ltd., Yurai-chô 126, Shinjuku-ku, Tokyo, 185-0806 Japan.

²⁾ Forestry and Forest Products Research Institute, Ibaraki, 305-0903 Japan.

Paramimistena brevis was described based on a single male specimen collected from a primary forest on Bukit Soeharto in eastern Kalimantan. This species belongs to the group of *P. enterolobii* of the genus *Paramimistena*, and is characterized by the short broad body form, broad subbasal pale bands on the long parallel-sided elytra, and stout appendages. In the course of continuous survey in the same field, MAKIHARA succeeded in collecting an additional specimen of this very rare paramimistenine.

***Paramimistena brevis* NIISATO et MAKIHARA, 1999**

Paramimistena brevis NIISATO et MAKIHARA, 1999, *Elytra*, Tokyo, **27**, pp. 331–333, fig. 2; type locality: Bukit Soeharto, Kalimantan Timur, Indonesia.

Additional specimen examined. 1♂, Bukit Soeharto, Kalimantan Timur, Indonesia, 19–IV–1999, H. MAKIHARA leg.

Distribution. Eastern Kalimantan, Indonesia. “Western Kalimantan” recorded in the original description is a lapsus calami by the senior author, NIISATO.

Notes. The specimen examined is almost identical with the holotype, but the pronotum is a little more widened at the base, and the elytra are fairly short, 2.03 times as long as the humeral width (2.25 times in the holotype), and with pale maculations moderately arcuate at the anterior margins. The standard ratios of body parts are as follows: HW/PA 1.32, HW/PW 1.06, FL/FB 0.58, CL/CB 0.34, PL/PA 2.00, PL/PW 1.60, PB/PA 1.14, PB/EW 0.84, PL/EL 0.73, EL/EW 2.03. Body length is 4.1 mm.

The additional specimen of *P. brevis* was collected by a Malaise trap set at a height of 40 m above the ground as in the case of the holotype.