

Additional Records of *Merionoeda fusca* GERESSITT et RONDON (Coleoptera, Cerambycidae)

Tatsuya NIISATO¹⁾ and Yaheita YOKOI²⁾

¹⁾Bioindicator Co., Ltd., Nikko-Kagurazaka Building, Iwato-chō 18, Shinjuku, Tokyo, 162-0832 Japan

²⁾Beerenkotten 33, 40882 Ratingen, Germany

Merionoeda fusca is a characteristic species in having the short elytra with black external margins and largely black hind femur beyond the club part (GERESSITT & RONDON, 1970), and has so far been known from Laos and Yunnan (LÖBL & SMETANA, 2010; NIISATO & LIN, 2013). We examined the additional specimens of this species including the new record from Thailand as listed below.

We wish to express our thanks to Messrs. Katsumi AKITA and Kaoru SAKAI for their kind offer of material used in the present examination.

Merionoeda fusca GERESSITT et RONDON, 1970

Merionoeda fusca GERESSITT et RONDON, 1970: 121, fig. 22 c; type locality: Pakkading, 156 m, Borikhane Prov., Laos; Luang Prabang, 340 m, Laos.

Specimens examined. 3 ♂♂, 1 ♀, Nan Province, Thailand, VI.1992, Local collector leg.; 1 ♂, Maetang, Chiang Mai Province, Thailand, VI.1998, Local collector leg.; 1 ♂, 4 ♀♀, Mt. Phukhaokhova, 700 m, Thupakhom dist., Vientiane Province, Laos, 2-3.V.1995, K. AKITA leg.

Distribution. Laos, Thailand (new record) and China: Yunnan.

References

- GERESSITT, J. L., & J. A. RONDON, 1970. Cerambycids of Laos (Disteniidae, Prioninae, Philiinae, Aseminae, Lepturinae, Cerambycinae). *Pacific Insects Monograph*, 24: 1-314.
LÖBL, I., & A. SMETANA, 2010. Catalogue of Palaearctic Coleoptera, 6. Chrysomeloidea. 924 pp. Apollo Book, Stenstrup.
NIISATO, T., & M.-Y. LIN., 2013. Collection list of the genus *Merionoeda* (Coleoptera: Cerambycidae: Cerambycinae: Stenopterini) preserved in the Institute of Zoology, Chinese Academy of Sciences, Beijing. Pp. 69-83. In LIN, M.-Y., & C.-C. CHEN (eds.), *In Memory of Mr. Wenshin LIN*. 233 pp. Formosa Ecological Company, Taiwan.

Manuscript received 7 November 2015;
revised and accepted 8 November 2015.