

## Occurrence of *Merus (Merus) piceus* (ROELOFS) (Coleoptera, Curculionidae, Molytinae) on Okinoerabujima Is., the Amami Isls., Southwestern Japan

Hiraku YOSHITAKE

Institute for Agro-Environmental Sciences, NARO, 3–1–3 Kannondai, Tsukuba, Ibaraki, 305–8604 Japan

*Merus (Merus) piceus* (ROELOFS, 1875), belonging to the subfamily Molytinae, family Curculionidae, is known to be a gall-maker on oak, chinquapin, and chestnut trees (MORIMOTO, 1984) and distributed in China (Fujian, Hubei & Zhejiang), Japan, South Korea, and Taiwan (ALONSO-ZARAZAGA, 2013). In Japan, this species has hitherto been recorded from the Izu Isls. (Mikurajima Is. & Hachijojima Is.), Honshu, Sadogashima Is., the Oki Isls. (Dôgo Is.), Shikoku, Kyushu, and the Ryukyus (Okinawajima Is.) (KADOWAKI, 2001; ÔGAI, 2014; KOJIMA, 2015). As far as I know, however, *M. (M.) piceus* has never been recorded from the Amami Isls. located north of Okinawajima Is. In this short report, I record this species for the first time from Okinoerabujima Is., the Amami Isls., filling a gap in its known distribution. I thank M. HAYASHI, H. KOJIMA, and N. TSUJI for their help with literature. I also thank H. SUENAGA for his donation of specimens.

### *Merus (Merus) piceus* (ROELOFS, 1875)

*Alcides piceus* ROELOFS, 1875, 152 (type locality: “Japon”).

*Mecyslobus* [sic!] *piceus*: MORIMOTO, 1962, 347 (catalogued).

*Mecysolobus piceus*: MORIMOTO, 1984, 288, pl. 56, fig. 23 (diagnosis, distribution & ecology).

*Specimens examined.* Japan: the Ryukyus. 1 ex., Amami Isls., Okinoerabujima Is., China-chô, Mt. Ooyama, 6.VII.2015, H. SUENAGA leg.; 2 exs., same locality, 21.V.2016, H. YOSHITAKE leg.

*Distribution.* Japan: Izu Isls. (Mikurajima Is. & Hachijojima Is.), Honshu, Sadogashima Is., the Oki Isls. (Dôgo Is.), Shikoku, Kyushu, and the Ryukyus (Okinawajima Is. & Okinoerabujima Is. — new record); China; South Korea; and Taiwan.

### References

- ALONSO-ZARAZAGA, M. A., 2013. Molytinae. Pp. 475–497. In LÖBL, I., & A. SMETANA (eds.), *Catalogue of Palaearctic Coleoptera*, 8. 700 pp. Brill, Leiden/Boston.
- KADOWAKI, H., 2001. [Beetles collected in the Oki Isls. and identified by Dr. Takehiko NAKANE.] *Sukashiba, Izumo*, (49): 39–55. (In Japanese.)
- KOJIMA, H., 2015. A preliminary checklist of the superfamily Curculionoidea (Insecta, Coleoptera) (excepting Scolytidae and Platypodidae) from Mikura-jima Is., the Izu Islands, Japan. *Mikurensis*, 4: 17–44.
- MORIMOTO, K., 1962. Preliminary check list of the families Curculionidae (II) and Rhynchophoridae of Japan. *Science Bulletin of the Faculty of Agriculture, Kyushu University*, 19: 341–368.
- MORIMOTO, K., 1984. Curculionidae. Pp. 269–345 [incl. pls. 53–68]. In HAYASHI, M., K. MORIMOTO & S. KIMOTO (eds.), *The Coleoptera of Japan in Color*, 4. 438 pp. Hoikusha, Osaka. (In Japanese, with English book title.)
- ÔGAI, H., 2014. [On beetles new to the fauna of Okinawa-Hontô Is. (2).] *Gekkan-Mushi, Tokyo*, (524): 52–53. (In Japanese.)
- ROELOFS, W., 1875. Curculionides recueillis au Japon par M. G. LEWIS. *Annales de la Société entomologique de Belgique*, 18: 149–194.

Manuscript received 17 November 2017;  
revised and accepted 21 November 2017.