A New Record of *Procirrus lewisii* SHARP (Coleoptera, Staphylinidae, Paederinae) from Tsushima Island, Japan

Yoshihiro Senda


c/o Hiwa Museum of Natural Science, Hiwa 1119–1, Hiwa, Shôbara City, Hiroshima, 727–0301 Japan E-mail: geostix666@gmail.com

Procirrus lewisii was originally described from Kyushu, Japan (Sharp, 1889). Since then, this species has been recorded also from Hong Kong and Thailand (Herman, 2010). In Japan, this species seems to be rare, but is known to be widely distributed in Honshu, the Izu Islands (Izu-Ôshima Island), the Ogasawara Islands (Haha-jima Island), Shikoku, Kyushu, and the Ryukyus (the islands of Kuchinoerabu-jima, Tokunoshima, and Ishigaki-jima) (KISHIMOTO, 1995; WATANABE, 2001; YAMAJI, 2005; KAMEZAWA & NOMURA, 2013). Recently, I examined a specimen of this species collected from Tsushima Island which is located in the Korea / Tsushima Strait, approximately halfway between the Japanese mainland and the Korean Peninsula. Here I newly record it as below. I thank Dr. Hiroyuki YOSHITOMI (Ehime University Museum) and Mr. Osamu YAMAJI (Okayama City) for their kind helps in various ways.

Procirrus lewisii Sharp, 1889

(Figs. 1 & 2)

Specimen examined. 1 ex., Kechi, Mitsushima-chô, Tsushima Is., Japan, 14.VII.2000, T. Kan leg. (Fig. 1) Distribution. Japan: Honshu, Izu Isls. (Izu-Ôshima Is.), Ogasawara Isls. (Haha-jima Is.), Shikoku, Kyushu, and the Ryukyus (Kuchinoerabu-jima, Tokunoshima, and Ishigaki-jima Iss.); Tsushima Is. — new record (Fig. 2). Note. The specimen is preserved in the Ehime University Museum, Matsuyama, Japan.



Figs. 1–2. *Procirrus lewisii* —— 1, Dorsal habitus, specimen collected from Tsushima Is., Japan; 2, distribution in Japan (circle: previous records; star: the present new record).

36 Yoshihiro Senda

References

- HERMAN, L. H., 2010. Generic revision of the Procirrina (Coleoptera: Staphylinidae: Paederinae: Pinophilini). *Bulletin of the American Museum of Natural History*, New York, **347**: 1–78.
- KAMEZAWA, H., & S. NOMURA, 2013. [New records of two species of Pinophilini, Paederinae from Izu-Ôshima Is.] *Sayabane*, (n. ser.), *Tokyo*, (10): 25–26. (In Japanese.)
- KISHIMOTO, T., 1995. New record of *Procirrus lewisii* (Coleoptera, Staphylinidae, Paederinae) from Ishigaki-jima of the Ryukyu Islands, Southwest Japan. *Elytra*, *Tokyo*, **23**: 262.
- SHARP, D., 1889. The Staphylinidae of Japan. Annals and Magazine of Natural History, London, (6) 3: 319-334.
- Yamaji, O., 2005. [Records of the beetles collected from Okayama Prefecture.] Suzumushi, Kurashiki, (140): 79–74. (In Japanese.)
- WATANABE, Y., 2001. Occurrence of *Procirrus lewisii* Sharp (Coleoptera, Staphylinidae) on Haha-jima Island of the Ogasawara Islands. *Elytra*, *Tokyo*, **29**: 118.

Manuscript received 5 January 2019; revised and accepted 3 March 2019.