

A New Record of *Longitarsus holsaticus* (LINNAEUS) (Coleoptera, Chrysomelidae, Galerucinae, Alticini) from the Okinawa Isls., Central Ryukyus, Southwestern Japan

Hiroaki SHIGETO¹⁾ and Hiraku YOSHITAKE²⁾

¹⁾Naha Plant Protection Station, Ministry of Agriculture, Forestry, and Fisheries of Japan,
2–11–1 Minatomachi, Naha, Okinawa, 900–0001 Japan

²⁾Kyushu Okinawa Agricultural Research Center (Itoman residence),
NARO, 820 Makabe, Itoman, Okinawa, 901–0336 Japan

Longitarsus holsaticus was described from Europe by LINNAEUS (1758) as *Chrysomela holsatica*, and has so far been known from Europe, Russia, China, South Korea, and Japan (Hokkaido to Amami-Ōshima Is. in the central Ryukyus). In Japan, this species was known as *L. haemorrhoidalis* JACOBY, 1885, which was described from Yokohama, Honshu, but WARCHALOWSKI (1969) synonymized it with *L. holsaticus* based on Korean specimens. The second author collected this species from Izena-jima Is. in the Okinawa Isls., central Ryukyus, southwestern Japan.

In this short paper, we report the southernmost record of the species. Scientific and Japanese names follow KIMOTO & TAKIZAWA (1994). We thank Mr. Haruki SUENAGA (Kurashiki) for his constructive comments on the manuscript.

Longitarsus holsaticus (LINNAEUS, 1758)

[Japanese name: Inunofuguri-tobi-hamushi]

Specimens examined. Japan: the Ryukyus. [Okinawa Isls.] 4 exs., Okinawa-ken, Izena-mura, Uchihana, Sakata, 6.III.2020, H. YOSHITAKE leg.

Distribution. Japan: Hokkaido, Okushiri-tō Is., Honshu, the Izu Isls. (Miyake-jima Is. and Hachijō-jima Is.), Oki Isls. (Dōgo Is.), Shikoku, Kyushu, Tsushima Is., the Ōsumi Isls. (Tanegashima Is.), and the Ryukyus (Nakanoshima Is., Amami-Ōshima Is., and Izena-jima Is.); Europe, Russia, China, and South Korea (DÖBERL, 2010; NAKANE & KIMOTO, 1961; OHNO, 1968).

Notes. New to Izena-jima Is. This is the first record of this species from the Okinawa Isls., representing extension of the southernmost limit in its distribution range. The examined specimens were collected by light trap.

References

- DÖBERL, M., 2010. Alticinae. Pp. 491–563. In LÖBL, I., & A. SMETANA (eds.), *Catalogue of Palaearctic Coleoptera*, **6**. 924 pp. Apollo Books, Stenstrup.
- KIMOTO, S., & H. TAKIZAWA, 1994. Leaf Beetles (Chrysomelidae) of Japan. 539 pp. Tokai University Press, Tokyo. (In Japanese, with English title, keys & synonym list.)
- LINNAEUS, C., 1758. *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymies, Locis*. Tomus I. Editio decima, reformata. 824 pp. Laurentii Salvii, Holmiae.
- NAKANE, T., & S. KIMOTO, 1961. Entomological results from the scientific survey of the Tokara Islands. Coleoptera: Chrysomelidae. *Bulletin of the Osaka Museum of Natural History, Osaka*, (13): 71–79.
- OHNO, M., 1968. A revision of *Longitarsus*-species occurring in Japan (Coleoptera, Chrysomelidae, Alticinae). *Journal of the Toyo University, General Education, Natural Science, Tokyo*, (9): 1–56.
- WARCHALOWSKI, A., 1969. Beitrag zur Kenntnis der koreanischen Halticinen (Coleoptera, Chrysomelidae). *Annales Zoologici, Warszawa*, **27**: 225–236.