

A New Distributional Record of *Cybister sugillatus* ERICHSON, 1834 (Coleoptera, Dytiscidae) from Kita-daitô Island, Japan

Tai KOMORIYA

College of Biological Sciences, University of Tsukuba, Ibaraki, 305–8572 Japan
E-mail: kmry.inslrs@gmail.com

Cybister sugillatus ERICHSON, 1834 (Dytiscidae; Tobiiro-gengorô in Japanese) is widely distributed from southeastern Asia through Sri Lanka, Nepal, the Philippines, Taiwan, China, and Japan (MORI & KITAYAMA, 2002). In the Daitô Islands, this species was known only from Minami-daitô Island (MATSUI, 1990).

In this short paper, I recorded *C. sugillatus* for the first time from Kita-daitô Island.

Before going further, I thank Mr. Ryousuke MATSUSHIMA for his kind help in the course of this study. All examined specimens are kept in my personal collection.

Specimens examined. 2 ♀♀, Nakano, Kita-daitô-son, Kita-daitô Island, Okinawa Prefecture, Japan, 25°56'3"N, 131°18'1"E, 20.I.2020, T. KOMORIYA leg.

Distribution. Japan: the Ryukyus (Nakanoshima Island in the Tokara Islands and southward).

Notes. New to Kita-daitô Island. These specimens were collected from a pond bushed by aquatic plants, together with *Hydrophilus bilineatus cashimirensis* REDTENBACHER, 1844 (Hydrophilidae; Kogata-gamushi in Japanese), *Laccophilus sharpi* RÉGIMBART, 1889 (Dytiscidae; Sharp-tsubugengorô in Japanese) and *Canthhydrus nitidulus* SHARP, 1882 (Noteridae; Tsuya-kotsubugengorô in Japanese).

References

MORI, M., & A. KITAYAMA, 2002. Dytiscoidea of Japan (Revised edition). 232 pp. Bun-ichi Sogo Shuppan, Tokyo. (In Japanese.)
MATSUI, E., 1990. [Collecting records of water beetles from the Ryukyus.] *Kitakyushu-no-Konchu*, **37** (2): 69–76, pls. 9–10 (In Japanese.)

Manuscript received 5 July 2020;
revised and accepted 27 July 2020.