May 25, 2016

## An Additional Record of *Merionoeda kinoshitai* (Coleoptera, Cerambycidae)

## Tatsuva Niisato<sup>1)</sup> and Yaheita Yokoi<sup>2)</sup>

<sup>1)</sup> Bioindicator Co., Ltd., Nikkô-Kagurazaka Building, Iwato-chô 18, Shinjuku, Tokyo, 162–0832 Japan <sup>2)</sup> Beerenkothen 33, 40882 Ratingen, Germany

Merionoeda kinoshitai Yokoi et Niisato, 2009 was described solely on account of the holotype male from "Chiang Mai District" without details in collection data (Yokoi & Niisato, 2009). No additional specimen of the species were found thereafter, though expected from somewhere in northern Thailand or its vicinities. Recently we had an opportunity to examine the Merionoeda specimens in Masamichi Yagi collection preserved in the Osaka Museum of Natural History, and identified one specimen as a male of M. kinoshitai. We would like to thank Mr. Shigehiko Shiyake for his permission to examine the valuable specimen in the museum collection.

## Merionoeda kinoshitai Yokoi et Niisato, 2009

(Figs. 1-3)

Merionoeda kinoshitai Yokoi et Niisato, 2009: 329, figs. 3, 10-15; type locality: Chiang Mai District.

Specimen examined. 1  $\circlearrowleft$ , "Chiengmai / Thailand / V. 1988 / N. Koyama leg." "12–4 (yellow card)" "Masamichi YAGI / collection / [OMNH 07-14]".


Distribution. Thailand.

*Notes*. The specimen examined agrees well with the holotype of *M. kinoshitai*, though the body size is a little smaller (7.5 mm; 8.1 mm in the holotype). Measurements of other body parts are as follows (in mm): head width across eyes 1.53; length of pronotum 1.38; maximum width of pronotum across lateral swellings 1.22; apical width of pronotum 0.95; basal width of pronotum 1.20; length of elytra 3.55; humeral width of elytra 1.85.

## Reference

Yokoi, Y., & T. Niisato, 2009. Three New *Merionoeda* (Coleoptera, Cerambycidae) from northern Indochina. *Elytra*, *Tokyo*, 37: 343–353.

Manuscript received 30 April 2016; revised and accepted 8 May 2016.



Figs. 1–3. *Merionoeda kinoshitai* YOKOI et NIISATO, 2009,  $\delta$ , from Chiang Mai, northern Thailand. —— 1, Habitus; 2, pronotum (SEM image); 3, abdominal sternites 7–8 and apical part of median lobe (SEM image).