

Occurrence of *Ceutorhynchoides squamatus* YOSHITAKE et COLONNELLI (Coleoptera, Curculionidae) on Okinawajima Island, the Ryukyus, Japan

Hiraku YOSHITAKE

Natural Resources Inventory Center, National Institute for Agro-Environmental Sciences,
Kannondai 3-1-3, Tsukuba, Ibaraki, 305-8604 Japan

Ceutorhynchoides squamatus YOSHITAKE et COLONNELLI, 2005 belonging to the tribe Ceutorhynchini, the subfamily Ceutorhynchinae was described on the basis of the type series collected from Iriomotejima Is. and Ishigakijima Is., the Yaeyama Isls., the Ryukyus. No information on its host plant association is available, except that several *C. squamatus* adults were obtained from *Styrax japonica* SIEBOLD et ZUCC. (Styracaceae) on Ishigakijima Is. (I. MATOBA, pers. comm., 2011). As far as I know, no additional record of this species has been reported after the original description. Recently, I had an opportunity to examine a specimen of *C. squamatus* collected for the first time on Okinawajima Is., as recorded below.

Ceutorhynchoides squamatus YOSHITAKE et COLONNELLI, 2005

Ceutorhynchoides squamatus YOSHITAKE et COLONNELLI, 2005, 121.

Specimen examined. 1 ♂, Mt. Tanodake, Nago, Okinawajima Is., Ryukyus, 21-III-2010, Takashi & Toshiko KURIHARA leg.

Distribution. Japan (Ryukyus: Okinawajima Is., Ishigakijima Is., Iriomotejima Is.).

I thank T. KURIHARA (Tochigi Prefectural Museum, Utsunomiya) for the donation of the specimen examined and I. MATOBA (Wakayama Prefectural Museum of Natural History, Kainan) for the kind offer of biological information.

Reference

YOSHITAKE, H., & E. COLONNELLI, E., 2005. Taxonomic study of the genus *Ceutorhynchoides* COLONNELLI, 1979 from Japan, with establishment of an allied new genus (Coleoptera: Curculionidae: Ceutorhynchinae). *Esakia, Fukuoka*, (45): 117-153.

Manuscript received 7 March 2011;
revised and accepted 26 April 2011.