

On Several Species of Histerids from the Island of Hokkaido, Japan (Coleoptera, Histeridae)

Masahiro ÔHARA¹⁾ and Hiroshi SHIDO²⁾

¹⁾ Hokkaido University Museum, Hokkaido University, North 10, West 8,
Sapporo, 060–0810 Japan

²⁾ 10–20, 11 jō, 16 chōme, Shin-kotoni, Sapporo, 001–0911 Japan

On the island of Hokkaido, a total of 60 species of histerid beetles has been recorded so far (ÔHARA, 1994, 1999). We have an opportunity to examine 22 specimens of histerid beetles in Hokkaido collected by the second author. Of these, several species newly recorded from the local districts of Hokkaido are treated in this paper.

1. *Onthophilus flavigornis* LEWIS, 1878
Ishikari: 1 ex., Maruyama-kōen, Sapporo, 29–VI–2005.
The last record of this species from the same locality is at 1939 (ÔHARA & NAKANE, 1986).
2. *Hololepta depress* LEWIS, 1884
Iburi: 1 ex., Shikotsu-ko, Bifue-gawa, Chitose, 22–V–2005. New to Iburi district.
3. *Platylistes (Platylistes) pini* (LEWIS, 1884)
Ishikari: 4 exs., Maruyama-kōen, Sapporo, 18–VII–1999, 26–VI–2001, 21–VI–2002, 15–VII–2005. New to Ishikari district.
4. *Margarinotus (Ptomister) yezoensis* M. ÔHARA, 1989
Ishikari: 1 ex., Maruyama-kōen, Sapporo, 27–VI–2001.
5. *Margarinotus (Ptomister) weymarni* WEZNEL, 1944
Ishikari: 1 ex., Shin-kotoni, Sapporo, 9–VI–2009.
6. *Margarinotus (Ptomister) agnatus* (LEWIS, 1884)
Hiyama: 1 ex., Dobashi-shizen-kōen, Assabu, 12 to 16–VIII–2001. New to Hiyama district.
7. *Margarinotus (Grammostethus) niponicus* (LEWIS, 1885)
Ishikari: 2 exs., Maruyama-kōen, Sapporo, 19–VII–1999, 6–VIII–2005.
8. *Atholus duodecimsteriatus quatuordecimstriatus* (GYLLENHAL, 1808)
Iburi: 1 ex., Morino, Shiraoi-gawa, Shiraoi, 9–V–2000.
9. *Atholus pirithous* (MARSEUL, 1873)
Ishikari: 1 ex., Shiratsukari-hama, Ishikari (old name: Atsuta), 8–V–2001.
10. *Merohister aino* (LEWIS, 1884)
Ishikari: 1 ex., Maruyama-kōen, Sapporo, 10–VII–1999.
11. *Hetaerius optatus* LEWIS, 1884
Ishikari: 1 ex., Maruyama-kōen, Sapporo, 3–VI–2005.
12. *Dendrophilus xavieri* MARSEUL, 1873
Iburi: 1 ex., Hobetsu Sky Slope, Hobetsu, Yūfutsu, 6–VII–2002. New to Iburi district.
13. *Anapleus semen* (LEWIS, 1884)
Hiyama: 1 ex., Yunotai, Kaminokuni, 20–V–1995. New to Hiyama district.
14. *Hypocaccus (Hypocaccus) subaeneus* (SCHMIDT, 1890)
Iburi: 4 exs., Shikotsu-ko, Bibi-gawa, Chitose, 2–V–1992; 2 exs., Morino, Shiraoi-gawa, Shiraoi, 4–V–2000, 4–V–2009. New to Iburi district.
15. *Saprinus niponicus* DAHLGREN, 1962
Ishikari: 1 ex., Maruyama-kōen, Sapporo, 13–VII–2008.

References

- ÔHARA, M., 1994. A revision of the superfamily Histeroidea of Japan (Coleoptera). *Ins. matsum.*, New series, (51): 1–283.
_____, 1999. A revision of the superfamily Histeroidea of Japan (Coleoptera), Supplementum 1. *Ibid.*, (55): 75–122.
_____, & T. NAKANE, 1986. On the genus *Onthophilus* from Japan (Coleoptera, Histeridae). *Ibid.*, (37): 1–15.

Manuscript received 22 March 2012;
revised and accepted 22 April 2012.